

# Phytochemical Screening and Antimicrobial Activities of *Lepidium sativium* and *Rumex nervosus* in Eritrea

## ABSTRACT

Some of the widely used plants for the treatment of eye diseases, especially animals, include *Lepidium sativium* and *Rumex nervosus*. Therefore, the phytochemical screening and antimicrobial activities of the crude extracts of the seeds *L. sativium* and leaves of *R. nervosus* were investigated. Freshly matured seeds of *L. sativium* and leaves of *R. nervosus* were collected, and sequentially extracted by soxhlet extractor using different polarity solvents including n-hexane, methanol, ethanol and water. The extracts were thus screened for the presence of the prominent secondary metabolites which display characteristic bioactivities. Comparatively, the methanol and ethanol extracts displayed most of the metabolites including alkaloids, flavonoids, tannins, phenols, glycosides and steroids. Moreover, the antimicrobial activities of the extracts were tested against *Staphylococcus aureus*, *Escherchia coli* and *Candida albicans* using agar well diffusion method. The ethanolic extracts of *R. nervosus* leaves showed the highest activity (18 mm) against *S. aureus* and the water extracts gave the lowest zone of inhibition (6 mm) against *E. coli*. In all the extracts, the growths of the two bacterial and fungus strains were inhibited by the methanol and ethanol extracts. The best antifungal activity was observed in the methanol extract of the seeds of *L. sativium* against *Candida albicans* (20 mm). The results revealed that, the ethanol and methanol extracts of both plants were the two best extractive solvents with potential inhibitory activity against microbial growth. The observed activities were related to the presence of the noticeable phytochemicals in those plants. Therefore, this finding strongly supports the claim of the local community to use *L. sativium* and *R. nervosus* for the treatment of different pathogenic bacterial infections associated to eye diseases. However, the traditional aqueous extraction practice should be modified by addition of alcohol in order to maximize the extraction efficiency and thus the bioactivity of the plants.

Key words: *Lepidium sativium*, *Rumex nervosus*, phytochemicals, antimicrobial, eye disease

## 1. INTRODUCTION

Infectious diseases remain the most common cause of several illness and death worldwide, the causative agents of infectious include bacteria, viruses, fungi and parasites (Amit *et al.*, 2014). Medicinal plants have source for the invention of novel drugs and 25 % of modern drugs contain one or more active principles of plant origin (Abiyu *et al.*, 2014). According to WHO, medicinal plants would be the best source to obtain a variety of drugs. Therefore, medicinal plants should be investigated to better understand their properties, safety and efficacy (Bruno, 2013). Many plants have been used because of their antimicrobial traits, which are due to compounds synthesized in the secondary metabolism of the plant. These products are known by their active substances, for example, phenolic compounds, which are part of essential oils, tannins, terpenoids, alkaloids, and flavonoids. These metabolites have been found *in vitro* to have antimicrobial properties.

Many plants have been evaluated not only for their inherent antimicrobial activity, but also for their action as a resistance-modifying agent. The enhancement of antibiotic activity or the reversal of antibiotic resistance by natural or synthetic nonconventional antibiotics has led to the classification of these compounds as modifiers of antibiotic activity. The antimicrobial compounds produced by plants are active against plant and human pathogenic microorganisms (Hero and Jwan, 2012). Herbal molecules are safe and would overcome resistance produced by the pathogens as they exist in a combined form or in a pooled form of more than one molecule in the protoplasm of the plant cell (Endalkachew and Negasse, 2016).

Eye disease is a serious problem in Eritrea especially for animals due to the bacterial infection, therefore there is a need to identify the microbes as well as develop cheap and less hazardous drugs from medicinal plant to minimize this bacterial infection. With novel mechanism of action that is less susceptible to resistance by microbes, eye diseases in remote areas of Eritrea are still traditionally cured by medicinal plants. Based on some ethnobotanical information, the treatment of eye disease in Eritrea, especially in the treatment of animals, includes the use of plants such as *Barleria eranthemoides*, *Lepidium sativium*, *Cynoglossum lanceolatum*, *Zehneria scarab*, *Ocimum hadiense* and *Rumex nervosus*.



**Figure 1:** *Lepidium sativium* (left) and *Rumex nervosus* (right)

The seed of *Lepidium sativium* is used to treat backache, tympanitis pain and nervous disorders (Rehab and Salem, 2013). The various parts of the plant have been used for the treatment of jaundice, liver problems, diarrhea, dysentery spleen diseases, gastrointestinal disorders, menstrual problems, fracture, arthritis, and other inflammatory conditions (Haider, 2016; Indumathy and Ajithadus, 2013). The seeds boiled in milk are used for nursing mothers to increase the secretion of milk and relieving the flatulence (Asra *et al.*, 2017). *Lepidium sativium* is largely recommended by traditional herbal healers for diabetes control, hypertension, renal disease and physiotherapy (Mussie *et al.*, 2015; Shahla *et al.*, 2014). The roots and aerial parts of *Rumex nervosus* have been used traditionally for a variety of therapeutic uses, such as antioxidant, cytotoxic, antifertility, anti-inflammatory, antimicrobial, antidiarrheal and antiviral activities (Rao *et al.*, 2011). The juice of *Rumex nervosus* is used in Ethiopia to seizure bleeding during male circumcision (Zewdu, 2013). Traditionally in Eritrea, the leaves, stems and sometimes roots of *Rumex nervosus* are used as traditional medicines, for the eye disease, taeniocapitis, haemorrhoids, infected wounds, arthritis, eczema, abscess and gynecological disorders. In this report, the seeds of *Lepidium sativium* and leaves of *Rumex nervosus* were selected based on their wider applications and thus the phytochemical and antimicrobial activities of the plants were investigated.

## 2. MATERIALS AND METHODS

### 2.1. Plant Collection and Sampling

Seeds of *Lepidium sativium* were purchased from a market in Asmara and the fresh matured leaves of *Rumex nervosus* were collected from Adi-hawisha, Zoba Maekel, Eritrea during the month of February of 2018. The plant species were identified and authenticated by a Botanist in the Department of Biology of EIT, Eritrea. The plants were thoroughly washed in running tap water to remove debris and dust particles and then rinsed using distilled water and then finally air dried for 14 days under shade. The shade dried plant materials were powdered using pestle and mortar and thus kept in polythene bags until further use.

### 2.2. Preparation of Plant Extracts

The powdered plants (200 g) were extracted sequentially using a soxhlet apparatus with with n-hexane (98%), ethanol (90%), methanol (99%) for 2 hrs each and the final residue was extracted with distilled water for 6 hrs. All the extracts were concentrated using Buchi Rota evaporator under reduced pressure and the water bath was fixed at 40 °C. The crude extracts obtained were stored in sterile capped reagent bottles and refrigerated at 4 °C until further use. From the extracts, 0.5 g sample was reconstituted in the

respective solvents (in 20 mL) to get a final concentration 25 mg/mL and thus was used for the phytochemical screening. Moreover, each crude extract (0.25 g) was dissolved in 1 mL DMSO to give 250 mg/mL solution for antimicrobial assay (Shama *et al.*, 2014).

### 2.3. Phytochemical Screening

The crude extracts of the two plants were tested for the presence different phytochemicals based on previously reported methods (Trease *et. al*, 1989; Harborne, 1998). The common types of reactions including Salkowski test for terpenoids and steroids, alkaline reagent test for flavonoids, ferric chloride test for phenols, Wagner's reagent for alkaloids, foam test for saponins, Keller-Killian's test for glycosides and lead acetate test for tannins were employed.

### 2.4. Antimicrobial Screening

#### 2.4.1. Chemicals and Materials

The materials and chemicals used include McFarland solution, saline water, incubator, safety cabinet, an autoclave for sterilizing, Mueller-Hinton agar, Saboradi agar, 6 mm diameter well borer, sterile micro syringe, and DMSO.

#### 2.4.2. Media Preparation

The mixture of Mueller-Hinton agar powder and sterile distilled water were stirred with a sterilized glass rod and covered with a cotton wool, over which an aluminum foil was tightly wrapped and then autoclaved for 15 min at 121 °C. Soon after autoclaving, the agar was allowed to cool and placed inside a water bath at about 50 °C to maintain the media in a molten stage in order to minimize the amount of condensation that forms. Then, the agar medium was allowed to cool to room temperature in the laminar flow hood prior to pouring it into the Petri plate. Plates were dried faster in lower humidity by keeping them in a laminar flow hood. The freshly prepared and cooled medium was poured into flat-bottomed petri-dishes on a level, horizontal surface to give a uniform depth of approximately 4 mm (Aragaw and Nega, 2016).

#### 2.4.3. Preparation of Bacterial Suspension

Using a sterile inoculating loop, one or two colonies of the organism to be tested were taken from the subculture plate. The organism was suspended in 3 mL of physiological saline. The test tube containing the saline was then vortexed to create an overall smooth suspension. McFarland standards are used as reference to adjust the turbidity of any given bacterial suspension. McFarland solution is an essential material needed before testing the microorganisms for their sensitivity. This was done to make sure that the number of bacteria is within a given range to standardize the microbial testing. This would also help avoid any error in result, because if the suspension is too heavy or too diluted, an error might occur for any given antimicrobial agent (Nishat, 2017).

#### 3.4.4. Well Diffusion Method

Optimally, within 15 minutes after adjusting the turbidity of the inoculum suspension, a small volume about 0.1 ml of the bacterial suspension was inoculated onto the dried surface of Mueller-Hinton agar plate and Saboradi agar of the fungal suspension and streaked by the sterile cotton swab over the entire sterile agar surface. This procedure was repeated by streaking two more times, rotating the plate approximately 60 °C each time to ensure an even distribution of inoculum and finally the rim of the agar was swabbed. The lid was left ajar for 3 to 5 minutes, but no more than 15 minutes, to allow for any excess surface moisture to be absorbed before applying the crude extracts on the respective well (Aragaw and Nega, 2016).

#### 3.4.5. Test Bacterial Organisms

Two species of bacteria, *Staphylococcus aureus* (gram positive, ATCC 25923) and *Escherichia coli* (gram negative, ATCC 25922) and *Candida albicans* (fungus, ATCC 10231) were used as indicator microorganisms to detect the anti-microbial activities. The bacterial strains were selected due to their effect associated with eye infections. The analysis was done in the microbiology laboratory, National Health Laboratory, Asmara, Eritrea. All the strains were obtained as actively growing cultures from the microbial culture collection of the microbiology laboratory.

## 4. RESULTS AND DISCUSSIONS

The phytochemicals including triterpenoids, steroids, flavonoids, phenols, alkaloids, saponins, glycosides, tannins were investigated. As presented in Table 1, most of the phytochemicals were present in large or small amount in most of the extracting solvents. Most of the phytochemicals were present in the ethanol and methanol extracts of both plants. Steroids and glycosides were not present in the hexane extracts of both plants and this could be due to the non-polar nature of the solvent. Alkaloids, flavonoids and phenols which are polar compounds were found in almost the solvents but not appreciably much in the n-hexane extracts. Moreover, tannins and phenols were found in the water extracts of both plants. These results revealed that these plants have quite a number of chemical constituents, which may be responsible for the different pharmacological actions related to those plants and the existing secondary metabolites promote characteristic antimicrobial activities. The results infer that the two plants are rich in potentially bioactive metabolite and thus their use as herbal medicine by the public can be demonstrated. Moreover, the phytochemicals present in the plants can serve as a valuable source of information and provide appropriate standards to establish a base for identification and elucidation of the different types of bioactive chemicals using advanced spectroscopic methods.

**Table 1:** Qualitative Phytochemical Analysis of the Crude Plant Extracts

Medicinal plant	Solvent	Phytochemicals							
		Alkaloids	Terpenoids	Steroids	Flavonoids	Tannins	Saponins	Phenols	Glycosides
<i>Lepidium sativum</i>	n-hexane	+	-	-	+	-	+	-	-
	Methanol	+	+	+	+	+	+	+	+
	Ethanol	+	+	+	+	+	+	+	+
	Water	+	+	+	+	-	+	+	-
<i>Rumex nervosus</i>	n-hexane	+	+	-	-	-	-	+	-
	Methanol	+	+	+	+	+	+	+	+
	Ethanol	+	-	+	+	+	+	+	+
	Water	-	+	-	+	+	-	+	+

Key: '+'- Present, '-' Absent of the particular compound

Previously reported phytochemicals found in *L. sativum* showed that the plant contains triterpenes, alkaloids, flavonoids, tannins and coumarins but not saponins (Nuha 2006; Haider, 2016). The seeds show the presence of volatile essential aromatic oils (Hero and Jwan, 2012) carbohydrates, phenolic compounds, flavonoids, alkaloids, proteins, saponins and lipids. It also contains good amount of lignans and antioxidants. The plant is known to contain imidazole, lepidine,  $\beta$ -carotenes, ascorbic acid, linoleic acid, oleic acid, palmitic acid and stearic acid. *L. sativum* contains significant amount of iron, calcium, folic acid, beside vitamins A and C (Abdel *et al.*, 2017). Moreover, recent studies showed that the leaves of the *R. nervosus* contain palmitoleic acid and palmitic acid (Mohammed *et al.*, 2018).

### 4.2. Antimicrobial Activities of the Crude Extracts


The antimicrobial activity of the extracts was determined by growth inhibition activities of the extracts and thus the diameter of the zone of inhibition exhibited by the extracts against the different microbes was measured. As shown in Table 2 and Figure 2, the methanol and ethanol extracts of *L. sativum* exhibited good inhibition activities against all the pathogens. The methanol extract displayed the highest zone of inhibition (20 mm) followed by the ethanol extract (16 mm) against *C. albicans*. However, the ethanol extract displayed better activity against *S. aureus* and *E. coli*, with zones of inhibition of 15 and 13 mm respectively, compared to the other extracts. Although the n-hexane extracts displayed a low zone of inhibition (8 mm) against both *S. aureus* and *C. albicans* but there was no measurable activity against *E. coli*. Similarly, the aqueous extracts indicated some activities against *E. coli* (10 mm) and *S. aureus* (8 mm) but no characteristic activity was observed against *C. albicans*. As previously reported the petroleum ether extract of seeds of *L. sativum* showed antimicrobial activity against some standard human pathogens. The antimicrobial activity of the water extract was low against *Bacillus subtilis* and *Escherchia coli*. The extract had no effect against *S. aureus* and *C. albicans* (Afaf *et. al.*, 2015). All the extracts had lower activities relative to the standard drug. Though there are other reported bioactivities

related to *L. sativum*, the type of solvents used, the extraction process and the stains employed were different (Pragya *et. al.* 2011; Pragya *et. al.* 2012; Shahla *et. al.* 2014)

**Table 2:** Zone of Inhibition in (250 mg seeds powder/mL) the solvent extracts of *L. sativum*

Microorganisms	Zone of inhibition (in mm)				
	Stand.	n-Hexane	Ethanol	Methanol	Water
<i>E. coli</i>	24 ± 0.5	-	13 ± 0.21	8 ± 0.10	10 ± 0.00
<i>S. aureus</i>	20 ± 0.2	8 ± 0.12	15 ± 0.32	10 ± 0.05	8 ± 0.05
<i>C. albicans</i>	*	8 ± 0.15	16 ± 0.24	20 ± 0.43	-

NB: **Stand.** = Chloramphenicol    '\*' The Standard was not available    '-' No observed inhibition  
Results presented as mean ± SD (triplicate)



**Figure 2:** Bioactivities of the different extracts of *L. sativum* against *E. coli*, *S. aureus* and *C. albicans*


Similarly, as shown in Table 3 and Figure 3, the ethanol extract of *Rumex nervosus* was found to have the highest activity against *S. aureus* and *C. albicans* with zones of inhibition of 18 mm each. The ethanol extract also displayed characteristic activities against all pathogens and its activity against *E. coli* was the highest (15 mm) compared to the other extracts. The n-hexane and aqueous extracts showed lower activities against *S. aureus* and *E. coli*; there was no observed activity by both extracts against the fungal strain. Comparing the effect of solvent on the bioactivity of both plants showed similar results against all pathogens. In both cases, the ethanol extract showed a better inhibition on bacterial and fungal growth. The extraction was initially done using ethanol followed by methanol and thus some of the active polar metabolites could have been extracted via ethanol and thus the better activity observed can be attributed to the order of extraction. The effect of extracting solvent on the bioactivity of plant extracts was reported by Beata *et. al.*, 2007. The results obtained revealed that the tested polar extracts possess more potential antimicrobial activities against *E. coli*, *S. aureus* and *C. albicans* when tested using Agar well plate method. As indicated in Table 1, the different active metabolites present in the methanol and ethanol extracts were responsible for the observed activities. Figure 4 shows some representative pictures that show the characteristic zones of inhibitions for the bacterial and fungal strains.

**Table 3:** Zone of Inhibition in (250 mg leaves powder/mL) the solvent extracts of *R. nervosus*


Microorganisms	Zone of inhibition (in mm)				
	Stand.	Methanol	n-Hexane	Ethanol	Water
<i>E. coli</i>	24 ± 0.60	15 ± 0.33	8 ± 0.10	10 ± 0.13	6 ± 0.15
<i>S. aureus</i>	20 ± 0.32	16 ± 0.25	10 ± 0.22	18 ± 0.41	10 ± 0.00
<i>C. albicans</i>	*	15 ± 0.22	-	18 ± 0.50	-

NB: **Stand.** = Chloramphenicol    '\*' The Standard was not available    '-' No observed inhibition  
Results presented as mean ± SD (triplicate)





**Figure 3.** Bioactivities of the different extracts of *R. nervosus* against *E. coli*, *S. aureus* and *C. albicans*



**Figure 4:** Ethanol extract of *L. sativium* against *S. aureus* (A), methanol extract of *R. nervosus* against *C. albicans* (B) and *S. aureus* (C)

## 5. CONCLUSIONS

The phytochemical analysis showed the presence of alkaloids, flavonoids, tannins, saponins and others in most of the extracts; this indicates that the extracts contain some of the prominent secondary metabolites. The different solvents exhibited a wide range of antimicrobial activities and based on the results observed the ethanol and methanol extracts were the two best extractive solvents for antimicrobial activities of the seeds of *L. sativium* and leaves of *R. nervosus*. Therefore, the results establish a good support of the use of *R. nervosus* and *L. sativium* in traditional medicines especially for the people living in remote areas as an alternative treatment for eye diseases.

## CONSENT

It is not applicable

## ETHICAL APPROVAL

It is not applicable

## REFERENCES

- Abdel KM, Safian, A, Kamal MS, Inas O. GC-MC Analysis and Antimicrobial Activity of Fixed oil from Saudi *Lepidium sativum* seeds. Inter. J. Adv. Res. 2017; 5(3):1662-1670.
- Abiyu E, Zemedu A, Ensermu K, Raja N. Ethnobotanical Study of Traditional Medicinal Plants in and around Fiche District, Central Ethiopia. Curr. Res. J. Biol. Sc. 2014; 6(4): 154-167.
- Afaf IA, Manuaki IA, Hassan A. Antimicrobial Activities of Extracts for Some Medicinal Plants. Inter. J. Adv. Appl. Sc. 2015; 2(2):1-5.
- Amit T, Sanchita S, Suchita S. Chemical Analysis of Leaf Extracts of *Calotropis procera*. Inter. J. Sc. Res. Publ. 2014; 4(1): 2250-3153.
- Aragaw Z, Nega B. The Antimicrobial Effect of *Lepidium sativum* Extract on Some Clinical Isolated Bacteria from University of Gondar Teaching Hospital. Biotech. Inter. 2016; 9(3): 69-82.
- Asra J, Rani S, Mohammed I, Abdul SM. A Review on *Lepidium Sativum*. Indo Am. J. P. Sci. 2017; 4(8): 2223-2227.
- Beata, D, Agnieszka S, Rafał W. The Influence of Time and Type of Solvent on Efficiency of the Extraction of Poly-Phenols from Green Tea and Antioxidant Properties Obtained Extracts. Acta. Sci. Pol. Tech. Aliment. 2007; 6(1), 27-36.
- Bruno E. Research in Clinical Phytopharmacology to Develop Health Care in Developing Countries: State of the art and perspectives. Phytopharmacol. 2013; 4(2):149-205.
- Endalkachew A, Negesse M. *In vitro* Antibacterial Activity of *Rumex nervosus*, *Plantago anceolata*, *Solanum incanum* and *Lepidium sativum* against selected bacterial pathogens of human and animals. Ethio. Veter. J. 2016; 20 (2):119-131.
- Haider MH. Analysis of Trace Heavy Metals and Volatile Chemical Compounds of *Lepidium sativum* Using Atomic Absorption Spectroscopy, Gas Chromatography–Mass Spectrometric and Fourier Transform Infrared Spectroscopy. Res. J. Pharm. Biol. Chem. Sc. 2016; 7(4): 25-29.
- Harbone JB. Methods of extraction and isolation in phytochemical methods, Chapman and Hall, London.1998; 60-66.
- Hero FS, Jwan DT. Antibacterial Activity of *Lepidium sativum* and *Allium porrum* Extracts and Juices against Some Gram Positive and Gram Negative Bacteria. Med. J. Islam. 2012; 20(1):10-16.
- Indumathy R, Ajithadas A. Free Radical Scavenging Activities, Total Phenolic and Flavonoid Content of *Lepidium sativum*. Intern. J. Pharm. Pharmaceut. Sc. 2013; 5(4): 634-637.
- Mohammed MQ, Rasool K, Mujeeb R, Abdulrahman A. Chemical Composition and *In vitro* Anticancer, Antimicrobial and Antioxidant Activities of Essential Oils and Methanol Extract from *Rumex nervosus*. Nat. Prod. Res. 2018; 19:1-6.
- Mussie SD, Kareru PG, Keriko JM, Berhane GN. Ethnobotanical Survey and Preliminary Phytochemical Studies of Plants Traditionally Used for Diabetes in Eritrea. Europ. J. Med. Plants. 2015; 9(2), 1-11.
- Nishat AS. Comparative Study on the Antibacterial Activities of Neem oil, Mustard oil and Black Seed Oil Against *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Bacillus cereus*, *Salmonella typhi* and *Pseudomonas aeruginosa* MSc Thesis in Microbiology, University of BRAC, Dhaka, Bangladesh. 2017.
- Pragya B, Dinesh B, Anita P, Anita RS. Antimicrobial Activities of *Lepidium sativum*: Medicinal Plant Used in Folklore Remedies in India. J. Pharm. Res. 2012; 5(3):1643-1645.
- Pragya B, Dinesh B, Yadav OP, Anita P. *In vitro* Antioxidant Activity and Phytochemical Analysis of Seed Extracts of *Lepidium sativum*- a Medicinal Herb. J. Biosci.Tech. 2011; 2(6):410- 415.
- Rao KNV, Sunitha C, David B, Sandhya S, Mahesh V. A Study of the Nutraceuticals from the genus *Rumex*. J. Drugs Med. 2011; 3(1):76-88.
- Rehab A, Salem E. Phytotoxicity of Synthesized Butenolide: Response of Medicinal Plant *Lepidium sativum* L. Seed Germination and Seedling Development to Different Concentrations. Eur. J. Exp. Biol. 2013; 3(2):396-401.
- Shahla S, Farshad G, Mehdi H, Zahra S. *In- vitro* Antimicrobial Assessment of *Lepidium Sativum* L. and *Capsicum annuum* L Extracts. J. J. Microbiol. Path. 2014; 1(1): 002.
- Shama, IA, Abdullah AA, Adam KMO, Aldai MB, Omer AA, Warda SA. *In vitro* Antimicrobial Activity of *Rosmarinus officinalis* Leave Extracts. J. Agri-Food. App. Sc. 2014; 2(1): 15-21.
- Trease GE, Evans WC. Pharmacognosy, 11<sup>th</sup> Edn, Bailliere Tindall, London. 1989; 45-50.
- Zewdu B. Traditional Use of Medicinal Plants by the Ethnic Group of Gondar Zuria District, North-Western Ethiopia. J. Nat. Rem. 2013; 13 (1): 47-53.