

Original Research Article

Can Calcium and Magnesium Infusion Before and after Oxaliplatin Administration Improves Its Neurotoxic Effect?

ABSTRACT

Introduction: The oxaliplatin neurotoxicity is one of two distinct phases: an acute and a dose-limiting cumulative phase. Large doses of I.V Ca & Mg is given before and after oxaliplatin in FOLFOX protocol. **Aim:** we aimed to evaluate the efficacy of Ca & Mg in improving or eliminating oxaliplatin-induced neurotoxicity. **Study design:** Randomized double controlled placebo study. **Place and Duration of Study:** Clinical Oncology department at Mansoura university hospital, Egypt at the period from July 2014 up to December 2016 inclusive. **Methodology:** In our study, 140 Patients with adenocarcinoma of the colon received 6 months of FOLFOX protocol. Patients were choiced to receive I.V calcium gluconate plus magnesium sulfate (1 g for both of them) in 100 mL of D5W for about 30 minutes immediately before and after each cycle of oxaliplatin (70 patients), or an identical looking placebo shortly before and after each cycle of oxaliplatin (70 patients) in a random manner. The primary end point was a comparison between neurotoxicity assessed by Common Terminology Criteria for Adverse Effects (CTCAE) version 4.0 in both groups. **Results:** there was statistically significant difference as regard grade of neuralgia. Grade II neuralgia is higher in the study group II (45.7%) compared to (22.9%) in the study group I. There was statistically significant difference as regard grade of paresthesia. Grade II & III are higher in study group II (51.4%, 14.3%) compared to (30.0%, 4.3%) in study group I. **Conclusion:** Ca and Mg administration with oxaliplatin have been well tolerated. Our results support the role of Ca & Mg in decreasing neurotoxicity associated with oxaliplatin.

Key words: Colorectal cancer, Neurotoxicity, Oxaliplatin, Ca gluconate

INTRODUCTION

Oxaliplatin is a member of widely accepted standard protocols for treatment of stage III and stage IV colorectal cancer (CRC). Also, it is now commonly used in the treatment of other malignancies including esophagogastric and pancreatic cancers (1).

Chemotherapy protocols that contain oxaliplatin are widely used and having a good effects with response rate as high as 53% and a quite low rate of side effects as nausea, vomiting and leucopenia. However, neurotoxic side effects restrict its use and cause patient irritability. The oxaliplatin causing neurotoxicity is one of two distinct phases: a transient, acute phase that can appear during or after administration of oxaliplatin with a short time and a dose-limiting cumulative sensory phase of neuropathy (2).

The neuropathy associated with oxaliplatin is different from the neuropathy that happens with cisplatin. The two agents will lead to a glove stocking neuropathy that increases with elevating the dose. But oxaliplatin is accompanied by an acute neuropathy in the form of muscle twitches, sensation of discomfort in the throat and intolerability to the coldness which is generally associated with each oxaliplatin dose (3). The acute form occurs within short minutes of administration of oxaliplatin and is able to sustain for one to two days in the form of paresthesia and tingling. Sometimes, there is an unordinary sensation in the tongue or limb stiffness, and may be increased by cold exposure. The rate of acute form of neuropathy ranges from 81.5 up to 98% (4). The sustained sensory chronic neuropathy appears as paresthesia and proprioceptive changes. They do not disappear in between cycles of chemotherapy. It happens in about 15% of patients after total doses of oxaliplatin ranges from 780-850 mg/m² (2).

Large doses of I.V Ca & Mg is given before and after oxaliplatin in FOLFOX protocol. At the clinical settings; they represent the most used regimen for the prevention of FOLFOX neuropathy. The oxalate was released from the metabolism of oxaliplatin. This oxalate is chelating the Ca & Mg. They are involved in the function of ion channels in nerve membranes. So, Ca & Mg may eliminate or improve this induced neuropathy (5). Unlike the acute form neurotoxicity, the cumulative toxicity of oxaliplatin appears to be due to direct toxic effect on the nerve itself (6).

The benefit was noticed for both the acute and chronic neuropathy, with no evident antagonistic effect on the antitumor effect of FOLFOX. Subsequently, a lot of oncologists used I.V Ca & Mg as a part of the pre-treatment medications for patients under FOLFOX protocol. Divalent cations had the power to change voltage-gated sodium channels. It is hypothesized that the acute neuropathy of oxaliplatin is related to calcium chelation caused by oxalate. Increasing the extracellular calcium has been found to increase the sodium channel closure. And, this leads to decrease the over excitability of the peripheral neurons which seen in the neuropathy caused by oxaliplatin. Magnesium supplementation has been studied in preventing hypomagnesemia caused by cisplatin (7), (6).

AIM

In this prospective randomized double controlled placebo study, we aimed to evaluate the efficacy of Ca & Mg in improving or eliminating oxaliplatin-induced neurotoxicity.

PATIENTS AND METHODS

In our randomized double controlled placebo study, 140 Patients with adenocarcinoma of the colon who attended to clinical oncology & nuclear medicine department at Mansoura university hospital at the period from July 2014 up to December 2016 inclusive. The patients were treated with curative-intent resection. They were scheduled

to receive 6 months (12 cycles) of FOLFOX protocol as adjuvant treatment (ie. oxaliplatin, leucovorin and fluorouracil, involving oxaliplatin 85 mg/m²) every 2weeks.

The studied patients needed to have adequate hematological parameters, serum total bilirubin and serum creatinine to allow chemotherapy and normal levels of calcium and magnesium. Women at the age of childbearing period must have a negative pregnancy test. A central venous line was inserted before the beginning of chemotherapy protocol. Patients were excluded from the study if they had a pre-existing peripheral neuropathy of any grade or had received neurotoxic chemotherapy like oxaliplatin, cisplatin, taxanes or vinca alkaloids previously prior to existing treatment.

At the beginning of our study and before each 2-week cycle of chemotherapy; we performed a brief history, physical examination, and CBC for each patient. Neurotoxicity assessments were obtained in each setting. The primary neuropathy assessment was described by Common Terminology Criteria for Adverse Effects (CTCAE) version 4.0.

Questionnaires were obtained from the patients before each dose of FOLFOX to provide data regarding grades of peripheral motor neuropathy, peripheral sensory neuropathy, neuralgia, and paresthesia.

Patients were choiced to receive I.V calcium gluconate plus magnesium sulfate (1 g for both of them) in 100 mL of D5W for about 30 minutes immediately before and after each cycle of oxaliplatin (70 patients), or an identical looking placebo shortly before and after each cycle of oxaliplatin (70 patients) in a random manner. Patients were classified by age (less than 50 years vs. 50 years and more) and gender. The primary end point was a comparison between neurotoxicity assessed by Common Terminology Criteria for Adverse Effects (CTCAE) version 4.0 in both groups.

Statistical analysis

SPSS (version 21) were used to analyze data. The normality of data was tested with one-sample Kolmogorov-Smirnov test. Qualitative data were described by using number and percent. Chi-square was used to test association between categorical variables. Continuous variables were presented as mean $\pm$ SD (standard deviation). Student – test was used to compare means of 2 groups.

The level of significance:

For all above mentioned statistical tests done, the threshold of significance is fixed at 5% level (p-value).The results were considered:

- When the probability of error is more than 5%, so it is non-significant ($p > 0.05$).
- When the probability of error is less than 5%, so it is significant ($p \leq 0.05$).
- When the probability of error is less than 0.1%, so it is highly significant ($p \leq 0.001$).

The results are more significant when the smaller p- value is obtained.

RESULTS

Two arm, placebo-controlled, randomized, blind, parallel group design was used. One hundred and forty patients started FOLFOX therapy between July 2014 till December 2016 was enrolled in the present study and divided into two groups: the study group and the placebo one; each of them has 70 patients per each. Our data shows that the two study groups were matched regarding sex and age; (50%) were males and (50%) were females for both groups, while age for study group I was 48.30 (8.38) and for study group II was 48.95 (8.12) with no statistically significant difference (*P-value* >0.05) as shown in table 1.

As regard toxicity; there was no statistically significant difference (*P-value* >0.05) regarding grade of sensory neuropathy and grade of motor neuropathy. However it is not statistically significant; grade II& III sensory neuropathy were higher in the study group II (31.4%, 7.1%) compared to (18.6%, 4.3%) in the study group I. Also, grade II & III motor neuropathy was higher in the study group II (30.0%, 7.1%) compared to (20.0%, 5.7%) in the group I as mentioned in the table1.

On the other hand, there was statistically significant difference (*P-value* <0.05) as regard grade of neuralgia. Grade II neuralgia is higher in the study group II (45.7%) compared to (22.9%) in the study group I (OR=2.9). In addition, there was statistically significant difference as regard grade of paresthesia. Grade II & III are higher in study group II (51.4%, 14.3%) compared to (30.0%, 4.3%) in study group I (OR =3.3, 6.4) respectively as described by table 1 and figures 1&2.

TABLES

Table (1): Distribution of patient's sex, age and grades of neurotoxicity

Items	Study Group I n=70		Placebo Group II n=70		χ^2 (<i>P value</i>)	OR (95%CI)
	No	%	No	%		
Sex						
Male	35	50.0	35	50.0	-	-
Female	35	50.0	35	50.0		
Age / years						
Mean (SD)	48.30 (8.38)		48.95 (8.12)		t=0.471	0.638
Min-Max	33-63		33-61			
G of Sensory neuropathy						
1	54	77.1	43	61.4	-	r (1)
2	13	18.6	22	31.4	3.5 (0.06)	2.1 (0.96-4.7)
3	3	4.3	5	7.1	0.98(0.32)	2.09 (0.47-9.2)
G of Motor neuropathy						
1	52	74.3	44	62.9	-	r (1)
2	14	20.0	21	30.0	2.1 (0.15)	1.8 (0.8-3.9)
3	4	5.7	5	7.1	0.31(0.57)	1.5 (0.4-5.9)

G of Neuralgia						
1	51	72.9	35	50.0	-	r (1)
2	16	22.9	32	45.7	8.3 (0.004)	2.9 (1.4-6.1)
3	3	4.3	3	4.3	0.2 (0.65)	1.5 (0.3-7.6)
G of Paresthesia						
1	46	65.7	24	34.3	-	r (1)
2	21	30.0	36	51.4	10.5 (0.001)	3.3 (1.6-6.8)
3	3	4.3	10	14.3	8.2 (0.004)	6.4 (1.4-3.5)

FIGURES




Figure (1): Distribution of the grade of neuralgia in both groups




Figure (2): Distribution of the grade of paresthesia in both groups

DISCUSSION

The question of the usage of Ca and Mg as a neuroprotectant for oxaliplatin is difficult to be definitively answered. Many studies are necessary to obtain the answer of this debate.

There have been 3 published retrospective trials for the utility of Ca & Mg as a neuroprotectant against oxaliplatin neurotoxic effect. Gamelin et al studied (7) 161 patient in a retrospective cohort trial and they reported positive results as compared with our study results. Knijn et al (8) studied 732 patients entered in a clinical study randomly with Xeloda, oxaliplatin, and bevacizumab versus Xeloda, oxaliplatin, and cetuximab, with the use of Ca & Mg. The retrospective analysis, for 551 patients who received Ca and Mg during their initial treatment cycle, there was a slightly lower incidence of any physician-judged neurotoxicity (85% vs. 92%, respectively; ***P* 0.02**) and physician-judged grade 2 neurotoxicity (40% vs. 45%; ***P* 0.22**), in patients received Ca and Mg versus the patients who did not take Ca and Mg in association with their cycles (181 patients). In another retrospective study (9), ninety patients receiving FOLFOX protocol also administrated Gosha-Inki gan, a traditional Japanese herbal compound, Ca & Mg alone, Ca & Mg plus Gosha-Inki gan, or neither agent. No benefit was gained for Ca & Mg in this study which is contrary to our study results.

Two prospective double-blind published trials. The CONCEPT trial did not show any significant advantage which is against our results, but the N04C7 trials suggested benefit for Ca & Mg as we demonstrated in our patients

(3) . In another double-blinded, placebo-controlled clinical study which had 3 study arms, with about thirty patients in each study arm; the patients received Ca, Mg, and glutathione, or placebo. There were no significant differences statistically in the neurotoxic effect happened among the study groups (10).

Nikcevich et al performed a randomized controlled trial that studied the effect of Ca and Mg on neuropathy in stage II or III colon cancer patients treated with FOLFOX as adjuvant settings. Their study was also stopped after the release of the interim results of the CONCEPT study (11).

Also, Chay and his colleagues reported in 2010 that their study was stopped early based on the first negative results of the CONCEPT trial. Median follow up was 8.7 months. Overall 22 out of 27 patients developed neuropathy. Arm A showed a rate of 77% of subjective neuropathy & Arm B showed 86% ($P = 0.6$). There was no statistical significant difference in neuropathy between both arms, during or at the end of treatment. Median score of objective neuropathy was 6 in Arm A & 0 in Arm B ($P = 0.02$) (12).

Gamelin and his colleagues reported in 2008 another study suggesting that Ca & Mg was beneficial. This report discussed preliminary data from the NEUROXA trial involving fifty two patients suffering from metastatic colorectal cancer; 50% of patients received Ca & Mg and 50% did not. They noticed a difference in the incidence of grade 3 neurotoxicity between the two study groups (5% vs. 24%; $P-0.001$) (13).

Also, Gobran published in 2013 findings supporting that the reduction in neurotoxicity in patients receiving Ca and Mg with oxaliplatin infusions was statistically significant. The study was performed from July 2008 to February 2011. They studied 30 patients in arm 1 who received calcium/ magnesium and another 30 patients in the placebo arm. All the patients received regimen containing oxaliplatin as adjuvant treatment. They used the NCI-CTCAE version 3.0 criteria for neuropathy assessment. Results concluded that 23.3% of patients receiving Ca and Mg developed neuropathy by the completion of therapy, vs. 46.6% in the placebo arm, a statistically significant difference (14). This was comparable with our results as the study group developed sensory neuropathy in about 22.9% (GII, GIII), vs. 38.5% (GII, GIII) in placebo arm but it is not statistically significant. Also, motor neuropathy was 25.7% in study group vs. 37.1% in the placebo arm but also with no significance statistically. As regard neuralgia in our study, the placebo group showed 45.7% of patients, vs. 22.9% in the study group I and it shows good significance statistically.

During the last few years, many studies suggest that, with regional changes worldwide, many oncologists have been using Ca & Mg in their routine work and this is clear from many studies mentioned previously. Our results support this issue which currently notes that Ca & Mg can be used for the prevention of neuropathy caused by oxaliplatin or to decrease its degree.

CONCLUSION

Ca and Mg administration with oxaliplatin protocols have been well tolerated. Our study results support the role of Ca & Mg administration in decreasing neurotoxicity associated with oxaliplatin. Further studies will be needed to adjust the optimal dose, repetition, and infusion time. Also, these studies will be helpful to determine the role of these administrations in oxaliplatin causing neuropathy.

184 **DISCLOSURE**

185 Authors have declared that no competing interests exist.

186 **ETHICAL APPROVAL**

187 The study has the approval of the IRB committee of the Faculty of Medicine (MFM-IRB), Mansoura
188 University, Egypt. The code number is R/17.05.66.

189 **REFERENCES**

- 189 1. Vatandoust S, Joshi R, Pittman KB, Esterman A, Broadbridge V, Adams J, et al. A descriptive
190 study of persistent oxaliplatin-induced peripheral neuropathy in patients with colorectal cancer. Supportive
191 Care in Cancer. 2014;22(2):513-8.
- 192 2. Ao R, Wang Y-H, Li R-W, Wang Z-R. Effects of calcium and magnesium on acute and chronic
193 neurotoxicity caused by oxaliplatin: a meta-analysis. Experimental and therapeutic medicine.
194 2012;4(5):933-7.
- 195 3. Grothey A, Nikcevic DA, Sloan JA, Kugler JW, Silberstein PT, Dentchev T, et al. Intravenous
196 calcium and magnesium for oxaliplatin-induced sensory neurotoxicity in adjuvant colon cancer: NCCTG
197 N04C7. Journal of Clinical Oncology. 2010;29(4):421-7.
- 198 4. Cersosimo RJ. Oxaliplatin-associated neuropathy: a review. Annals of Pharmacotherapy.
199 2005;39(1):128-35.
- 200 5. Loprinzi CL, Qin R, Dakhil SR, Fehrenbacher L, Flynn KA, Atherton P, et al. Phase III
201 randomized, placebo-controlled, double-blind study of intravenous calcium and magnesium to prevent
202 oxaliplatin-induced sensory neurotoxicity (N08CB/Alliance). Journal of Clinical Oncology.
203 2013;32(10):997-1005.
- 204 6. Saif MW, Reardon J. Management of oxaliplatin-induced peripheral neuropathy. Therapeutics
205 and clinical risk management. 2005;1(4):249.
- 206 7. Gamelin L, Boisdron-Celle M, Delva R, Guérin-Meyer V, Ifrah N, Morel A, et al. Prevention of
207 oxaliplatin-related neurotoxicity by calcium and magnesium infusions. Clinical Cancer Research.
208 2004;10(12):4055-61.
- 209 8. Knijn N, Tol J, Koopman M, Werter M, Imholz A, Valster F, et al. The effect of prophylactic
210 calcium and magnesium infusions on the incidence of neurotoxicity and clinical outcome of oxaliplatin-
211 based systemic treatment in advanced colorectal cancer patients. European Journal of Cancer.
212 2011;47(3):369-74.
- 213 9. Kono T, Mamiya N, Chisato N, Ebisawa Y, Yamazaki H, Watari J, et al. Efficacy of goshajinkigan
214 for peripheral neurotoxicity of oxaliplatin in patients with advanced or recurrent colorectal cancer.
215 Evidence-based complementary and alternative medicine. 2011;2011.
- 216

- 217 10. Dong M, Xing P, Liu P, Feng F, Shi Y. Assessment of the protective effect of calcium-magnesium
218 infusion and glutathione on oxaliplatin-induced neurotoxicity. Zhonghua zhong liu za zhi [Chinese journal
219 of oncology]. 2010;32(3):208-11.
- 220 11. Nikcevich D, Grothey A, Sloan J, Kugler J, Silberstein P, Dentchev T, et al. Effect of intravenous
221 calcium and magnesium (IV CaMg) on oxaliplatin-induced sensory neurotoxicity (sNT) in adjuvant colon
222 cancer: results of the phase III placebo-controlled, double-blind NCCTG trial N04C7. Journal of Clinical
223 Oncology. 2008;26(15_suppl):4009-.
- 224 12. CHAY WY, TAN SH, LO YL, ONG SYK, NG HC, Gao F, et al. Use of calcium and magnesium
225 infusions in prevention of oxaliplatin induced sensory neuropathy. Asia-Pacific Journal of Clinical
226 Oncology. 2010;6(4):270-7.
- 227 13. Gamelin L, Boisdron-Celle M, Morel A, Poirier AL, Berger V, Gamelin E, et al. Oxaliplatin-related
228 neurotoxicity: interest of calcium-magnesium infusion and no impact on its efficacy. Journal of Clinical
229 Oncology. 2008;26(7):1188-9.
- 230 14. Gobran NS. Role of calcium and magnesium infusion in prevention of oxaliplatin neurotoxicity. A
231 phase III trial. The Chinese-German Journal of Clinical Oncology. 2013;12(5):232-6.

232

233