

Colon Cancers: Epidemiological And Histopathological Aspects In Cameroon

ABSTRACT:

Objective: To determine the epidemiological and histological profile of colon cancer in Cameroon.

Material and methods: It was a retrospective descriptive and analytical study on cancers of the colon, histologically proven for 13 years (2004-2016), listed in the registers laboratories of pathology and Oncology of the national territory. The variables studied were the frequency, age, gender, risk factors, location and histopathologic type.

Results: We identified 1047 cases of digestive cancers. The colon with 366 cases (26. 01%) was the second most common location behind the stomach. The average age of the patients was 52.82 ± 15.92 years, with extremes ranging from 6 to 89 years old. The male was the most represented with 52. 73% (193 cases), with a sex ratio M/F of 1.12. A tumour was sigmoid localisation in 37. 59% of the cases (100 cases on 266 listed locations). Adenocarcinomas were the found histological type with 311 cases (84. 97%) followed by the Kaposi Sarcoma (21 cases; 5.74 percent) and lymphomas (17 cases; 4.65%).

Conclusion: Colon cancer remains a relatively common pathology in Cameroon where it ranks second among malignant tumours of the digestive tract. Women remain less affected than men by this pathology whose dominant histological type is adenocarcinoma. However, these data remain relative given the absence of a cancer register at the national level.

Keywords: Cancer; Colon; epidemiology; histopathology; Cameroon

1. INTRODUCTION

Colon cancer is the most common malignant tumour of the digestive tract, developing from cells lining the inner lining of the colon [1, 2]. It accounts for 9.7% of all cancers globally [3-5] and ranks third after lung (13%) and breast (11.9%) cancers [6-8]. Malignant colon tumours represent the third most common disease in men and the second in women with an estimated incidence of 1.4 million cases in 2012, the sex ratio M / F is 3/2 with almost 95% of patients diagnosed after 50 years [1-3]. Incidence rates vary from continent to continent, and high incidence countries are Western Europe, North America, Australia, New Zealand and Japan [4-8]. In France, the sex ratio M/ F is 1.22 with a mean age of diagnosis of 71

34 years in men and 73 years in women [3, 9, 10]]. In Canada it is the 2nd most diagnosed form
35 of cancer (13% of all cancers combined) with more than 90% of patients older than 55 years;
36 the sex ratio M / F is 1.25 [11, 12]. In Africa the incidence is lower; this is the case of Togo,
37 the sex ratio M / F is 3/2 with an average age of 46.7 years [13]. In Morocco the mean age of
38 onset 54.48 ± 14.75 with slight female predominance [14]. In 2012, according to the
39 International Agency for Research on Cancer (IARC), colon cancer was the 4th leading cause
40 of death in the world with 694,000 deaths behind lung, liver and stomach cancers [1]. In West
41 Africa, there is a low mortality rate for colon cancer of 3.5 per 100,000 men and 3 per
42 100,000 women [15]. Although colonic polyposis is considered to be the major risk factor for
43 the development of colon cancer (HNPCC: hereditary non-polyposis colorectal cancer) [16,
44 17], other risk factors are incriminated in the genesis of this pathology in particular, age > 50
45 years, obesity, a diet low in fruits and vegetables, sedentary lifestyle and smoking [18, 19].
46 The diagnosis of certainty is colonoscopic associated with an anatomopathological
47 examination. The most common histological types are adenocarcinoma in more than 90% of
48 cases (Liberkhuinen, colloid or mucinous, with signet ring cell contingents) [20]. The most
49 frequent locations are the ascending colon, the descending colon, and the sigmoid [21, 22].
50 The primary means of treatment are surgery for locoregional forms, chemotherapy and
51 radiotherapy for extended forms, including palliative care [23]. In Cameroon, the incidence
52 of colon cancer is estimated in Yaoundé at 2.9% [24]. In the littoral region, it ranks 2nd in
53 digestive cancers behind stomach cancer with an incidence of 22.79%; the mean age of onset
54 is 51.83 ± 17.34 years with a sex ratio of 1.69 [25]. However, it should be noted that these
55 two studies are regional and we do not have data at the national level. It is therefore essential
56 for us to study through this work the epidemiological profile and histopathological aspects of
57 this severe condition over a period of 13 years (2004-2016) in Cameroon to have recent data,
58 which would allow a better control of this pathology in our context.

59 **2. MATERIAL AND METHODS**


60 This is a retrospective descriptive and analytical study of histologically proven malignant
61 colon tumors, diagnosed between January 2004 and December 2016. The study took place in
62 the main public and private pathological anatomy laboratories in Cameroon. We needed the
63 reports of histopathological examinations of the various laboratories solicited, all the
64 necessary documentation relating to our subject (books, journals, specific publications ...),
65 and a well-defined office equipment. The samples generally come from previously
66 unresolved surgery, cancerology or gastroenterology departments. Once in the pathology

67 departments, they are fixed at 10% formalin, and then the macroscopic study in which the
 68 pieces are cut. The pieces are dehydrated by passing through several tanks of alcohol at
 69 increasing concentrations, then included in paraffin, then cut with a microtome to a thickness
 70 of 5 micron. They are then deparaffinized by xylene lightening, and the staining is done with
 71 haematin-eosin followed by a reading made using a microscope. Only patients for whom the
 72 diagnosis was confirmed by histology were included in the study. The information obtained
 73 included frequency, age, sex, histological type of the tumor. Data entry was done using
 74 computer based statistical Package for Social Sciences (SPSS) version 20. The elements of
 75 descriptive statistics were used to calculate the frequencies and proportions.

76 3. RESULTS

77 3.1. Frequency

78 At the end of our study, we collected 1407 digestive cancers, 574 of these cancers were
 79 stomach cancers with a frequency of 40.80% making it the first digestive cancer of both sexes
 80 men 312 cases and women 262 cases (Figure 1).



81

82 **Figure 1. Distribution of cancers according to the segment of the digestive tube**

83 The frequency of onset was 28.15 cases / year. We have seen an increase over the years from
 84 about 22 cases / year in the first seven years of our study to 35 cases / year for the last five
 85 years. (See Figure 2)




Figure 2: Evolution of colon cancer in the years from 2004 to 2016 (n = 366)

3.2. Distribution by sex

Of the 366 cases of colon cancer found, the male sex is represented by 193 cases (52.73%) or 13.72% of all digestive cancers and the female sex represented by 173 cases (47.27% or 12.30% of all digestive cancers. The sex ratio **M / F** is 1.12. (Table I)

Table I: Distribution of digestive tract cancers by sex


Organ	Stomach		Colon		Rectum		Anus		Esophagus		S intestine		Total
Sex	H	F	H	F	H	F	H	F	H	F	H	F	
Effective	312	262	193	173	130	110	42	49	67	22	24	23	1407
% Effective	22.17	18.62	13.72	12.30	9.24	7.82	2.99	3.48	4.76	1.56	1.71	1.63	100
Total	574		366		240		91		89		47		1407
% Total	40.80		26.01		17.06		6.47		6.33		3.34		100

S-Small

3.3. Distribution by age

96 As shown in the figure 3, the average age of diagnosis is 52.82 ± 15.92 years, with extremes
 97 ranging from 6 to 89 years. The majority of patients were between 50 and 59 years old (86
 98 cases; 24.57%).

99



100

101 **Figure 3: Distribution of patients by age group. n = 366**

102 In men the average age is 53.62 ± 15.38 years with extremes ranging from 6 to 89 years; the
 103 majority of cases are between 50 and 59 years old, ie 52 patients (27.6%). In women, the
 104 mean age of diagnosis is 51.90 ± 16.49 years with extremes ranging from 7 to 88 years; the
 105 majority of cases are between 40 and 59 years old with 68 patients (41.46%).

106 **Distribution according to risk factors**

107 Risk factors were highlighted in 235 patients. The most common are cold cuts (21.28%),
 108 alcohol (17.02%), obesity (13.62%) and smoking (12.34%). (Figure 4)




Figure 4: Distribution of Risk Factors n = 235

3.4. Tumor localization

In the study, 266 localizations were highlighted in this study. As showed in figure 5, the tumor was sigmoid localization in 37.59% of cases (100cas / 266).




Figure 5: Distribution by localization n = 266

3.5. Anatomopathology

3.5.1. Types o sampling

Of the 366 cases of colon cancer identified in our study, the type of sample was specified on 357 cases, of which 191 (52.19%) were derived from operative specimens and 166 (45.35%) were biopsies.

3.5.2. Histological type

The most common varieties are adenocarcinoma, the most common subtype being Liberkhunien. In patients whose age range varies between 0 and 9 years. It should be noted also the census of a case of the type Leiomyosarcoma or 0.27% of all histological types encountered. (Table II)

Table II: Distribution of Histological Types by Age Group

	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	≥80	Total
Lieberkühn's ADK	2	2	6	34	57	64	61	27	11	264
Mucinous ADK	1	2	3	9	9	13	6	4		47
UDT Carcinoma					1	1				2
squamous cell carcinoma					1	1				2
AQM carcinoma				1						1
Carcinoid tumor			1	1	2	1	3	1		9
NHML		2	2	2	1	3	4	2	1	17
Kaposi's sarcoma			1	1	2	8	7	2		21
Leiomyosarcoma						1				1
Liposarcoma									1	1
Stromal tumor								1		1
	3	6	13	48	73	92	81	37	13	366

NHML - Non-Hodgkin's Malignant Lymphoma; ADK – Adenocarcinoma; UDT - Undifferentiated; AQM - Adenosquamous

4. DISCUSSION

In the study, colon cancers accounted for 26.01% of all malignant digestive tumors, ranking 2nd after the stomach. Several other studies in the African territory rank them second but with varying frequencies (Peghini et al, Ayite et al, Harouna et al, Sani et al) [26-29]. Similarly, in Niger they represent 28.8% of the digestive tract behind the stomach [30-31]. Globally and in the West, particularly in France, they occupy the first place in digestive cancers [2, 4, 32]. This epidemiological difference could be explained by the high incidence of colic polyposis, whether familial or not in the West, which would raise colon cancer to the highest level of digestive tumoral pathologies on a global scale [16-18].

We collected a total of 366 cases of colon cancer in Cameroon over a period of 13 years or 28.15 cases / year. Indeed, in 2000, Takongmo et al found an incidence of 7 cases / year over 10 years (1987-1996) in Yaoundé [33]. More recently Engbang et al found 7.5 cases / year over 12 years (2004-2015) in the Littoral region [25]. It should be noted, however, that

these two studies are regional whereas ours is of a national character. This epidemiological observation can be explained on the one hand by an increase in the reference due to training and awareness activities on cancer and on the other hand a better knowledge of the diagnostic methods of the pathology, as well as the presence of cancer centers. pathology in the territory.

Moreover, their distribution per year shows a growing increase in the number of cases from 22 cases / year in the first seven years to (2004-2010) to 35 cases / year for the last five years (2011-2016), a rate of increase of 37.14%. African cohort studies are piecemeal; that of Bamako has shown an increasing trend in the number of cases from 0 cases in 2005 to 34 cases in 2011 [22]; in Uganda, Wabinga et al also demonstrated in 2000 an increase in the incidence of this pathology [34].

We noted a male predominance with a sex ratio M / F of 1.12. Globally, the sex ratio is 1.5 [35]. In western Algeria there is a male predominance with 272 cases of colon cancer in men and 229 in women out of 501 cases recorded from 2000 to 2007, a sex ratio of 1.2 [21]. In Dakar in 2014, Ibo et al had found a sex ratio M / F of 1.31 [36]. Given these figures, we conclude that our results are close to those of the literature review. This predominance could be explained by the presence of high-risk factors in men as in women.

The patients in the study were aged 6 to 89 years. The mean age was 52.82 ± 15.92 years. Engbang et al in the Littoral-Cameroon region had a mean age of patients of 51.83 ± 17.10 years with extremes ranging from 7 to 86 years. Similar retrospective studies in Congo, South Africa, and Tunisia found similar comparable age averages of 55, 59, and 58 years, respectively [37-39]. In Togo there is a lower average age of 48.7 years whereas in Senegal there was a higher average age [13; 36]. Thus, colon cancer appears at a relatively lower age among Africans than among Westerners whose peak frequency is between 60 and 70 years [40, 41]. This difference could be explained by the low life expectancy in our regions, but should also look for local factors of carcinogenesis.

The most reached age group was between 50 and 59 years old with 86 cases listed. There is also a high representativeness of young subjects with an unexpected peak of incidence between 30 and 39 years. Subjects under 40 years of age represent 20.06% of cases. The figures found by Takongmo et al found a significantly higher incidence in the under 40 years of 47.61% [33], as well as two Nigerian studies also reporting a significant subgroup of patients under 30 years old [42, 43]. The fact that the cases of colon cancer observed in young people are often associated with genetic markers and particular clinical aspects, invites us to conduct prospective studies on the genetic factors associated with this pathology in Cameroon [16]. We also note in our results 51 cases among those over 70 years of age (14.41% of the population of age). Age is a very questionable prognostic factor; studies have concluded that the occurrence of colon cancer in an elderly person is a factor of poor prognosis [14, 18]. On the other hand, several authors agree to highlight the seriousness of colon cancers in subjects under 40 years of age due to the frequency of histologically aggressive forms such as mucinous and undifferentiated forms [13, 37].

The most common risk factors are excessive consumption of cold cuts, smoking and alcohol. These results are similar to the review of the literature [18, 44]. However, there are also 2 cases (0.85%) of HNPCC syndrome, which differs somewhat from Western analytical studies according to which this syndrome is one of the important risk factors for colon cancer [16,

185 45]. Indeed the risk of occurrence of a malignant colon tumor is very high in case of a history
186 of polyposis or lynch syndrome. Few studies have been done on the incidence of this
187 polyposis in our country. In 2014 in the city of Douala, 86 cases of rectocolic polyps were
188 identified and the most common types were inflammatory (48.83%) and adenomatous
189 (30.23%) polyps [46].

190 Several mechanisms have been proposed to explain the association between red and
191 processed meat with colon cancer. Potential factors such as heterocyclic amines (HCA) and
192 polycyclic aromatic hydrocarbons (PAH) arising in meat cooked at high temperatures, heme
193 iron or nitrates and nitrites used in meat processing have all been hypothesised to play a role
194 [47-50]. Red meat is abundant in heme iron which has been suggested to mediate the
195 formation of intestinal carcinogenic compounds. [47] Egeberg et al has hypothesised that
196 since different red meats contain differing amounts of heme iron, the risk for colon cancer
197 may therefore vary according to the red meat subtype.[51] A more recent hypothesis
198 proposed to explain the relation of red meat consumption and colon cancer suggests that a
199 specific bovine infectious factor may be involved in colon cancer development. Based on the
200 fact that chemical carcinogens (HCA and PAH) as the sole player in colon cancer risk have
201 been questioned and the fact that the increased risk for colon cancer is restricted to
202 populations with high beef consumption the author concludes that a specific beef factor may
203 contaminate the meat which could then be potentially carcinogenic upon transmission to
204 humans. [47, 52]

205 Concerning alcohol, that product can act as a pro-oxidant in tissues, including lung tissue [53,
206 54], and on lipids, including lung membrane lipids [55, 56]. Alcohol can induce the
207 expression of enzymes that are related to carcinogen metabolism , and compounds other than
208 ethanol that are contained in alcoholic beverages may have carcinogenic effects. some
209 mechanisms may explain the alcohol-colon cancer relationship. First, acetaldehyde, an
210 oxidation product of alcohol, may be responsible for colorectal carcinogenesis [55]. Homann
211 et al reported that high levels of acetaldehyde in rat colon degrade folate, a nutrient that is
212 hypothesized to reduce the risk for colorectal cancer [57]. Second, alcohol is an antagonist of
213 methyl-group metabolism and may contribute to abnormal DNA methylation, an early step in
214 colonic carcinogenesis [58]. Finally, greater alcohol intake may increase the risk for
215 colorectal cancer indirectly through immune suppression, delay of DNA repair, activation of
216 liver procarcinogens by induction of cytochrome P-450 enzymes, or changes in bile acid
217 composition [55].

218 Colon cancer risk is 18% higher in men who are overweight (body mass index [BMI] 25-
219 29.9) and 48% higher in men who are obese (BMI 30+), compared with men of a normal
220 weight (BMI 18.5-24.9).[59] Colon cancer risk is 12% higher in women who are obese,
221 compared with women of a normal weight.[59] the association in obese women may be
222 stronger in premenopausal than postmenopausal women[60].The study results showed that
223 there was a strong risk for proximal colon cancer or distal colon cancer higher with high BMI
224 or WC levels [61]. Obesity is considered one important risk factor for many types of solid
225 cancers, especially for colon [61]. Previous reviews have indicated that obesity is associated
226 with 7% to 60% greater risk of colon cancer compared with normal weight individuals [62].
227 Currently, several possibilities have been hypothesized. Two hormonal systems – the
228 insulin/insulin-like growth factor (IGF) axis and adipokines (adiponectin and leptin) – are the

most studied candidates. First, the involvement of insulin and IGF-1 in colorectal carcinogenesis has been supported by experimental and clinical studies [63]. Circulating total IGF-I, a major determinant of free IGF-I concentrations, is associated with increased risk of colorectal advanced adenomas and cancer [61]. The main reason is that increased free IGF-I with concomitant changes of environment mitogenesis and anti-apoptosis in the cellular favouring tumour formation . Moreover, there is an increased risk of colon cancer development associated with type 2 diabetes [61]. Previous studies have demonstrated that the fat itself can also influence colon cancer risk [64, 65]. Adipocytes and preadipocytes could promote proliferation of colon cancer cells [66]. For Ogino et al, fatty acid synthase overexpression has been shown to be associated with colon cancer phenotype [67]. Adipokines such as adiponectin, leptin are also associated with the risk of colon cancer. Adiponectin as an insulin-sensitizing agent and a negative regulator of angiogenesis is secreted mainly from visceral adipose tissue, which could inhibit colon cancer growth in animal models, and its circulating concentrations was associated with colon cancer risk in clinical trials [68]. Leptin could also favour that pathology growth in vivo and in vitro experiment as a pleiotrophic hormone being mitogenic, anti-apoptotic, pro-angiogenic, and pro-inflammatory in various cellular systems[69].The relationship between circulating leptin concentrations and colon cancer risk has been demonstrated [70].

The association between cigarette smoking and colon cancer risk has been shown to be dose-dependant [71]. cigarette smoking strongly associated with molecularly defined subtypes of colorectal cancer, such as MSI-high, CIMP-positive, and BRAF mutation-positive, that originate through epigenetically mediated carcinogenic pathways. These subtypes are more prevalent among women and are more often located in the proximal than the distal colon [72, 73]. For Samadder et al, cigarette smoking may be a stronger risk factor for KRAS mutation-negative tumors located in the proximal colon than in the distal colon [74]. Evidence is emerging in support of a strong association between smoking and proximal colon cancer, especially among female smokers [71].

We found in our study 266 localizations out of 366 registered cases. The sigmoid location ranked first with 37.59% (100cas / 266) followed by descending and ascending settlers representing 22.93% and 22.18% of the total proportion. Ele et al., Barth et al. And Brunet et al. Find frequencies approximating ours [37, 75, 76]. There is no clear explanation because there is little data available on the factors associated with the occurrence of colon cancer by sub-location [77].

Of the 366 cases reported, liberkhunien adenocarcinoma is the histological variety that dominates colon cancer in our series with 72.13% (264 cases). In parallel with the literature, this type of histology is the most important, but nevertheless Amegbor, Diallo Owono and Darré report higher percentages, especially 89.9%, 98% and 91.2% [13, 78, 79].

Lymphomas account for 46.5% (17 cases) of colonic malignancies, superior to data in the literature which states that they are rare with a rate of 2% [80]. They are observed at any age with great frequency between 40 and 60 years [66]; In our study non-Hodgkin's Malignant Lymphomas are observed at younger age groups (<20 years).

According to the literature, sarcomas are extremely rare histological varieties of primary colorectal cancers; Viguier et al reported a frequency of 0.09% [81]. In contrast to these studies, we find 22 cases of sarcoma including 21 of Kaposi (5.74% of the total proportion). Studies on etiological factors of sarcoma would allow us to explain and understand a high incidence of this histological type.

5. CONCLUSION

This study conducted on colon cancer in Cameroon has noted that that pathology has a high incidence of digestive tumors (26.01%) with male predominance. This pathology occurs mainly in people in their fifties with a relatively high frequency in people under 40 years. Predominant sigmoid localisation, they are primarily dominated by adenocarcinoma. However, these data remain relative given the absence of exhaustive information on several clinical and epidemiological aspects, hence the need for confirmation of these results by the installation of a national cancer registry. Besides, the fact that colon cancer cases are observed in young subjects invites us to conduct prospective studies on the genetic, environmental and food factors associated with this pathology

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

REFERENCES

1. World Health Organization. WHO classification of tumours of the digestive system. Fourth edition. Lyon; 2010
2. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. Mars 2015; 65(2):87–108.
3. Haute Autorité de Santé. Dépistage et prévention du cancer colorectal. Paris: Imprimerie nationale; 2013. 64 p.
4. Dernieres statistiques mondiales sur le cancer. IARC (International Agency for Research on Cancer. 2013 . Disponible sur: <http://www.iarc.fr>
5. Siegel R, DeSantis C, Jemal A. Colorectal cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(2):104–117.
6. Institut National de la Santé et de la recherche médicale. Épidémiologie France et monde: Lyon; Référentiel de pratique nationale; 2011. 70 p.
7. Welch HG, Robertson DJ. Colorectal Cancer on the Decline — Why Screening Can't Explain It All. *N Engl J Med*. 28 avr 2016;374(17):1605–7.
8. Dancourt V, Faivre J. Épidémiologie et dépistage des cancers colo-rectaux: *Cancers colo-rectaux*. *Rev Prat*. 2004;54(2):135–142.
9. Bouvier A-M, others. Épidémiologie descriptive du cancer colorectal en France. *BEH*. 2009;23:14–6.
10. Guérin S, Hill C. L'épidémiologie des cancers en France en 2010: comparaison avec les États-Unis. *Bull Cancer (Paris)*. 2010;97(1):47–54.

11. Ramji F, Cotterchio M, Manno M, Rabeneck L, Gallinger S. Association between subject factors and colorectal cancer screening participation in Ontario, Canada. *Cancer Detect Prev.* 2005;29(3):221–226.
12. Statistiques sur le cancer colorectal - Société canadienne du cancer. [Cité 9 juin 2017]. Disponible sur: <http://www.cancer.type/colorectal/statistics/>
13. Darré T, Amégbor K, Napo-Koura G, Bagny A, Bouglouga O, Lawson AL, et al. Profil histo-épidémiologique des cancers colorectaux au Togo. *J Afr Hépatogastroentérologie.* 1 déc 2014;8(4):226–9.
14. Chbani L, Hafid I, Berraho M, Mesbahi O, Nejari C, Amarti A. Aspects épidémiologiques et anatomopathologiques des cancers dans la région de Fès-Boulemane (Maroc). *East Mediterr Health J.* 2013;19(3):263.
15. Parkin DM, Bray F, Ferlay J, Jemal A. Cancer in Africa 2012. *Cancer Epidemiol Prev Biomark.* 1 juin 2014;23(6):953–66.
16. Umar A, Boland C, Terdiman J. Revised Bethesda Guidelines for hereditary nonpolyposis colorectal cancer (Lynch syndrome) and microsatellite instability. *J Natl Cancer Inst.* 2004;96(4):261–8.
17. Bonaïti-Pellié C, Eisinger F, Feingold J, Frébourg T, Grandjouan S, Lasset C, et al. Prédispositions héréditaires au cancer colorectal. *Gastroentérologie Clin Biol.* 2005;29(6–7):701–710.
18. Irabor DO. Colorectal Carcinoma: Why is there a lower incidence Nigerians when compared to Caucasiens? *J Cancer Epidemiol.* 2011;
19. Hagggar FA, Boushey RP. Colorectal Cancer Epidemiology: Incidence, Mortality, Survival, and Risk Factors. *Clin Colon Rectal Surg.* nov 2009;22(4):191–7.
20. Zeitoun J-D, Chrysostalis-Kulundzic A, Lefèvre J. Hépatogastro-entérologie, chirurgie digestive. Paris: Éd. Vernazobres-Grego; 2013.
21. Meddah D, Meddah B, Tir Touil A, Ghalek M, Sahraoui T. Étude épidémiologique du cancer du côlon chez des patients de l'Ouest algérien. *J Afr Cancer.* 2009;(1):31–5.
22. Gaudre N, Ly M, Badiaga Y, Dembele A, Bathily M, Kone A, et al. Particularités épidémiologiques et cliniques du cancer colorectal dans le service d'Hématologie Oncologie Médicale du point G de Bamako au Mali de 2005 à 2011: 113 cas. *Mali Med.* 2013;28(3):32–6.
23. Laurent-Puig P, Kirzin S. Comment la découverte d'une prédisposition familiale modifie la prise en charge des malades atteints par un cancer colorectal. *E-Mém Académie Natl Chir.* 2004;3(2):12–14.
24. Enow Oroch GE, Ndom P, Doh AS. Current cancer incidence and trends in Yaounde, Cameroon. *Oncol Gastroenterol Hepatol Rep.* 2012;1(1):58–63.
25. Engbang J-P, Fewou A, Moune A, Fonkwa C. Etude des caractéristiques épidémiologiques des cancers du colon au Cameroun: cas de la région du Littoral. XXIIIème journées scientifiques de la société camerounaise de gastro-entérologie; 2016 sept 22; Douala.
26. Peghini M, Rajaonarison P, Pecarrere J., Razafindramboa H, Richard J, Morin D. Épidémiologie des cancers du tube digestif à Madagascar. Apport de 14000 endoscopies effectués au centre hospitalier de Soavinandriana à ANTANANARIVO. *Médecine Afr Noire.* 1997;44(10):518–21.

27. Ayite A., Dosseh ED, Senah K, Etey k, James k, Napo-koura G. Epidémiologie descriptive des cancers du tube digestif au Togo. *J Afr Chir Dig.* 2001;1(0):10□1.
28. Harouna Y., Ilo A, Seybou A, Diakité I, Goza A. Les cancers colorectaux, Notre expérience à propos de 42 cas. *Médecine Afr Noire.* 2008;55(4):197□202.
29. Sani R, Dantata A., Bade M., Hassane N, Bazira L. Les cancers du tube digestif, Revue de 195 dossiers au service de Chirurgie digestive à l'Hopital National de Niamey Niger. *Médecine Afr Noire.* 2004;51(11):585□8.
30. Mamoudou G, Hami H, Soulaymani A, Quyou A. Les cancers digestifs au Niger.Fréquence relative sur une étude retrospective de 1992 à 2009. *Eur Sci J.* mars 2014;10(9).
31. Salamatou MG, Hinde H, Abdelmadjid S, Ali Q, Harouna MZ, Hassan N. les cancers digestifs au Niger. Fréquence relative sur une étude rétrospective de 1992 à 2009. *Eur Sci J.* 31 mars 2014 [cité 9 juin 2017]; Disponible sur: <http://les-cancers-digestifs-au-niger-frequence-relative-sur-une-etude-retrospective-de-1992-a-1999>
32. Bennon F, Sant M, Verdecchia. Cancer incidence in five continenets. *Int Agency Res Cancer.* 2002
33. Takongmo S, Essame-Oyono J., Binam F, Sadou, Malonga E. Les cancers colorectaux du sujet de moins de 40 ans à Yaoundé: des particularités anatomo-cliniques? *Médecine Afr Noire.* 2000;47(2).
34. Wabinga H, Parkin D, Wabwire-Mangen F, Namboze F. Trends in incidence in Kyadondo County, Uganda, 1960-1997. *Br J Cancer.* 82:1585□92.
35. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin.* mars 2015;65(2):87□108.
36. Ibou T, Kwame D, Cherif D, Abdou-Magib G, Chahir K. Morpho-epidemiological profile of elderly subject cancer in Dakar. *Ger Psychol Neuropsychiatr Vieil.* 2014;12(4):354□60.
37. Ele N, Okiemy G, Lebeau R, Nkoua-Mbon, J., Mbombi P, Massengo R. Le cancer du colon gauche au Chu de Brazzaville. Résultats du traitement chirurgical. *Mali Med.* 2006;21(1):1□4.
38. Amira A, Lilia K, Mahmoud B, Khiari M, Lahmer A, Gharbi L, et al. Etude épidémiologique, anatomopathologique et évaluation des facteurs pronostiques des adénocarcinomes colorectaux mucineux vs non mucineux. (A propos d'une série de 196 patients). *Tunis Med.* 88(01):12□7.
39. Wentink M, Rakers M, Stupart D, Algar U, Ramesar R, Goldberg P. Incidence and histological features of colorectal cancer in the Nothern Cape Province, South Africa. *SAJS.*2010; 48 (4):10-113
40. Cress R, Morris C, Ellison G, Goodman M. Secular changes in colorectal cancer incidence,stage at diagnosis and race/ethnicity 1992-2001. *Subsite Cancer.* 2006;1142□52.
41. Giovannucci E. Modifiable risk factors for colon cancer. *Gastroenterol Clin North Am.* 2002;(31):925□43.
42. Irabor D, Adedeji O a. Colorectal cancer in Nigeria: 40 years on. A review. *Eur J Cancer Care (Engl).* 1 mars 2009;18(2):110□5.
43. Sule A, Mandong B. Malignant colorectal tumours in patients 40 years and below: a review of 35 cases. *Cent Afr J Med.* 1999;45(8):209□12.

44. Bouregba S, Boulenouar FZ. cancer du colon [Internet]. 2015. Disponible sur: <http://dspace.univ-tlemcen.dz/handle/112/8305>
45. Chevreul K. Colorectal cancer in France. *Eur J Health Econ HEPAC Health Econ Prev Care*. janv 2010;10 Suppl 1:S15-20.
46. Yokane’Ndebi GE. Etude histopathologique des polypes rectocoliques en milieu hospitalier à Douala. [Douala]: Faculté de Médecine et des Sciences Pharmaceutiques de l’université de Douala; 2014.
47. Carr PR, Walter V, Brenner H, Hoffmeister M. Meat subtypes and their association with colorectal cancer: Systematic review and meta-analysis. *Int J Cancer*. 2016; 138(2):293-302.
48. Sugimura T, Wakabayashi K, Nakagama H, et al. Heterocyclic amines: mutagens/carcinogens produced during cooking of meat and fish. *Cancer Sci* 2004;95:290-9.
49. Bastide NM, Pierre FH, Corpet DE. Heme iron from meat and risk of colorectal cancer: a meta-analysis and a review of the mechanisms involved. *Cancer Prev Res (Phila)* 2011;4:177–84.
50. Joosen AMCP, Kuhnle GGC, Aspinall SM, et al. Effect of processed and red meat on endogenous nitrosation and DNA damage. *Carcinogenesis* 2009;30:1402–7.
51. Egeberg R, Olsen A, Christensen J, et al. Associations between red meat and risks for colon and rectal cancer depend on the type of red meat consumed. *J Nutr* 2013;143:464–72.
52. zur Hausen H. Red meat consumption and cancer: reasons to suspect involvement of bovine infectious factors in colorectal cancer. *Int J Cancer* 2012;130:2475–83.
53. Dupont I, Lucas D, Clot P, Menez C, Albano E. Cytochrome P4502E1 inducibility and hydroxy-ethyl radical formation among alcoholics. *J Hepatol* 1998; 28: 564-71.
54. Yang M, Coles BF, Delongchamps R, Lang NP, Kadlubar FF. Effects of the ADH3, CYP2E1, and GSTP1 genetic polymorphisms on their expressions in Caucasian lung tissue. *Lung Cancer* 2002; 38: 15-21
55. Wang Y, Duan H, Yang H, Lin J. A pooled analysis of alcohol intake and colorectal cancer. *Int J Clin Exp Med* 2015;8(5):6878-6889
56. Manautou JE, Carlson GP. Ethanol-induced fatty acid ethyl ester formation in vivo and in vitro in rat lung. *Toxicology* 1991; 70: 303-12.
57. Homann N, Tillonen J, Salaspuro M. Microbially produced acetaldehyde from ethanol may increase the risk of colon cancer via folate deficiency. *Int J Cancer* 2000; 86: 169-73.
58. Choi SW, Stickel F, Baik HW, Kim YI, Seitz HK, Mason JB. Chronic alcohol consumption induces genomic but not p53-specific DNA hypomethylation in rat colon. *J Nutr* 1999; 129: 1945-50.
59. Xue K, Li FF, Chen YW, et al. Body mass index and the risk of cancer in women compared with men: a meta-analysis of prospective cohort studies (link is external). *Eur J Cancer Prev*. 2017 Jan;26(1):94-105.
60. Ning Y, Wang L, Giovannucci EL. A quantitative analysis of body mass index and colorectal cancer: findings from 56 observational studies (link is external). *Obes Rev*. 2010 Jan; 11(1):19-30.
61. Ma Y, Yang Y, Wang F, Zhang P, Shi C, Zou Y, Qin H. Obesity and risk of colorectal cancer: a systematic review of prospective studies. *PLoS One*. 2013;8(1):e53916.

62. Bergstrom A, Pisani P, Tenet V, Wolk A, Adami HO (2001) Overweight as an avoidable cause of cancer in Europe. *Int J Cancer* 91: 421–430.
63. Clayton PE, Banerjee I, Murray PG, Renehan AG (2011) Growth hormone, the insulin-like growth factor axis, insulin and cancer risk. *Nat Rev Endocrinol* 7:11–24.
64. Dong Y, Zhou J, ZhuY, He T, Hu H, et al. Abdominal Obesity and Colorectal Cancer Risk: Systematic Review and Meta-Analysis of Prospective Studies. *Biosci Rep*. 2017. pii: BSR20170945. doi: 10.1042/BSR20170945
65. Chung Y.W.; Han D.S.; Park Y.K.; Son B.K.; Paik C.H.; Lee H.L.; Jeon Y.C.; Sohn J.H. Association of obesity, serum glucose and lipids with the risk of advanced colorectal adenoma and cancer: a case-control study in Korea. *Dig Liver Dis*. 2006 38:668-72.
66. Amemori S.; Ootani A.; Aoki S.; Fujise T.; Shimoda R.; Kakimoto T.; Shiraishi R.; Sakata Y.; Tsunada S.; Iwakiri R.; Fujimoto K. Adipocytes and preadipocytes promote the proliferation of colon cancer cells in vitro. *Am J Physiol Gastrointest Liver Physiol*. 2007 292:G923-9.
67. Ogino S.; Kawasaki T.; Ogawa A.; Kirkner G.J.; Loda M.; Fuchs C.S. Fatty acid synthase overexpression in colorectal cancer is associated with microsatellite instability, independent of CpG island methylator phenotype. *HUM PATHOL*. 2007 38:842-9.
68. Renehan A.G.; Roberts D.L.; Dive C. Obesity and cancer: pathophysiological and biological mechanisms. *Arch Physiol Biochem*. 2008 114:71-83.
69. Stattin P.; Lukanova A.; Biessy C.; Soderberg S.; Palmqvist R.; Kaaks R.; Olsson T.; Jellum E. Obesity and colon cancer: does leptin provide a link? *INT J CANCER*. 2004 109:149-52.
70. Stattin P.; Palmqvist R.; Soderberg S.; Biessy C.; Ardnor B.; Hallmans G.; Kaaks R.; Olsson T. Plasma leptin and colorectal cancer risk: a prospective study in Northern Sweden. *ONCOL REP*. 2003 10:2015-21.
71. Parajuli R, Bjerkaas E, Tverdal A, Selmer R, Le Marchand L, et al. The increased risk of colon cancer due to cigarette smoking may be greater in women than men. *Cancer Epidemiol Biomarkers Prev*. 2013 May;22(5):862-71
72. Boland CR, Goel A. Clearing the air on smoking and colorectal cancer. *J Natl Cancer Inst* 2010;102:996–7.
73. Limsui D, Vierkant RA, Tillmans LS, Wang AH, Weisenberger DJ, Laird PW, et al. Cigarette smoking and colorectal cancer risk by molecularly defined subtypes. *J Natl Cancer Inst* 2010;102:1012–22.
74. Samadder NJ, Vierkant RA, Tillmans LS, Wang AH, Lynch CF, Anderson KE, et al. Cigarette smoking and colorectal cancer risk by KRAS mutation status among older women. *Am J Gastroenterol* 2012;107:782–9.
75. Barth X, Lanoviron A, Repellin P, Spay G. Les occlusions aiguës par cancer colique, analyse d’une série de 163 observations. *Lyon Chir*. 1990;86(1):12–7.
76. Brunet C, Thirion Z, Gregoire R, Farisse J. Occlusions par cancers coliques: traitement en urgence (62cas). *J Chir*. 1995;32:30–3.
77. Lelong B, Moutardier V, Delpere J. Prise en charge des tumeurs primitives colorectales. *Rev Prat*. 2004;54:155–66.
78. Diallo F, Nguema R, Ibaba J. Epidemiological and diagnostic features of colorectal cancer in libreville. *Gabon Med TROP*. 6:605–7.

- 493 79. Bagny A, Bouglouga O, Darre T, Lawson-Ananissoh LM, Kaaga YL, Sonhayé L, et
494 al. Profil épidémiologique et diagnostique des cancers digestifs au CHU Campus de
495 Lomé: à propos de 250 cas. J Afr Hépato-Gastroentérologie. 2015;9(2):80–84.
496 80. Ruskone-Founestraux A, Lavergne-Slove A, Delmer A. Lymphomes gastro-
497 intestinaux. Gastroenterol Clin Biol. 26:233–41.
498 81. Viguier J, Bourlier P, Karsenti D. Cancer du colon. In: Encycl méd chir, Gastro-
499 entérologie. 2003. p. 18.
500