

Assessment of genetic variability and correlation coefficient to improve some agronomic traits in rice

ABSTRACT

This study was carried out to evaluate genetic components, heritability and correlation coefficient effects of the parents and their F1 for some agronomic as well as yield and yield traits. A line x tester cross was conducted among the ten parents (four lines and six testers) in 2010 to produce (24) crosses. The results indicated that GCA showed highly significant for all studied traits. The mean squares of lines x testers interaction (SCA) exhibited highly significant differences for all studied traits. Higher estimates for non-additive or dominance genetic variance was computed for number of panicles / plant, number of filled grains/ panicle and grain yield /plantin comparison with its corresponding evaluates of additive genetic variance. Concerning to heritability estimates, high values were observed in broad sense heritability. Meanwhile,evaluate of narrow-sense heritability were low to moderate for number of panicles / plant, number of filled grains / panicle, 1000-grain weight and grain yield / plant. Heritability estimates in narrow sense were relatively low for panicle weight and spikelet fertility percentage. As for correlation for grain yield/ plant the results recorded highly positive correlation coefficient with each of days number to heading, plant height, number of panicles / plant, panicle length and fertility percentage.

Keywords: Heritability, Correlation, Genetic parameters, Yield, Rice.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is central to the lives of billions of people around the world. Rice provides 27% of dietary energy and 20% of dietary protein in the developing world. Rice is cultivated in at least 114, mostly developing countries and is the primary source of income and employment for more than 100 million households in Asia and Africa (FAO, 2004). Rice crop plays a significant role in Egypt, as strategic crop for sustaining the food self-sufficiency and for increasing the export., The total rice production in Egypt reached 7.1 million tons with a national average of 9.9 tons/ha in 2012 growing season. This average ranked the first among the rice producing countries in the world (RRTC, 2013).

Genetic parameters such as genotypic coefficient of variation and phenotypic coefficient of variation are useful in detecting the amount of variability present in the germplasm. Heritability coupled with high genetic advance helps in determining the influence environment on the expression the genotypic and reliability of characters (Nirmala et al., 2015). Yield traits breeding for grain yield increasing would be most effective, if the traits involved are highly heritable and genetically independent or positively correlated with grain yield. However, it is very difficult to judge whether observed variability is highly heritable or not (Osman et al., 2012). Moreover, knowledge of heritability is important for selection based improvement as it indicates the extent of transmissibility of a character into future generations (Sabesan et al., 2009). The utilization of heritability and genetic advance of yield traits and inferences from significant genotypic correlation between yield and its components should permit selection of predictable rice genotypes for upland ecosystem (Osman et al., 2012).

Productivity improvement for rice has become of immense necessary to feed nearly half of the world's population. Genetic resource of rice is the primary material for improving in rice (Perera et al., 2014). Over dominance pseudo-over dominance is the most important contributor to heterosis of yield, number of grains /panicle and grain weight. Dominance x dominance interaction is important for heterosis of tillers / plant and grain weight and has roles in yield and grain number. Cumulative effects of these components may adequately explain the genetic basis of heterosis in the hybrid (Zhou Gang et al., 2012). With the above background information this study aimed to study the magnitude of the genetic variance components i.e. additive genetic variance (σ^2_A), dominance genetic variance (σ^2_D) and broad and narrow sense heritability for agronomic characteristics and yield as well as to estimate the correlation between studied traits.

2. MATERIAL AND METHODS

The the present study was carried out at the Experimental Farm of the Rice Research and Training Center (RRTC), Sakha, Kafr EL-Sheikh, Egypt, at 2010 and 2011 successive seasons. The research involving four rice genotypes, i.e. three commercial varieties (Giza178, Sakha103 and Sakha106), one promising line (Gz 9057-6-1-3-2) as well as six bold grain lines (SKC 23819-189-1-1-1-3-1-2-4-2, SKC 23819-192-2-1-2-2-4-2-1-2, SKC 23819-192-2-1-2-3-1-1-1-2, SKC 23819-192-2-1-2-4-5-3-2-1, SKC 23819-192-2-2-1-1-2-1-2-2 and SKC 23819-192-2-2-1-1-2-2-1-2). The parental genotypes have a wide range of variations for all studied traits. Origin and parentage of the ten parents were illustrated in (Table 1).

Table 1: Names, origin and parentage of the parents utilized in this study

No.	Entry	Origin	Parentage
1	Giza 178	Egyptian	(Giza175/Milyang49)
2	Sakha 103	Egyptian	(Giza 177/Suweon 349)
3	Sakha 106	Egyptian	(Giza 177/Hexi 30)
4	GZ9057-6-1-3-2 (Giza 179)	Egyptian	(Gz 6296/Gz 1368-S-5-4)
5	SKC 23819-189-1-1-1-3-1-2-4-	America/ Egypt	(L 204/Giza177)
6	SKC 23819-192-2-1-2-2-4-2-1-	Egypt /America	(L 204/Giza177)
7	SKC 23819-192-2-1-2-3-1-1-1-	Egypt /America	(L 204/Giza177)
8	SKC 23819-192-2-1-2-4-5-3-2-	Egypt/America	(L 204/Giza177)
9	SKC 23819-192-2-2-1-1-2-1-2-	Egypt/America	(L 204/Giza177)
10	SKC 23819-192-2-2-1-1-2-2-1-	Egypt/America	(L 204/Giza177)

The ten parental genotypes in this study were sown in the growing season of 2010 in three sowing dates with 15 days intervals to overcome the difference of heading date among the parental varieties. After 30 days from the sowing, the seedling of ten parents were transplanted to the experimental field in three rows, each row was five meters long and (20 cm x 20 cm) spacing between plants and rows. A line x tester cross was conducted among the ten parents (four lines and six testers) in 2010 to produce (24) crosses. The hybridization technique of Jodon (1938) and modified by Butany (1961) was used and the hot water method of emasculation was utilized. The parental varieties and their resulting 24 crosses were arranged in a randomized complete block design (RCBD) with three replications. Each replication contained 34 rows and each row contains 25 individual plants for each genotype. The studied traits were number of days to heading (day), plant height (cm), number of panicles/ plant, number of filled grains/ panicle, panicle length, panicle weight (g), spikelets fertility percentage, grain weight (g), and grain yield / plant(g).

Genotypes means were used for the analysis of variance Singh and Chaudhary (1985). The correlation coefficients were tested using the formula according to Gomez and Gomez (1983). The data were subjected to analysis of variance as suggested by Panse and Sukhatme (1957) and the analysis of variance for line x tester crossing followed the design of Kempthorne (1957).

3. RESULTS AND DISCUSSION

3.1 Analysis of variance

The results on analysis of variance revealed that highly significant differences were observed among genotypes for all studied traits (Table 2). The parental lines and the crosses showed highly significant differences for all studied characters. Parents vs. crosses mean squares indicated that average heterosis was highly significant for all traits. Obtained result agreement with Singhet *al.* (2011) and Abdelkhalik *et al.* (2013).

Table 2: Analysis of variance and mean square from line x testers' analysis for rice studied traits

S.O.V	D.F	Days to heading(days)	Plant height (cm)	No. of panicles plant ⁻¹	No .of filled grains panicle ⁻¹	Panicle length (cm)
Reps.	2	0.992	0.381	0.191	26.90	0.123
Genotypes	33	74.09**	303.7**	96.02**	2600**	9.753**
Parents	9	121.6**	379.3**	24.01**	515.3**	9.769**
P.Vs.C	1	706.6**	2839.2**	481.6**	0.308**	0.470**
Crosses	23	644.0**	163.9**	107.4**	3530**	10.15**
Lines (gca,L)	3	320.1**	806.4**	721.7**	6216**	64.22**
Testers(gca,T)	5	90.52**	60.68**	5.840**	4459**	2.040**
Line x testers(sca)	15	233.4**	69.93**	18.45**	2682**	2.041**
Error	66	21.74	0.268	0.496	15.99	0.215

*and**, Significant at 5% and 1% levels of probability, respectively.

Table 2: continue.

S.O.V	D.F	Panicle weight	Spikelets fertility%	1000-grain weight (g)	Grain yield plant ⁻¹ (g)
Reps.	2	0.024	0.281	0.003	1.569
Genotypes	33	0.503**	610.1**	43.41**	272.0**
Parents	9	0.312**	25.25**	125.3**	174.2**
P.Vs.C	1	0.425**	936.0**	37.83**	2478**
Crosses	23	0.581**	824.8**	11.59**	214.3**
Lines (gca,L)	3	1.434**	3033**	49.97**	376.0**
Testers (gca,T)	5	0.275**	815.6**	13.62**	234.3**
Line x testers (sca)	15	0.513**	386.2**	3.249**	175.3**
Error	66	0.019	0.964	0.032	10.45

*and**, Significant at 5% and 1% levels of probability, respectively.

The results also illustrated that the general combining ability variance of lines (female parents) showed highly significant for all studied characters. The general combining ability variances of testers (male parents) were highly significant for all characters. The mean squares of lines x testers interaction (SCA) exhibited highly significant differences for all characters, this might be due to the wide genetic diversity of lines and testers. These results were agreement with Singhet *et al.* (2011), Abdelkhaliket *et al.* (2013) and Anis *et al.* (2016b).

3.2 Genetic components of variance and heritability

Results in (Table 3) showed that the additive genetic variance as a portion of the total genetic variance was smaller than the non-additive or dominance genetic variance for all studied characteristics except for number of panicles/ plant, number of filled grains / panicle and panicle length characters. This result indicated that dominance genetic variance played the major role compared to the additive variance in the inheritance of these traits and phenotypic variance for these traits was due to dominance genetic variance and environmental effects.(Fazza *et al.* 2016)) These findings led to conclusions the selection for such traits must be done in the late segregating generations. This result agreed with Zhou Gang *et al.* (2012), Anis (2013) and Perera *et al.* (2013).

Both additive and dominance genetic variances (σ^2_A and σ^2_D) were positive for all traits. Their respective values for σ^2_A and σ^2_D were (1.167 and 5.076) for days to heading, (8.324 and 23.22) for plant height, (7.374 and 5.983) for number of panicles plant⁻¹, (96.58 and 88.9) for number of filled grains panicle⁻¹, (0.675 and 0.608) for panicle length,(0.01 and 0.164) for panicle weight, (39.45 and 128.4) for spiklets fertility percentage, (0.707 and 1.072) for 1000-grain weight at last (4.992 and 54.95) for grain yield / plant, respectively. These results indicate that the two genetic variance components might be

important in the inheritance of these traits. These results were in general agreement with those reported by Anis (2013) and Perera *et al.* (2013).

Table 3: Estimation of additive genetic variance (σ^2A), dominance genetic variance (σ^2D), environmental variance (σ^2E) and heritability in broad and narrow senses for studied characters

Traits	σ^2A	σ^2D	σ^2E	Heritability %	
				$h^2_b\%$	$h^2_n\%$
Days to heading (day)	1.167	5.076	0.329	94.98	17.76
Plant height (cm)	8.324	23.22	0.268	99.15	26.16
No. of panicles plant ⁻¹	7.374	5.983	0.496	96.42	53.22
No. of filled grains panicle ⁻¹	96.58	88.9	15.99	98.40	9.643
Panicle length (cm)	0.675	0.608	0.215	85.64	45.06
Panicle weight	0.010	0.164	0.019	90.12	5.607
Spikelets fertility%	39.45	128.4	0.964	99.42	23.37
1000-grain weight (g)	0.707	1.072	0.963	98.23	39.06
Grain yield plant ⁻¹ (g)	4.992	54.95	10.45	85.15	7.090

Concerning to heritability estimates in broad sense ($h^2_b\%$), the results indicated that the heritability values were more than 90% for all studied traits except for panicle length and grain yield plant⁻¹ the values were 85.64% and 85.15% , respectively (Table 3). On the other hand, heritability estimates in the narrow sense ($h^2_n\%$) were low to moderate for the same traits. These results indicated that a major part of the total genotypic variance was non-additive in nature for these traits. These results were in general agreement with those reported by Anil and Mani (2010), Fahlani *et al.* (2010), Kayande and Debaje (2010), and Mohan and Chauhan (2011) and Anis *et al.*, (2016 a).

Heritability in narrow sense is an indicator of the efficiency of selection procedure in identifying the superior genotypes. In (Table 3) it was observed that heritability estimates in broad sense were high for all traits, while heritability estimates in narrow sense were relatively low for days to heading, plant height, number of filled grains panicle⁻¹, panicle weight, spikelets fertility percentage, and grain yield plant⁻¹, and moderate for panicle length and 1000-grain weight. These results are in full agreement with previous results obtained by Chakraborty and Chakraborty (2010) and Adilakshmi and Rani (2012).

3.3 Correlation coefficient

The study of correlation coefficients among various agronomic traits is of great importance, since correlation can enable researchers to predict the performance of complex and quantitative traits that is difficult to measure by using linked simple measured traits. Thus, enhancing significantly the selection efficiency since the influence of environment will be minimal in case of the simple traits (Anis *et al.* 2016a). The estimates of correlation coefficients among all studied characters are presented in Table 4.

Table 4: Estimates of phenotypic correlation coefficients among each pair of studied traits

Traits	Heading date	Plant height	No. of panicles plant ⁻¹	No. of filled grains Panicle ⁻¹	Panicle length	Panicle weight	Spikelets fertility%	1000-grain weight	Grain yield plant ⁻¹
Heading date	0.00								
Plant height	0.142	0.00							
No. of panicles plant ⁻¹	0.479**	0.580**	0.00						
No. of filled grains Panicle ⁻¹	0.002	0.215	0.401**	0.00					

Panicle length	-0.007	0.492**	0.278	0.169	0.00				
Panicle weight	0.027	-0.338*	-0.236	0.253	-0.022	0.00			
Spikelets fertility%	-0.234**	-0.369*	-0.179	0.497**	-0.476**	0.356*	0.00		
1000-grain weight	0.436**	-0.535**	0.015	-0.103	-0.319	0.417**	0.201	0.00	
Grain yield plant ⁻¹	0.253**	0.206**	0.181**	0.080	0.020**	0.230	0.151**	-0.133	0.00

130

131 Concerning to heading date, data showed that highly significant positive correlation with each of
 132 number of panicles/ plant, 1000-grain weight (g) and grain yield/ plant, (0.479, 0.436 and 0.253,
 133 respectively) and highly significant negative correlation with spikelets fertility percentage (-0.234). These
 134 results are in similar to those of Anis (2009) and Minnie *et al.* (2013).

135 As for plant height gave highly significant positive correlation coefficient with each of number of
 136 panicles / plant (0.580), panicle length (0.492) and grain yield/ plant (0.206). Also it was found highly
 137 significant negative correlation between plant height and 1000-grain weight (-0.535) while found
 138 significant negative correlation with panicle weight (-0.338) and fertility percentage (-0.369). This result is
 139 in accordance with to those of El-Sherif (2011) and Essa (2012).

140 With regard to number of panicles/ plant highly significant positive correlation coefficient was
 141 showed with number of filled grains panicle⁻¹ (0.401), and grain yield plant⁻¹ (0.181). The obtained results
 142 are in harmony with those reported by El-Sherif (2011). Regarding number of filled grains /panicle, data
 143 cleared highly significant positive correlation coefficient with fertility percentage the value was (0.497)
 144 similar results were previously obtained by Nandeshware *et al.* (2010) and Fazza *et al.* (2016a,b).

145 Significant and positive correlation coefficient was recorded between panicle length and grain yield/
 146 plant with value of 0.020, and highly significant negative correlation with spikelet fertility percentage with
 147 value of -0.476. The obtained results are in harmony with those reported by Chakraborty and Chakraborty
 148 (2010) ;Fazaa *et al.*, (2016a, b) and Aniset *et al.*, (2016 b). On the other hand, panicle weight showed
 149 positive highly significant correlation coefficient with 1000- grain weight and the value was (0.417) and
 150 significant positive correlation coefficient with spikelet fertility percentage with value (0.356). The result is
 151 in agreement with Nandeshware *et al.* (2010) and Essa (2012). Highly significant positive correlation
 152 coefficient for fertility percentage was observed with grain yield plant⁻¹ (0.151). Similar results were
 153 previously obtained in different crops by Sedeek *et al.* (2007); Abd el-wahed *et al.*, (2015) and Barutçular
 154 *et al.*, (2016).

155

156

157 4. CONCLUSION

158

159 The results of the present study suggested that the hybrid combinations viz., Giza 177 x Chinese2
 160 followed by the hybrid, Giza 177 x Sakha 106, Chinese2 x Gz8479-6-2-3-1 and Sakha 101 x IR67701B
 161 were existing rice hybrids due to good heterosis values for grain yield trait. Thus these genotypes with
 162 higher magnitude of this trait could be either evolved by breeding program for genetic improvement of
 163 yield in rice.

164

165

166 REFERENCES

167

168 Abdlekhaliq AF, El kallawy WH, Abd El-Lattef ASM, Anis GB. 2013. Estimation of genetic components,
 169 heritability and cluster analysis for some agronomic characters in rice. J. Plant Production, Mansoura
 170 Univ. 2013; 4(9): 1297-1307.

171

172 Abd el-wahed MH, EL Sabagh A, Zayed A, Ahmed S, Saneoka H, Barutçular C. Improving yield and
 173 water productivity of maize grown under deficit-irrigated in dry area conditions. Azarian Journal of
 Agriculture. 2015; 2 (5), PP: 123-132.

- 174 Adilakshmi D,Rani MG.Variability, character association and path analysis in rice varieties under
175 submergence. Crop Research (Hisar).2012; 44(1/2):146-151.
- 176 Anil, K., Mani, S.C. (2010). Gene action for grain yield, its components and quality traits in Basmati rice
177 (Oryza sativa L.). Pantnagar J. Res., 8(1):26-31.
- 178 Anis, G.B. (2009).Breeding for earliness and some agronomic characters in rice (Oryza sativa L.).An
179 M.Sc. Thesis, Fac. of Agric. Kafr El-Sheikh Univ., Egypt.
- 180 Anis GB.Development and evaluation of new restorer and maintainer lines and their relationship to
181 heterosis in rice. A Ph. D.2013; Thesis, Agric. Sci., Genetics Dept., Mansoura Univ., Egypt.
- 182 Anis G, El Sabagh A, El-Badry A, Barutçular C.Improving Grain Yield in Rice (Oryza Sativa L.) by
183 Estimation of Heterosis, Genetic Components and Correlation Coefficient. International Journal of Current
184 Research.2016a; Vol. 8(01):25080-25085.
- 185 Anis G, ELSabagh A, Ghareb A, EL-Rewainy I. Evaluation of promising lines in rice (Oryza sativa L.) to
186 agronomic and genetic performance under Egyptian conditions. International Journal of Agronomy and
187 Agricultural Research.2016b; 8(3), 52-57.
- 188 Barutçular C, Yıldırım M, Koç M, Akıncı C, Toptaş I, Albayrak O, Tanrikulu A, EL Sabagh A. Evaluation of
189 SPAD chlorophyll in spring wheat genotypes under different environments.Fresenius Environmental
190 Bulletin.2016; 25(4):1258-1266.
- 191 Butany WT. Mass emasculation in rice. Inter. Rice Com. Newsletter.1961; 9: 9-13.
- 192 Chakraborty R, Chakraborty S. Genetic variability and correlation of some morphometric traits with grain
193 yield in bold grained rice (Oryza sativa L.) gene pool of Barak valley. American-Eurasian J. Sustainable
194 Agric.2010; 4(1):26-29.
- 195 El-Sherif AIAZ. Genetic studies on blast disease in rice (Oryza sativa L.). M.Sc. Thesis, Fac. Agric.
196 Menofiya Univ.Shibin El-Kom, Egypt.2011.
- 197 Essa WMMM. Genetic and molecular studies on blast resistance in rice (Oryza sativa L.). M.Sc. Thesis,
198 Fac. of Agric., Kafr El-Sheikh Univ., Egypt.2012.
- 199 Fazaa M, EL Sabagh A, Anis G, EL-Rewainy I, Barutçular C, Hatipoğlu R, Islam MS. The Agronomical
200 Performances of Doubled Haploid Lines of Rice (Oryza sativa L.)Derived from Anther Culture. Journal of
201 Agriculture Science.2016a; Vol. 8(5): 177-183.
- 202 Fazaa M, EL Sabagh A, Anis G, EL-Rewainy I, Barutçular C, Yildirim M,Islam MS.. (2016b).Grain Quality
203 of Doubled Haploid Lines in Rice (Oryza sativa L.)Produced by Anther Culture.Vol. 2016b; 8(5): 184.190.
- 204 Fahlani RA, Khodambashi M, Houshmand S, Arzani A. Estimation of the heritability of agro-
205 morphological traits in rice (Oryza sativa L.) using F2:3 families. African J. Agric. Res.2010; 5(11): 1297-
206 1303.
- 207 Food and Agriculture Organization (FAO).Rice is Life. Italy: FAO.
208 [http://www.fao.org/newsroom/en/focus/20043_6887/](http://www.fao.org/newsroom/en/focus/20043_6887/index.html) index.html.International Year of Rice. 2004. "Rice
209 and human nutrition"
- 210 Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. 2nd Edition. John Wiley Sons,
211 NY, USA.1983.
- 212 Jodon NE. Experiments on artificial hybridization of rice. J. Mer. Soc. Agron.1938; 30: 249-305.

- 213 Kayande NV, Debaje PP. Genetic variation and association studies in segregating population of rice
214 (*Oryza sativa* L.). *Inter. J. Plant Sci. (Muzaffarnagar)*.2010; 5(2): 700-702.
- 215 Kempthorne O. An introduction to genetic statistics. John Wiley and Sons Inc., New York.1957.
- 216 Khush, G.S. Strategies for increasing the yield potential of cereals: case of rice as an example. *Plant*
217 *Breeding*.2013; 132(5): 433-436.
- 218 Minnie CM, Reddy TD, Raju CS. Correlation and path analysis for yield and its components in rice (*Oryza*
219 *sativa* L.). *J. Res. ANGRAU*.2013; 41(1): 132-134.
- 220 Mohan L, Chauhan DK. Studies of genetic variability, heritability and genetic advance in relation to yield
221 traits in rice. *Agric. Scie. Digest*.2011; 31(3): 220-222.
- 222 Nandeshwar BC, Pal S, Senapati BK, De DK. Genetic variability and character association among
223 biometrical traits in F2 generation of some rice crosses. *Electronic J. Plant Breed*.2010; 1(4): 758-763.
- 224 Osman KA, Mustafa AM, Ali F, Yonglain Z, Fazhan Q. Genetic variability for yield and related attributes of
225 upland rice genotypes in semi-arid zone (Sudan). *Afr. J. Agric. Res*.2012; 7(33):4613-4619.
- 226 Panse VG, Sukhatme PV. Statistical methods for agricultural workers. Indian Council Agric. Res.1957;
227 New Delhi, India.
- 228 Perera UIP, Bentota AP, Ratnasekara D, Senanayake SGJN. Heterosis in F1 generation of two Indica rice
229 crosses for growth and yield characteristics. *J. Agric. Sci. (Sri Lanka)*.2013; 8(3): 136-141.
- 230 Perera UIP, DisnaRatnasekera, Senanayake SGJN, Abey Siriwardena DS de Z. Genetic Parameters and
231 Correlations of Yield Attributing Characteristics of Weedy Rice in Sri Lanka. *JNSF, Sri Lanka*.2014; 42 (4):
232 299-306.
- 233 RRTC. 2013. National Rice Research Program: Final results of 2012 growing season. Sakha, Egypt.
- 234 Sabesan T, Suresh R, Saravanan K .Genetic variability and correlation for yield and grain quality
235 characters of rice grown in coastal saline lowland of Tamiluadu. *Electr. J. Plant Breed* .2009; 1:56-59.
- 236 Sedeek SEM, Hammoud SAA, Ammar MH, Metwally TF. Genetic variability, heritability, genetic advance
237 and cluster analysis for some physiological traits as well as grain yield and its components in rice (*Oryza*
238 *sativa* L.). *J. Agric. Res., Kafr El-Sheikh Univ*.2007; 35(3): 858-878.
- 239 Singh AK, Chaudhary BD. Biometrical Methods in Quantitative Genetic Analysis. 1985; Kalyani Publisher,
240 New Dehli, India.
- 241 Singh SK, Singh CM, Lal GM. Assessment of genetic variability for yield and its component characters in
242 rice (*Oryza sativa* L.). *Res. Plant Biol*.2011; 1(4):73-76.
- 243 Zhou G, Chen Y, Yao W, Zhang C., Xie W, Hua J, Xing Y, Xiao J, Zhang Q. Genetic composition of yield
244 heterosis in an elite rice hybrid. *Proceedings of the Nati. Acad. Sci. Uni. Sta. Ame*.2012; 109(39): 15847-
245 15852.