

Original Research Article

THE EFFICACY OF *ZINGIBER OFFICINALE* (GINGER-ZINGIBERACEAE) CRUDE EXTRACTS APPLIED AS INDIVIDUAL AND MIXED (WITH *DENNETTIA TRIPETALA* (PEPPERFRUIT-ANNONACEAE)) AGAINST *MUSCA DOMESTICA* (HOUSEFLY) LARVAE.

ABSTRACT

Musca domestica is one of the most common household insect pests, it feeds on human food thereby transmitting disease to man and livestock since it is a vector of many diseases. Crude extracts of *Ziniger officinale* was applied alone and mixed with *Dennettia tripetala* in equal proportion at various concentrations (200ppm to 1000ppm) to determine their toxic efficacy against freshly emerged larvae of *Musca domestica*. Each concentration was replicated twice. The bioassay was done using dipping and feeding techniques under laboratory conditions and was monitored every 12 hours for 72 hours. Distill water was used as control for the experiment. Mean mortality was recorded to obtain Median Lethal Concentration (LC₅₀) and Median Lethal Time (LT₅₀). The results showed that mortality was generally significant (P<0.05) as concentration increased from 200ppm to 1000ppm,. Mortality was not significant with time (P>0.05) as duration of exposure increased from 12-72 hours when *Zingiber officinale* was applied alone, but when it (*Z. officinale*) was mixed with *D. tripetala* in equal proportion mortality was significant (P<0.05) with time as duration of exposure increased from 12-72 hours. The mixture of plant extracts (*Z. officinale* and *D. tripetala* in equal proportion) had the lowest LC₅₀ and LT₅₀ (0.2754 mg/ml (275.4ppm) and 30 hours). It was observed that dipping the insects in plant extracts caused more mortality when compared to feeding the insects with plant extracts. This result showed that plant extracts used for this study could be promising bioinsecticides especially when mixed in equal proportions and should be considered in integrated pest management (IPM).

Key words: *Zingiber officinale*, *Dennettia tripetala*, *Musca domestica*, LC₅₀ and LT₅₀.

INTRODUCTION

Housefly is one of the most common household insect pests, it feeds on human food thereby transmitting disease to man and livestock since it is a vector of many diseases such as typhoid, cholera, dysentery, salmonella, etc. they also transmit the eggs of parasitic worms hence houseflies are seen as carries of easily communicable disease. Diseases are found on the legs and

hairs covering the body of houseflies. They come in contact with these pathogens when females lay their eggs on decaying organic matter. Houseflies transmit these pathogens to food substance or any surface they come in contact with within seconds. They are able to transmit these diseases because they liquefy solid food before feeding on them as they have sucking mouthparts and people get infested when they feed on it. Their presence in homes (around man) is irritating as they are considered as unhygienic insects. They tend to disturb men and livestock during leisure and work periods. Eliminating this insect pest totally maybe difficult, reducing its population below economic threshold and stopping its contact with human and livestock food and water is very necessary and a step to make homes and environment hygienic. Chemical insecticides which are usually used in controlling housefly (Herve *et al.*,) have great adverse effects on the ecosystem at large, as their residues left in the environment are hazardous. Chemical insecticides also make the insect pest become resistant (Scott, *et al.*, 2000; Shono and Scott 2003). Thus researchers are in continuous search of alternative methods for controlling insect pests. Plants have been used in medicine even before man understood the cause of infectious disease (El-Sherbini and Hanykamel 2014) this has led to the use of bio insecticides in controlling insect pest because they are better alternatives to chemical insecticides. *Zingiber officinale* is a flowering plant in the family Zingiberaceae whose rhizome or simply ginger is widely used as spice. It has gingerols and shogaol as its active constituents amongst other compounds (Gosh, 2011) which suffocate insects. *Dennettia tripetala* (pepper fruit) is an indigenous medicinal fruit tree of the family Annonaceae known to have β -phenylnitroethane (a nitro compound) as the compound responsible for its high insecticidal activity (Anyaele and Amusan, 2003). There are various works on the use of bioinsecticides in controlling housefly such the work of Asid *et al.* (2015), Soonwera (2015), Meenakshisundaram *et al.*, (2014)) but less research has been done on the comparison of plant extracts applied individually and mixed in equal proportions. Thus this work will be evaluating the efficacy of *Zingiber officinale* applied individually and mixed with *Dennttia tripetala* in equal proportions; their insecticidal strength will also be compared using dipping and feeding techniques.

MATERIALS AND METHODS

Insect collection and culture

Insects were collected near a garbage site by placing fresh fish in a small plastic plate inside the rearing cages. The fish served as a culture medium for laying eggs and larvae development. The cage (30x30x30xm) was built with wood, nets and wire gauze with a wooden door by its side, the floor was made with plywood. Freshly emerged larvae were used for the bioassay. The cage was placed on plastic stands which contained engine oil to prevent infestation of other insects and mites (Herve' *et al.*, 2008). The larvae was transferred into petri dishes containing larvae food (2ml of powdered milk mixed with distilled water) using fine art brush. This procedure was adopted from Herve *et al.*, (2008); Bosly (2013) and Ojianwuna *et al.*, (2011) with some modifications.

Preparation of test plant materials

The plants used for this study was purchased from a local market in Ukwuani Local Government area of Delta State and identified by a plant taxonomist in the department of botany Delta state university, Abraka. *Zingiber officinale* rhizome was sliced into tiny pieces while *Dennettia tripetala* skin was peeled off to obtain the seeds which were used for this study. They (*Z. officinale* and *D. tripetala*) were dried at ambient temperature ($28\pm 2^{\circ}\text{C}$) for four weeks, afterwards they were grounded with an electric blender (Philip) and sieved with 0.1mm mesh size sieve to obtain fine powder which was stored in an airtight container to prevent the active ingredients from evaporating and kept in a cupboard until the time for use. This procedure was adopted from Ojianwuna and Umoru 2010) and Ojianwuna *et al.*, (2011) with slight modifications.

Preparation of plant crude extract

The crude extracts were obtained by homogenizing 10g of dust (plant sample) with 100ml (80/20) of solvents (v/v) (hydro-ethanol) and left for 24hours thereafter, it was filtered with a filter paper and the residue (active compound) was evaporated over a water bath at 40°C . To obtain the final concentration, the extracts were dissolved in a known volume of distil water; to obtain 200, 400, 600, 800 and 1000ppm, 0.2, 0.4, 0.6, 0.8 and 1.0 mg of extract was dissolved in a ml of distil water. This method was adopted from Asid *et al.*, (2015).

Larvicidal Bioassay

Mortality and other effect on larvae were monitored every 12 hours of 72 hours and recorded. Two replicates of each concentration were made and the mean value was taken as the final result. Each group had 20 larvae and distill water was used as control experiment. The concentrations used for study are 200ppm, 400ppm, 600ppm, 800ppm, 1000ppm.

The toxicity of crude extracts was tested by these methods:

Dipping Technique

The larvae of each group were carefully dipped into various concentrations of crude extracts with a dip net and allowed to stand in it for 40seconds, and then introduced into petri dish with food.

Feeding technique

Various concentration of the crude extract was added to larvae food in petri dish and the larvae were introduced into the petri dish. This method used was adopted from Asid *et al* (2015) with slight modification.

Statistical analysis of Data

The results obtained were recorded and statistical analysis was performed using two-way ANOVA (Ms Excel), Descriptive statistics and Probit Analysis (SARS). Significance was determined at $P<0.05$.

RESULTS

Table 1 shows the mortality observed within a period of 72 hours, percent mortality and probit mortality of *Zingiber officinale* on *Musca domestica* larvae using dipping technique. The result shows that mortality was significant ($P < 0.05$) as concentrations increased from 200-1000ppm but mortality was not significant as duration of exposure increased from 12-72 hours. The LC_{50} and LT_{50} are 0.4898 mg/ml and 40 hours (Figures 1&2). No mortality was observed when the insects were fed with crude extracts but they (insect larvae) were weakened (their movement was sluggish and they didn't pupate).

Tables 2 and 3 shows the results obtained when *zingiber officinale* and *Dennettia tripetala* were mixed in equal proportions using dipping and feeding techniques. Mortality was generally significant as both concentrations and duration of exposure increased when the insects were dipped in crude extract and when they were fed with crude extract. When they were dipped in crude extract, a LC_{50} and LT_{50} of 0.2754 mg/ml and 30 hours was obtained (Figures 3 and 4) while LC_{50} and LT_{50} of 0.4786 mg/ml and 44 hours was obtained when they were fed with crude extract (Figures 5 and 6).

Table 1: percent mortality and probit mortality of *M. domestica* larvae exposed to *Z. officinale* for 72 hours by dipping method

Conc	Log	period of exposure(hours)							percent	probit	
Mg/ml of	conc								mortality	mean	mortality
H ₂ O		12	24	36	48	60	72				
0.2	-0.6690	0	1	2	2	2	2	45	1.5	4.87	
0.4	-0.3974	2	2	4	2	1	2	65	2.167	5.39	
0.6	-0.2218	2	3	2	3	2	3	75	2.5	5.67	
0.8	-0.0969	2	3	2	3	3	5	90	3	6.28	
1.0	0.0000	3	3	4	3	3	4	100	3.33	8.71	
Mean		1.8	2.4	2.8	2.6	2.2	3.2				




Figure 1: Relationship between probit mortality of housefly larvae and logarithm concentration to give ml concentration at (antilog) LC₅₀ when larvae were dipped in *Z. officinale*.




Figure 2: Relationship between probit mortality of housefly larvae and duration of exposure period to give time at LT₅₀ when larvae were dipped in *Z. officinale*.

Table 2: Percent mortality and probit mortality of *M. domestica* larvae exposed to *Z. officinale* and *D. tripetala* in equal proportions for 72 hours by dipping method

Conc	Log	period of exposure(hours)						percent	probit	
Mg/ml of	conc							mortality	mean	mortality
H ₂ O		12	24	36	48	60	72			
0.2	-0.6690	0	0	2	2	3	3	50	1.667	5.00
0.4	-0.3974	0	1	2	3	4	3	65	2.1667	5.39
0.6	-0.2218	1	2	3	4	4	1	75	2.5	5.67
0.8	-0.0969	2	3	4	4	3	3	95	3.1667	6.64
1.0	0.0000	2	4	3	5	3	3	100	3.333	8.71
Mean		1	2	2.8	3.6	3.4	2.6			




Figure 3: Relationship between probit mortality of housefly larvae and logarithm concentration to give ml concentration at (antilog) LC_{50} when larvae were dipped in crude extracts of *Z. officinale* and *D. tripetala*.




Figure 4: Relationship between probit mortality of housefly larvae and duration of exposure to give LT_{50} when larvae were dipped in crude extracts of *Z. officinale* and *D. tripetala*.

Table 3: Percent mortality and probit mortality of *M. domestica* larvae exposed to *Dennettia tripetala* and *Z. officinale* in a 50:50 proportion for 72 hours by feeding method

Conc	Log	period of exposure(hours)							percent	probit	
Mg/ml of	conc								mortality	mean	mortality
H ₂ O		12	24	36	48	60	72				
0.2	-0.6690	0	1	1	0	0	2	20	0.667	4.16	
0.4	-0.3974	0	3	0	2	0	0	25	0.833	4.33	
0.6	-0.2218	2	4	0	4	2	0	60	2	5.25	
0.8	-0.0969	4	5	5	2	0	0	80	2.667	5.84	
1.0	0.0000	5	5	4	3	0	0	85	2.833	6.04	
Mean		2.2	3.6	2	2.2	0.4	0.4				




Figure 5: Relationship between probit mortality of housefly larvae and logarithm concentration to give ml concentration at (antilog) LC_{50} when larvae were fed with *Z. officinale* and *D tripetala* in equal proportions.




Figure 6: Relationship between probit mortality of housefly larvae and duration of exposure to give LT_{50} when larvae were fed with *Z officinale* and *D tripetala* in equal proportions

187 DISCUSSION AND CONCLUSION

188 From the results obtained, crude extracts are toxic to freshly emerged larvae of *Musca domestica*.
 189 The high insecticidal activity and post application effect of these crude extracts is as a result of
 190 their constituent compounds; the volatile nature of the constituent compound could have also
 191 increased larvae mortality and insecticidal activities. It was also observed that test plant materials
 192 may have synergistic effects when they were mixed in equal proportions which led to its lower
 193 LC₅₀ and LT₅₀. Similar findings have shown that when extracts are mixed together, there may be
 194 synergistic effects. Akunne *et al* (2014) reported that *Piper guinense* and *Zingiber officinale* had
 195 additive effects against *Callosobruchus maculatus* when mixed in a 50:50 proportion.
 196 Ojianwuna and Umoru, (2010) also reported that *C. citratus* and *O. suave* had additive effect
 197 against *C. maculatus* in cowpea when they were mixed in a 60:40 proportion. Dawudo and
 198 Ofuya (2000) reported that the mixture of fruit powders of *Piper guineese* and *Dennettia*
 199 *tripetala* in equal proportion (50:50) had additive effects on *C. maculatus* and significantly
 200 caused their mortality, reduced their oviposition and adult emergence. Ojianwuna *et al* (2011)
 201 reported that *Ocimum suave* (wild basil) leaf oil had a LC₅₀ value of 0.09ml/50ml of water and
 202 LT₅₀ value of 4.40 hours when they studied toxicity of *Ocimum suave* leaf oil on adult housefly.
 203 They also reported that there was significant difference (P<0.05) as both concentrations
 204 increased from 0.05-0.20ml/50ml of water and duration of exposure increased from 1-6 hours.
 205 Soonwera (2015) also reported that the LT₅₀ and LC₅₀ values of *S. aromaticum* oil is 27.05 hours
 206 and 9.83% while *C. nardus* oil has LT₅₀ value of 38.99 hours and LC₅₀ value of 13.60% and *C.*
 207 *odorata* oil has LT₅₀ value of 52.08 hours and LC₅₀ value of 29.36% respectively when he
 208 evaluated larvicidal and oviposition deterrent activities of essential oils against housefly.

209 Dipping technique was observed to cause higher mortality than feeding technique as shown in
 210 the results obtained, this could be because larvae inhaled the extracts which may had acted as
 211 fumigants and contact poison to cause mortality. The potency of the extracts may have reduced
 212 when the food (mixed with plant extracts) came in contact with the digestive enzymes in the
 213 larvae. The larvae may have also been repelled and may not have fed on the food (mixed with
 214 plant extracts). This might be the reason why the dipping technique caused more mortality and
 215 had lower LC₅₀ when compared to the feeding method. This result is in accordance with the
 216 result obtained by Asid *et al* (2015) when he treated housefly larvae with *Citrullus colocynthis*
 217 (50% and 10%) using dipping and feeding method and observed that dipping method had more
 218 mortality and effects on larvicidal activity. Sripongun (2008) also observed that dipping method
 219 was very effective when he evaluated the contact toxicity of crude extract of Chinese star anise
 220 fruits to housefly larvae and their development. But it is contrary to the observation of Ghamdi *et*
 221 *al* (2014) when they carried out potential studies of non-conventional chemicals against housefly
 222 larva; their study revealed that feeding method had more effects on larvicidal activity than
 223 dipping method. The difference in results could be because of differential mode of action of plant
 224 extracts and their effective concentration.

Asid *et al.*, (2015), observed that *Citrullus colocynthis* with 50% concentration against *M. domestica* larvae caused larva mortality, pupa mortality and inhibit adult emergence. Bosly, (2013), observed that *Mentha piperita* and *lavandula angustifolia* essential oils have great insecticidal effect on *M. domestica* larvae; the extracts had mortality, prolonged larva and pupa duration and reduced adult emergence. Islam and Aktar, (2013) reported that *Calotropis procera*, *Piper longum* and *Polygonum hydropiper* had synergistic (additive) effects against *M. domestica* larvae and seven vital life-history traits of *M. domestica* such as fecundity, percent egg hatch, larva duration, numbers of pupae and adults, female ratio and adult longevity. Meenakshisundaram, *et al.*, (2014) evaluated the effect of Sweet Flag Rhizome (*Acorus calamus*) extract for Biological activity against housefly and observed that the extract had high mortality against housefly and offers an alternative source for the control of houseflies. From the results obtained, plant extracts were toxic especially when mixed in equal proportions (50:50). Thus they can be considered as bioinsecticides and could be useful in integrated pest management (IPM) as the results obtained are promising. Although, bioinsecticides are not as active as synthetic insecticides and do not act as fast as them (synthetic insecticides), they (bioinsecticides) are better alternatives because they are cost effective, environmental friendly and readily available. Bioinsecticides are prophylactic measure of protection against insects.

REFERENCES

- Ahmed, S., Malik, H., Riaz, M. A. and Akthar, M. (2013). Influence of plant extracts on the Life History and Population Development of Housefly, *Musca domestica* L. *J. Zoology* **45** (2), 345-349.
- Akunne, C. E., Afonta, C.N., Mogbo, T.C., Ononye, B.U. and Ngenegbo, U.C. (2014). Evaluation of the Efficacy of Mixed Powders of *Piper guineense* and *Zingiber officinale* against *Callosobrochus maculatus* (F.) (Coleoptera: Bruchidae). *American Journal of Biology and Life Sciences*, **2**(2), 63-67.
- Alan and HuiMeng, (2008). Life cycle of housefly. www.vtaide.com. Retrieved on 28th of June, 2015.
- Al-Ghamdi, Kh. M., Saleh, M. S., Mahyoub, J.A., Asid, A. N., Al-Najada, A. R., Mamdouh, I. N. and Alfarhan, B. Z. (2014). Potential Studies of Non-Conventional Chemicals against Housefly Larvae *Musca domestica* L. *Journal of Life Science* **11**(12), 1046-1048.
- Ande, A. T. (2001). Biological Activities of some Plant Materials against the Housefly – *Musca domestica*. *J. Biological Science* **1**(4), 293-296.
- Anyaele, O.O. and Amusan, A.A.S. (2003). Toxicity of Hexanolic Extract of *Dennettia tripetala* (G. Baxa) on Larvae of *Aedes aegypti* (L). *African Journal of Biomedical Research*, **6**, 49-53.

- 261 Asid, A. N., Al-Ghamdi, Kh. M., Mamdouh, I. N. and Mangoud, A. (2015). Assessment of Bitter
262 Apple, *Citrullus colocynthis* Extracts on the Housefly, *Musca domestica* L. *Journal of*
263 *Advances Environmental Biology*, **9**(8), 72-78.
- 264 Begum, N., Sharma, B. and Pandey, R. S. (2010). Evaluation of Insecticidal Efficacy of
265 *Calotropis procera* and *Annona squamosa* Ethanol Extracts against *Musca domestica*. *J.*
266 *Biofertilizers and Biopesticides* **1**(1), 1-6.
- 267 Bosly, A. H. (2013). Evaluation of Insecticidal Activities of *Mentha piperita* and *Lavandula*
268 *angustifolia* Essential Oils against Housely, *Musca domestica* L. *Journal of Entomology*
269 *and Nematology*, **5**(4), 50-54.
- 270 Buczacki, S. (2002). Housefly. www.arkive.org/house-fly/musca-domestica. Retrieved on 25ths
271 of April, 2015.
- 272 Dawudo, E. O. and Ofuya, T. I. (2000). Effects of Mixing Powders of *Piper guineense* Schum
273 and Thonn and *Dennettia tripetala* Baker on Oviposition and Adult Emergence of
274 *Callosobruchus maculatus* (F), Infesting Cowpea Seeds in Storage. *Appl. Trop. Agric.*, **5**
275 156-160.
- 276 El-Sherbini, T.G. and Hanykamel O.N. (2014). Insecticidal Effects of *Fortunella crassifolia*
277 Essential oil used against Housefly (*Musca domestica*). *Int. J. Current Microbiology and*
278 *Applied Sciences*, **3**(6), 1-9.
- 279 Gosh, A.K. (2011). *Zingiber officinale* a Natural Gold. *International Journal of Pharma and*
280 *Biosciences* **2**(1), 283-294.
- 281 Herve', B. D. B., Gbewonyo, S. W. K. and Obeng-Ofori, D. (2008). Toxicity, Growth
282 Regulatory and Repellent Activities of Medicinal Plant Extracts on *Musca domestica* L. *J.*
283 *Biotechnology* **7**(24), 4635-4642.
- 284 Idoko, J. E. and Adebayo, R. A. (2011). Effect of *Dennettia tripetala* Powder on Reduced Rates
285 of Pirimiphos – Methyl Singly and Combined in Management of *Callosobruchus*
286 *maculatus* (F). *International Journal of Biology* **3**(2), 174-178.
- 287 Ihemeje, A., Ojiannaka, M. C., Obi, K. C. and Ekwe, C. C. (2013). Biochemical Evaluation of
288 Pepper fruit (*Dennettia tripetala*) and its use as Substitute for Ginger in Zobo Drink
289 Production. *Int. J. Academic Research* **4**(6), 513-521.
- 290 Islam, M. S. and Aktar, M. J. (2013). Larvicidal Efficacies of some Plant Extracts and their
291 Synergistic Effects with Cypermethrin on the Life-History Traits of *Musca domestica* L.
292 *International Journal of Innovations in Bio-Sciences* **3**(3), 92-103.
- 293 Keiding, J. (1986). Division of Vector Biology and Control. *J. World Health Organisation*

- 294 Meenakshisundaram, T. J., Harikrishnan, N. R. and Anna, T. (2014). Evaluation of Extract from
295 Sweet Flag Rhizome for Biological Activity against Housefly. *Journal of Advanced*
296 *Veterinary Science and Technology* **3**(1), 140-144.
- 297 O'Toole, C. (2002). The New Encyclopedia of Insects and their Allies. [www.arkive.org/house-](http://www.arkive.org/housefly/musca-domestica)
298 [fly/musca-domestica](http://www.arkive.org/housefly/musca-domestica). Retrieved on 25th of April, 2015.
- 299 Ojjanwuna, C. C., Edafemakor, A. G. and Iloh, A. C. (2011). Toxicity of *Ocimum suave* (wild
300 basil) Leaf Oil on Adult Housefly (*Musca domestica*). *Journal of Agricultural Science and*
301 *Soil Science* **1**(10), 417-420.
- 302 Palacios, S. M., Berton, A., Rossi, Y., Santander, R. and Alejandro, U. (2009). Efficacy of
303 Essential Oils from Edible Plants as Insecticides against the House fly, *Musca domestica*
304 L. *Molecules* **14**, 1938-1947.
- 305 Pangnakom, U. and Kanlaya, S. (2014). Efficiency of Wood Vinegar Mixed with Some Plants
306 Extract against the Housefly (*Musca domestica* L.). *Journal of Biological, Food,*
307 *Veterinary and Agricultural Engineering*, **8**(9), 1-4.
- 308 Sinthusiri, J. and Soonwera, M. (2010). Effect of Essential Oils against Larvae, Pupae and Adults
309 of Housefly (*Musca domestica*). *Journal of Agricultural Technology*, 639-642.
- 310 Soonwera, M. (2015). Larvicidal and Oviposition Deterrent Activities of Essential Oils against
311 Housefly (*Musca domestica* L.). *J. Agricultural Technology* **11**(3), 657-667.
- 312 Sripongpun, G. (2008). Contact Toxicity of the Crude Extract of Chinese Star Anise Fruits to
313 Housefly Larvae and their Development. *J. Science and Technology* **30**(5), 667-672.
- 314 Stuart and Bennett, (2003). "Housefly" <http://www.thepiedpiper.co.uk/~th6a.htm>
- 315 Urzúa, A., Santander, R. I., Echeverriai, J., Cabezas, N., Palacios, S. M. and Rossi, Y. (2010).
316 Insecticide Properties of the Essential Oils from *Haplopappus foliosus* and *Bahia*
317 *ambrosoides* against Housefly, *Musca domestica* L. *Journal of Chilean Chemical Society*
318 **55**(3), 392-395.
- 319 www.britannica.com/plant/ginger. Retrieved 28th June, 2015.
- 320 www.healthtype.com/houseflydiseasestypes-spread-and-prevention.html. Retrieved 4th
321 February. 2016.
- 322 [www.qpm.ca/structure of housefly](http://www.qpm.ca/structure%20of%20housefly). Retrieved on 28th of June, 2015.
- 323 www.wikipedia.com/housefly. Retrieved on 28th of June, 2015.