

EGG QUALITY CHARACTERISTICS OF JAPANESE QUAILS (*Coturnix coturnix japonica*) FED VARYING LEVELS OF FERMENTED TARO COCOYAM (*Colocasia esculenta var esculenta*) MEAL.

Abstract.

A study was conducted to investigate the effect of FTCM on the egg quality parameters of Japanese quails. Two hundred and twenty five Japanese quails were randomly allotted to five dietary treatments (I-V) of 36 hens and 9 cockerels each. Each treatment was replicated thrice with 12 hens and three cockerels per replicate. In each of the five diets, FTCM was used to replace maize at 0%, 25%, 50%, 75% and 100% for treatments i, ii, iii, iv and v respectively. Quails were fed experimental diets over a period of 70 days. Feed intake was measured daily and quails were weighed weekly. Eggs were collected on weekly basis for egg quality analysis. Results showed that, mean weekly egg weight, shell weight, yolk weight, albumen weight, yolk index and egg breaking strength of quails fed treatments i and ii were significantly ($P < 0.05$) higher than those fed other treatments. Quails fed diets i, ii and iii recorded highest values of shell thickness. However, there were no significant ($P > 0.05$) differences in relative yolk and relative albumen weights across treatments. The results suggest that FTCM could replace maize up to 100% without adverse effect, but for premium egg quality, replacement should not exceed 25%.

KEY WORDS: Quails, FTCM, Maize and Egg quality

Introduction

Poultry production offers the greatest potential for bridging the gap of protein deficiency existing in Nigeria. The poor performance of non-ruminants (poultry, rabbits and pigs) is due to high cost of conventional ingredients such as maize, sorghum, millet, soybeans, groundnut etc. These conventional feedstuffs are expensive and are gravely faced by the problems of seasonality. This all year round unavailability has led to stiff competition between man and livestock industries as demand outstrips supplies. The need to search for alternative feed stuffs that are cheap and readily available cannot be over emphasized. A lot of Researchers have been able to use supplemental levels of non-conventional feed stuffs to reduce cost of production such as: Cocoyam [1 ; 2], Cassava [3], Rubber seed [4], Mango kernel [5; 6], Sweet orange peels [7] and *Prosopis* [8]. Taro cocoyam is one of the most widely spread root crop grown in humid tropics. About 60% of the world production is grown in Africa and 40% in Asia [9]. Fresh taro root is high in energy and contains about 25% starch and 1% protein. Taro has highest contents of P, Mg and Zn of all root/tuber crops and rich in all vitamins except for nicotinic acid [10]. Protein of taro cocoyam is well supplied with the essential amino acid but somehow low in histidine and lysine [11]. Cocoyam is cheaper and yields better than maize [12] and as such stands a better chance of replacing maize. However, the presence of anti-nutrients constitute a problem. Appropriate processing technique such as boiling, fermentation, toasting, sun drying

etc. could take care of its limitation [13]. This study was designed to investigate the effect of fermented taro cocoyam on egg quality parameters of Japanese quails.

Materials and Methods

The study was carried out in Cross River University of Technology Teaching and Research Farms Calabar, Calabar- Nigeria.

Experimental diets

Unpeeled taro cocoyam corms were bought from Bendeghe village, Etung, Cross River State, Nigeria. Corms were peeled and chopped into sizeable chips of about 14g each. These chips were submerged in a big plastic pot-like container with a tight lid and fermented in water at room temperature (28⁰c to 29⁰c) for 48hrs. Sun-drying lasted for a week (7days) and this reduced the moisture content to less than 10% for longer keeping. The 48hrs fermented taro cocoyam corms (FTCC) and other ingredients were milled separately and used to formulate experimental diets. The diets had varying protein (%) and energy (ME/Kcal/kg) levels of 21.20, 20.89, 20.78, 20.68, 20.59, and 2690.30, 2679.58, 2593.20, 2562.20 and 2552.90 for protein and energy respectively. These levels were for treatments 1, 2, 3, 4 and 5 respectively. The diets were analyzed according to the methods of [14]. Energy was calculated according to the methods of [15]. The percentage composition of the experimental diets is presented in Table one.

Table 1: Percentage composition of diet with fermented taro cocoyam (*Colocasia esculenta* var *esculenta*) meal

for laying Japanese quails (*Coturnix coturnix japonica*).

Levels of inclusion

Ingredient	0%	25%	50%	75%	100%
Maize	46.00	34.50	23.00	11.50	
Cocoyam	0.00				
Soybean	0.00	11.50	23.00	34.50	
Fish meal	46.00				
P.K.C	20.80	21.00	19.70	19.90	
Wheat offal	20.00				
Bone meal	4.20	4.60	5.80	6.00	
Salt	6.00				
Palm oil	10.90	10.00	10.00	10.00	
Vita/ min premix	10.00				
	10.00	10.00	9.75	9.10	
	8.30				
	7.00	7.00	7.00	7.00	
	7.00				

	0.50 0.50 0.50 0.50 0.10 1.20	0.50 0.50 0.50 0.40	0.50 0.50 0.50 7.50	0.50 0.50 0.50 1.00
Total	100.00 100.00	100.00	100.00	100.00
Analyzed nutrients:				
Crude protein (%)	21.20 20.59	20.89	20.78	20.63
ME/Kcals/Kg	2690.30 2552.90	2679.58	2593.20	2562.20

60

61

Experimental birds, design and management

62 The birds (quails) used for the experiment were a total of twelve (12) hens and three (3)
63 cockerels per replicate. In summary, a total of 225 quails (180 hens and 45 cockerels) were
64 studied over a period of ten weeks (70 days). Birds were randomly allotted to five dietary
65 treatments 1 , 2 ,3 , 4 and 5 formulated with 48hrs fermented taro cocoyam meal (FTCM) at 0% ,
66 25% , 50% ,75% and 100% replacement levels respectively (Table 1). Quails were housed in
67 three tier cages of dimension 0.75m² x0. 38m².The cages were made up of wood and wire mesh.
68 Same quantity of feed was weighed and served to the quails on weekly basis, the feed was
69 increased by 50g weekly. Fresh, clean water was supplied *ad libitum* daily. Adequate bio-
70 security and bio- sanitary measures were strictly adhered to. Records of feed intake were kept
71 daily, initial body weight was taken on commencement and subsequently on weekly basis. Total
72 number of eggs laid per replicate group were collected twice daily (8am and 4pm) and recorded.
73 At the end of every week fresh eggs were collected for egg quality analysis.

74

75

Determination of egg quality parameters

76 Nine fresh eggs (large, medium and small) were separated from each treatment (3 eggs from
77 each replicate) and used at weekly interval to determine egg quality parameters throughout the
78 experimental period. The eggs were weighed using a sensitive electronic weighing balance
79 (JJ560 USA) citizen scale to the nearest 0.01g. Each egg was weighed and broken around the
80 equator with care being taken to keep the yolk intact. Yolk was separated from the albumen
81 using an egg separator and weighed separately. Relative yolk and relative albumen weights were

82 calculated in percentage by relating the yolk or albumen weights measured to the nearest gram to
83 the whole weight of the particular egg and multiply by 100.

84 $\text{Relative yolk weight} = \text{Yolk/Egg wt.} \times 100$

85 $\text{Relative albumen weight} = \text{Albumen/Egg wt.} \times 100$

86 Egg shells were sun- dried for three days and their weights taken with the aid of a sensitive
87 electronic balance. Shell thickness was measured using micrometer screw gauge. The
88 measurement were taken from the pointed, middle and broad part of the egg and the mean thus
89 obtained and measured to the nearest 0.01mm. Yolk height was taken with the aid of a
90 Spherometer at the heighest point of its surface and yolk width was measured at long and short
91 axis using vernier caliper. The values obtained were used to calculate yolk index. Egg breaking
92 strength (EBS) was calculated using the formular suggested by [16] and represented as;

93 $\text{Yolk Index} = \text{mean height/mean width}$

94 $\text{EBS} = 50.86 \times (\text{EW})^{0.915}$, where EW =Egg weight.

95 Data obtained were subjected to analysis of variance using the completely randomized design
96 (CRD) as described by [17]. Least significant difference method was used to separate means that
97 differed significantly [17].

98 **Results and Discussion**

99 The egg weights of quails fed control diets and 25% inclusion levels of cocoyam had
100 significantly ($p < 0.05$) better egg weights compared to quails on other dietary treatments. [18]
101 indicated that egg mass or weight can be used as a criteria in assessment of nutritional status.
102 Similar results were reported in which egg weights increased significantly ($p < 0.05$) at higher
103 levels of dietary protein inclusion [19 and 20].The weight ranged between 9.40 to 10.80g. This
104 range was higher than the 8.92 to 10.13g reported by [20] and the 7.89 to 9.30g reported by [21].
105 [22] indicated that a good quail egg should weigh about 9.30g and above. Quails fed various
106 diets met this requirement. The egg shell weights of quails placed on treatments i and ii were
107 significantly ($p < 0.05$) heavier than those of other treatments. It was observed that beyond 25%
108 inclusion levels , the egg shell weights decreased with increased supplementation. This could be

109 attributed to decrease in dietary protein across the treatments. The egg shell weights ranged from
 110 1.12 to 1.30g. This range was higher than 0.90-0.93g reported by [19] and the 0.91 to 0.95
 111 reported by [23]. It could be as a result of the varying Ca:P ratio in the experimental diets of the
 112 various experiments. Result showed that shell thickness of quails fed control diets and 25%
 113 inclusion levels of cocoyam had significantly ($p<0.05$) thicker shells than quails fed other diets.
 114 Shell thickness ranged from 0.203 to 0.217mm. The result was lower than that reported by
 115 [19](0.24-0.26mm). The result contradicts with the report of [24] who asserted that, shell
 116 thickness declines with age due to increase in egg size which forces the constant amount of shell
 117 to spread thinner and thus reducing the thickness and quality. But was in agreement with the
 118 report of [25] who asserted that, the amount of shell materials deposited per egg does not change
 119 with age and size. Otherwise, treatments V and IV should have had significantly ($P<0.05$) thicker
 120 shells according to [24] . Egg breaking strength followed similar trend with quails placed on
 121 diets I and II having significantly ($P<0.05$) higher values compared with others. The result
 122 ranged from 442.1 to 502.77. This implies that eggs of quails placed on treatments I and II will
 123 not easily get cracked or broken when compared with eggs of other treatments. Generally, egg
 124 breaking strength (EBS) of quails is lower than that of chickens which could be as high as
 125 2322 ± 55.60 [26 ; 27]. By implication, the higher the EBS the better the egg quality. Yolk weight
 126 and albumen weights of quails fed control and 25% inclusion levels of cocoyam had significantly
 127 ($P<0.05$) higher weights than those of other treatments. Yolk and albumen weights ranged from
 128 3.28 to 3.60g and 5.00 to 5.91g respectively. The result of the yolk weight was in agreement with
 129 the report of [25] who noted that, albumen weight (AW) is negatively correlated with the yolk
 130 weight (YW).This infers that, eggs with higher values of AW will have lower yolk weights. The
 131 values recorded for AW in this finding were different from those recorded by [23] (4.42 to
 132 4.60g).The result was in agreement with the reports of [28] who indicated that AW was highly
 133 correlated with egg weight. It was observed that treatments I and II with highest values of egg
 134 weights (EW) had highest values of albumen weights (AW). The AW is sufficient to measure
 135 egg quality [29]. There were no significant ($P>0.05$) differences recorded in relative yolk
 136 weights (RYW) and relative albumen weights (RAW) across the treatments. RYW and RAW
 137 ranged from 33.33 to 34.89 and 52.75 to 54.72 respectively. The results of RYW and RAW are in
 138 agreement with the observation of [28] who observed that, as the egg size (ES) increases, the
 139 RYW decreases and the RAW increases. This result also agrees with the findings of [25] who

asserted that although the age of a layer has positive relationship with ES, it exerts a negative influence on the yolk properties. Result showed significant ($P<0.05$) difference in yolk index (YI) with treatment I and II having highest values. The YI ranged from 0.468 to 0.489. This range varied from the 0.35 reported by [19]. YI is the measure of the standing up quality of the yolk. Average values for fresh eggs usually fall between 0.40 to 0.42 [29]. The authors further stated that, the YI of 0.25 or lower indicates the weakness of the yolk and extreme difficulties in handling it for measurement without breaking. In this findings, one could deduce that the YI of all the treatments were within the range and could be handled for measurement without scattering easily.

Table 2: Summarized table of egg quality characteristics of laying quails fed vary levels of fermented taro cocoyam meal.

Parameter	0%	25%	50%	75%
100%				
<u>External characteristics</u>				
Egg weight (g)	8.636 ± 0.41 ^a	8.633±0.40 ^a	8.128±0.32 ^b	
7.757±0.41 ^c 7.516±0.52 ^c				
Egg shell weight (g)	1.29±0.08 ^a	1.30±0.08 ^a	1.19±0.09 ^b	1.15±0.08 ^c
1.12±0.10 ^d				
Egg breaking strength	502.36±19.11 ^a	502.77±18.58 ^a	472.68±14.87 ^b	451.23±19.00
^c 442.21±23.77 ^c				
Shell thickness(mm)	0.216±0.004 ^a	0.216±0.005 ^a	0.217±0.005 ^a	
0.203±0.006 ^b 0.205±0.006 ^b				
<u>Internal characteristics</u>				
Yolk weight (g)	3.60±0.10 ^a	3.45±0.07 ^b	3.34±0.07 ^c	3.28±0.10 ^c
3.28± 0.10 ^c				
Albumen weight (g)	5.90±0.25 ^a	5.91±0.24 ^a	5.56±0.16 ^b	5.21±0.62 ^c
5.00±0.35 ^c				
Relative yolk weight (%)	33.47±0.43	33.40±0.41	34.05±0.49	34.73±1.00
35.32±1.29				

Relative albumen weight (%)	54.61±0.39	54.65±0.35	54.73±0.34	53.52±0.84
	52.75±1.19			
Yolk index	0.488±0.06 ^a	0.489±0.06 ^a	0.475±0.005 ^b	0.472±0.002 ^b
	0.468±0.002 ^b			

Different superscripts (a, b, c and d) within the row indicates significant (p<0.05) differences.

Conclusion

The results of this study suggests that, fermented taro cocoyam meal could replace maize up to 100% without compromising egg quality but for premium egg quality, replacement should not exceed 25%.

References

- Agwunobi LN, Agwukan PO, Cora OO, Isika MA. Studies on the use of *Colocasia esculenta*(taro cocoyam) in the diets of weaned pigs. *Tropical Animal health and production*. 2002; 34(3):243-247.
- Okon BI, Obi MB, Ayuk AA. Performance of quails (*Coturnix coturnix japonica*) fed gradual levels of boiled sun-dried tarococoyam(*Colocasia esculenta var esculenta*) as a replacement of maize. *Agricultural Journal*. 2007; 2(6):654-657.
- Ogbonna JU. Nutritive value of cassava peels meal for poultry. *A Review of the proceeding of the 2nd annual conference of Animal Science Association of Nigeria*. 1997; Lagos, sept.

- 172 Agwunobi LN, Ina-ibar OB. Processed rubber seed meal in the diets of growing Japanese quail
173 (*Coturnix coturnix japonica*).*Proceedings of the 32nd Annual conference of the Nigerian society*
174 *for Animal production (NSAN). 2007;Calabar, march,Pp.386-388.*
- 175 Abang FB, Ejugwu G, Aker DT. Nutrient profile of Japanese quail(*Coturnix coturnix japonica*)
176 breast muscles and egg (whole) fed sun-dried mango(*Mangifera indica*) kernel meal. I.O.S.R
177 *Journal of Agriculture and Veterinary Science (IOSR-JAVS). 2015; 3(8):37-40.*
- 178 Diarra SS, Usman BA, Igwebuike JU. Replacement value of boiled mango kernel in broiler
179 finisher diets. *ARPRO Journal of Agriculture and Biological Science. 2008; 5:47-52.*
- 180 Oluremi OIA, Ojighen VO, Ejembi EH. The nutritive potentials of sweet orange (*citrus*
181 *sinensis*) rind in broiler production. *International Journal of Poultry Science. 2006;5(7):613-617.*
- 182 Yusuf ND, Ogali DM,Hassan DI, Musa MM, Doma UD. Effect of decorticated fermented
183 *Prosopis* seed meal(*Prosopis africana*) on growth performance of broiler chickens. *International*
184 *Journal of Poultry Science. 2008; 7(11):105-1057.ISSN.*
- 185 Viotti V. Taro production in hawaiian. Island. *Honolulu advertiser. 2004; March.*
- 186 Purseglove JW. *Tropical Crops Monocotyledons*.John willey and sons. 1992; New York.
- 187 Ohtsuka R, Kawabe T, Inaoka T, Suzuki T, Hong F, Akinchi OT, Sugaharu T. Composition of
188 local and purchased food consumed by the Gidra in lowland papua. *Ecology of Feed and*
189 *Nutrition. 1984; 15:159-169.*
- 190 Food and Agricultural Organization(FAO). Food and Agricultural Organization production year
191 book. Food and Agricultural Organization(FAO) of the United Nations. 2004; Rome, Italy.
- 192 Obi MB. Improving the nutritional value of taro cocoyam(*Colocasia esculenta var.esculenta*)
193 through processing for the feeding of growing and laying quails. Ph.D. thesis submitted to
194 Department of Animal Science University of Calabar. 2010; Pp210.
- 195 Association of Official Analytical Chemist . *Official methods of analysis* (17th edition). 2000;
196 Washington D.C.
- 197 Carpenter KJ, Clegg KM. The metabolisable energy of poultry feeding stuffs in relation to their
198 chemical composition. *Journal of science and Agriculture. 1956; 7:45-51.*
- 199 Arad z, Marder J. Difference in egg shell quality among the Sinai bedouing fowls, the
200 commercial white leghorn and their cross breeds. *British Poultry Science. 1982; 23:107-112.*
- 201 Steel RGD, Torrie JH. *Principles and proceedings of statistics,Mc Graw-hill book.1980;*
202 *Co.Inc.Newyork.*

- 203 Lin, H., Buyse, J., Sheng,Q.K., Xie,Y.M. and Song,J.L.(2004). Effects of ascorbic acid
204 supplementation on the immune function and laying performance of heat-stressed laying hens.
205 *Journal of Agriculture and Environmental*. 1:103-107.
- 206 Bawa,G.S. Lombin,H., Karson,P.,Musa, U., Pay,E. and Shamak,D. (2011). Response of Japanese
207 Breeder quails to varying Dietary Protein Levels in the Tropics. *Nigerian Journal of Animal*
208 *Production*. 38:43-54.
- 209 Babangida, S.and Ubosi,C.O.(2006). Effects of varying dietary protein levels on the performance
210 of laying Japanese quails (*Coturnix coturnix japonica*) in a semi-arid environment. *Nigeria*
211 *Journal of Animal Production*, 33(1):45-52.
- 212 Jika, C.P.(2015). Egg quality parameters of quails fed graded levels of *Prosopsis africana* seed
213 coat meal as a replacement for maize. B. Agric. Project submitted to the Department of Animal
214 Production, University of Agriculture, Makurdi.Pp5.
- 215 Yamane etal
- 216 Sachdev, A.K., Ahya, S.D and Cropals,R. (1975). The ability of hens to maintain calcium
217 deposition in the egg shell and egg yolk as the hen ages. *Poultry Science*.54:1729-1723-1729.
- 218 Roland (Sr), D.A., Putnem, C.E. and Hilburn,R.L.(1978). The influence of age on ability to hen
219 to maintain egg shell calcification when stressed with inadequate dietary calcium. *Poultry*
220 *Science*.57:1616-1621.
- 221 Essien, A I, Ekanem EO. Effect of the age of layers, rate of egg produce and time of oviposition
222 on some initial egg shell quality indices in a humid tropical environment. *Beitrage zur*
223 *Tropischen land wirl. Shaft veterinary medicine*.1990; 28:125-131.
- 224 Oluwofeso, O and Omisami,OO. (2008). Utilization of cassava peels treated with yeast culture
225 by egg laying chickens. *Proceedings of the 33rd annual conference of the Nigeria society for*
226 *Animal production*. Ogun,2008. Pp492-495.
- 227 Woolford, R. (1985). Nutrition and egg quality. *Poultry International* (May edition). 46 – 48.
- 228 Fletcher OI, Brittan WM, Rahn AO, Savage SL. The influence of layer flock age on egg
229 component yield and solid content. *Poultry Science*. 1980; 60(80):983-987.
- 230 Card, I E and Neshiem, MC(1972). *Poultry Production* 11th edition. Philadelphialea and
231 Febiger.
- 232 Abang,F.B, Ejugwu, G. and Aker, D.T.(2015). Nutrient profile of Japanese quail(*Coturnix*
233 *coturnix japonica*) breast muscles and egg (whole) fed sun-dried mango(*Mangifera indica*)
234 kernel meal. *I.O.S.R Journal of Agriculture and Veterinary Science(IOSR-JAVS)*. 3(8):37-40

- 235 Agwunobi, L.N., Agwukan, P.O.,Cora,O.O. and Isika M.A.(2002). Studies on the use of
236 *Colocasia esculenta*(taro cocoyam) in the diets of weaned pigs. *Tropical Animal health and*
237 *production*. 34(3):243-247.
- 238 Agwunobi,L.N. and Ina-ibar, O.B.(2007). Processed rubber seed meal in the diets of growing
239 Japanese quail (*Coturnix coturnix japonica*).*Proceedings of the 32nd Annual conference of the*
240 *Nigerian society for Animal production (NSAN).Calabar, march,2007.386-388.*
- 241 Arad,z. and marder,J. (1982). Difference in egg shell quality among the Sinai bedouing fowls,
242 the commercial white leghorn and their cross breeds. *British Poultry Science*.23:107-112.
- 243 Association of Official Analytical Chemist (2000). *Official methods of analysis* (17th edition).
244 Washington D.C.
- 245 Babangida, S.and Ubosi,C.O.(2006). Effects of varying dietary protein levels on the performance
246 of laying Japanese quails (*Cortunix coturnix japonica*) in a semi-arid environment.*Nigeria*
247 *Journal of Animal Production*, 33(1):45-52.
- 248 Bawa,G.S. Lombin,H., Karson,P.,Musa, U., Pay,E. and Shamak,D. Response of Japanese
249 Breeder quails to varying Dietary Protein Levels in the Tropics. *Nigerian Journal of Animal*
250 *Production*. 2011. 38:43-54.
- 251 Card, I.E and Neshiem,M.C.(1972). *Poultry Production* 11th edition.Philadelphia lea and
252 Febiger.
- 253 Carpenter, K.J and Clegg, K.M.(1956). The metabolisable energy of poultry feeding stuffs in
254 relation to their chemical composition. *Journal of science and Agriculture*.7:45-51.
- 255 Diarra,S.S, Usman, B.A. and Igwebuikie,J.U. (2008). Replacement value of boiled mango kernel
256 in broiler finisher diets. *ARPRO Journal of Agriculture and Biological Science*. 5:47-52.
- 257 Essien AI, Ekanem EO. Effect of the age of layers, rate of egg produce and time of oviposition
258 on some initial egg shell quality indices in a humid tropical environment. *Beitrage zur*
259 *Tropischen land wirl. Shaft veterinary medicine*.1990; 28:125-131.
- 260 Fletcher OI, Brittan WM, Rahn AO, Savage SL. The influence of layer flock age on egg
261 component yield and solid content. *Poultry Science*. 1980; 60(80):983-987.
- 262 Food and Agricultural Organization(FAO),(2004). Food and Agricultural Organization
263 production year book. Food and Agricultural Organization(FAO) of the United Nations. Rome,
264 Italy.
- 265 ***Jika, C.P.(2015). Egg quality parameters of quails fed graded levels of *Prosopsis africana*
266 seed coat meal as a replacement for maize. B. agric, project submitted to the Department of
267 Animal Production, University of Agriculture, Makurdi.Pp53.

- 268 Lin, H., Buyse, J., Sheng,Q.K., Xie,Y.M. and Song,J.L.(2004). Effects of ascorbic acid
269 supplementation on the immune function and laying performance of heat-stressed laying hens.
270 *Journal of Agriculture and Environmental*. 1:103-107.
- 271 Obi,M.B.(2010). Improving the nutritional value of tarococoyam(*Colocasia esculenta*
272 *var.esculenta*) through processing for the feeding of growing and laying quails. P.hD. thesis
273 submitted to Department of Animal Science University of Calabar, Calabar.Pp210
- 274 Ogbonna,J.U. (1997). Nutritive value of cassava peels meal for poultry. *A Review of the*
275 *proceeding of the 2nd annual conference of Animal Science Association of Nigeria. Lagos, sept*
276 *1999*.
- 277 Ohtsuka,R., Kawabe,T., Inaoka,T., Suzuki, F., Hong.O.T., Akinchi and Sugaharu,T.(1984).
278 Composition of local and purchased food consumed by the Gidra in lowland papua. *Ecology of*
279 *Feed and Nutrition*. 15:159-169.
- 280 Okon,B.I, Obi,M.B. and Ayuk,A.A.(2007). Performance of quails(*Coturnix coturnix japonica*)
281 fed gradual levels of boiled sun-dried tarococoyam(*Colocasia esculenta var esculenta*) as a
282 replacement of maize. *Agricultural Journal*.2(6):654-657.
- 283 Oluremi,O.I.A., Ojighen,V.O. and Ejembi, E.H. (2006). The nutritive potentials of sweet orange
284 (*citrus sinensis*) rind in broiler production. *International Journal of Poultry Science* 5(7):613-
285 617.
- 286 Oluwofeso, O. and Omisami,O.O. (2008). Utilization of cassava peels treated with yeast culture
287 by egg laying chickens. *Proceedings of the 33rd annual conference of the Nigeria society for*
288 *Animal production. Ogun,2008,Pp492-495*.
- 289 Oluyemi J.A. and Robert, F.A. (2000) Poultry production in the warm climate. Macmillian
290 publisher. L.t d.London.
- 291 Purseglove, J.W.(1992). *Tropical Crops Monocotyledons*.John willey and sons. New York.
- 292 Roland (Sr), D.A., Putnem, C.E. and Hilburn,R.L.(1978). The influence of age on ability to hen
293 to maintain egg shell calcification when stressed with inadequate dietary calcium. *Poultry*
294 *Science*.57:1616-1621.
- 295 Sachdev, A.K., Ahya, S.D and Cropals,R. (1975). The ability of hens to maintain calcium
296 deposition in the egg shell and egg yolk as the hen ages. *Poultry Science*.54:1729-1723-1729.
- 297 Steel,R.G.D. and Torrie,J.H.(1980). *Principles and proceedings of statistics,Mc Graw-hill book*
298 *Co.Inc.Newyork*.
- 299 Viotti,V.(2004). Taro production in hawaiian. Island. *Honolulu advertiser*,march,6,2004.

- 300 Woolford, R.(1985). Nutrition and egg quality. *Poultry International* (May edition). 46-48.
- 301 Yusuf, N.D., Ogali, D.M.,Hassan,D.I., Musa,M.M., and Doma, U.D.(2008). Effect of
 302 decorticated fermented *Prosopis* seed meal(*Prosopis africana*) on growth performance of broiler
 303 chickens. *International Journal of Poultry Science*.7(11):105-1057.ISSN.