

KUNYIMA PROCEDURE TO STUDY ENZYMATIC MECHANISM IN CASSAVA CYANIDE REMOVAL. CASE OF SASSOU VARIETY (DEMOCRATIC REPUBLIC OF CONGO).


Abstract

Despite many publications on cyanide elimination in cassava by soaking in water, it still remains our preoccupation because the techniques elaborations of cyanide removal depend on a good uptake of phenomena. The traditional methods of cassava processing (steeping, peeling, drying) permit to eliminate a large proportion of cyanhydric acid, but the technology of the cassava treatment is still nowadays to be improved. That suggests the importance to better its processing and its production on the technological level by a suitable and reliable fast method (a few minutes) in view to eliminate the cyanide completely. So enzymatic kinetic study of a cassava variety, Sassou, in Democratic Republic of Congo, has been undertaken in this paper at temperature of 25°C by means of only pH-meter. The procedure needs to be called **KUNYIMA Procedure** for the rising scientific generations in order to remember the cute idea framed by Dr. Anaclet KUNYIMA B., Ordinary Professor, University of Kinshasa Democratic Republic of Congo. It is needful to size a cassava reactor which will be a fast procedure of elimination of cyanide, economically payable and efficient with the production of cassava without cyanide according to the order of magnitude of the kinetic constants values found. In this paper cassava hydroxynitrilase has been successfully investigated and its K_M has been found $1,5 \cdot 10^{-2}$ M; $V_{max} = 0,0072 \text{ Mh}^{-1}$, $R^2 = 0,972$.

Keywords: KUNYIMA Procedure, Michaelis Constant, Cassava, Hydroxynitrilase, pH-meter.

1. Introduction

The cassava which constitutes the basic food for numerous population of DRC is toxic because it contains the cyanide whose toxicity lead to the diseases such as legs paralysis (KONZO), goitre, mental cretinism, dwarfishness [1,2,3]. This is due to the formation hindering of $T_4(T_3)$, essential thyroidean hormones for the differentiation and the maturation of foetal tissues needful for the growth of the skeleton and of almost all the organs and also for the development of the central nervous system. A thyroid insufficiency started during the foetal life or at the birth entails a regression of the cortical neurons [2,4,5,6]. That leads to the insufficiency of the brain development and to its definitive lesions. Theses hormones monitor metabolism of glucids, lipids, protids, and regulate the speed of the enzymatic reactions and the temperature of the human body [5,6,7]. The essential toxic principle which exists in variable quantities in all the parts of the plant of cassava is a chemical compound called linamarin (93%-97%). It often coexists with its methylic homologous called methyl linamarin or lotaustralin (3%-7%) [8,9,10,11,12]. The linamarin is a cyanogénétique glucoside which is transformed into hydrocyanic acid (toxic) as it is shown below.



The elimination of the cyanide from cassava has been realized many times and continues to be realized in Democratic Republic of Congo (DRC). In certain villages the cassava is soaked in flowing river water and a great quantity of cyanide is eliminated but the processing is not economically payable because not rapid (time is money). In the others villages the cassava is soaked in the butts water or exposed to the sun, the result is never satisfactory. Neither simple cooking nor addition of hot water to cassava produces totally satisfaction.

2. Materials and Methods

a) Materials

- Sassou with the following characteristics has been used [13]:

Family: *Euphorbiaceae*

Order: *Euphorbiales*

Class: Dicotyledones

Branching: Angiospermes

Gender: *Manihot esculenta crantz*

Variety: Sassou



Figure 1. Sassou Variety, DRC (Picture taken on the sample of experimentation after one week)

- pH-meter (Hanna, HI98127), a balance KERN (EMR 500-1), distillate water and cryothermostat Julobo have been used.

The figures have been plotted by means of origin 8.1, Matlab 7.12.0 (R 2011a) and IBM SPSS statistic viewer 19 programs.

b) Methods

66 **1°) Calculations**

67 Cassava roots have been peeled and sliced. The slices of 10g of fresh Sassou have been
68 soaked into 70 ml of distillate water; the acidity of solution has been followed as function of
69 time at 25°C. The concentrations of HCN and CN^- have been calculated as follow [14].
70

71
$$pH = \frac{1}{2} pKa - \frac{1}{2} \log Ca$$

72

73

$$Ca = [HCN] = 10^{(pKa-2pH)}$$



75
$$[HCN] = Ca$$

76
$$Ka = \frac{[H^+][CN^-]}{[HCN]} = \frac{[H^+]^2}{Ca}$$

77
$$[CN^-] = [H^+] = \sqrt{Ca \cdot Ka}$$

78
$$[CN^-] = [H^+] = \sqrt{10^{(pKa-2pH)} \cdot 10^{-pKa}}$$

79
$$[CN^-] = \sqrt{10^{-2pH}}$$




80
$$[CN^-] = 10^{-pH}$$

82 $pKa (HCN) = 9,31$ at 25°C [14,15].

83 - The velocity of HCN formation (v_a) and of CN^- formation (v_x) have been calculated
84 ($v = \frac{\Delta C}{\Delta t}$)

85 - Enzymatic kinetic of Michaelis-Menten and Lineweaver-Burk method have been used
86 [16,17].

87 - The two following reactions



88


89

90 being consecutives, the last one is the limiting reaction ($k_1 > k_2$).

91 It has been observed that the errors on the calculated parameters are of the order of magnitude
92 of 10^{-7} , so decision has been taken to neglect them in the tables.
93

2°) Approximation


The contribution of Iotaustralin (3%-7%) has been neglected for two reasons: the first is its presence in very small quantity and the second one is that the ethylic group on it makes it less competitive in the production of HCN. The reaction of production of HCN and acetone from only linamarin is the uppermost and promoted. Hence the large quantity of HCN, more than 90%, comes from linamarin [18].



As it can be seen stoichiometrically the $[HCN] = [cyanohydrin] = [S] = [Substratum] = Ca$. Globally it can be said that the quantity of HCN produced is proportional to the quantity of extractive substratum (cyanohydrin).

3°) Michaelis-Menten kinetic for a reaction with one substratum has been reported.

The kinetic model is based on the complex formation ES according to the following steps [16,17] :



A relation between the reaction velocity, both the substratum and the enzyme concentrations and individual step velocities is established. At the onset the reaction velocity is equal to $[ES]$ and the kinetic constant k_3 :

$$v = \frac{dP}{dt} = k_3[ES] \quad \dots\dots (2)$$

The complex formation velocity and its dissociation velocity are given by:

$$\text{Formation velocity of } ES = k_1[E][S] \quad \dots\dots\dots (3)$$

$$\text{Dissociation velocity of } ES = (k_2 + k_3)[ES] \quad \dots\dots\dots (4)$$

$$\text{Under stable state conditions: } k_1[E][S] = (k_2 + k_3)[ES] \quad \dots\dots\dots (5)$$

$$\text{By rearrangement of equation 5, one obtains: } [ES] = \frac{[E][S]}{(k_2 + k_3)/k_1} \quad \dots\dots\dots$$

(6)

Michaelis constant has been defined (K_M)

$$K_M = \frac{k_2 + k_3}{k_1} \quad \dots\dots\dots (7)$$

$$(7) \text{ and } (6) \text{ give rise to } [ES] = \frac{[E][S]}{K_M} \quad \dots\dots\dots (8)$$

Moreover $[S] = [S]_{total}$ if $[E] \ll [S]$

$$[E] = [E]_{total} - [ES] \quad (9)$$

$$[ES] = \frac{([E]_{total} - [ES])[S]}{K_M}$$

(10)

Rearrangement of equation 10 gives

$$[ES] = [E]_{total} \frac{[S]}{[S] + K_M}$$

(11)

$$(11) \text{ and } (2) \text{ give rise to } V = k_3[E]_{total} \frac{[S]}{[S] + K_M} \quad (12)$$

The maximum velocity, V_{max} , is reached when all the enzymatic sites are saturated of substratum, that means when $[S] \gg K_M$ and so

$$\frac{[S]}{[S] + K_M} \text{ tends towards } 1.$$

$$\text{Hence } V_{max} = k_3[E]_{total} \quad (13)$$


$$(13) \text{ and } (12) \text{ give } V = V_{max} \frac{[S]}{[S] + K_M} \quad (14)$$

Michaelis-Menten equation describes the kinetic curve of $V_a - [S]$ [4,16,17].

- For the weak concentrations of substratum, when $[S] \ll K_M$, $V = [S] \frac{V_{max}}{K_M}$ and the velocity is directly proportion to substratum concentration
- When $[S] \gg K_M$, $V = V_{max}$ and the velocity is independent on the substratum concentration
- The meaning of K_M is evident. When $[S] = K_M$, $V = \frac{V_{max}}{2}$. K_M is the needful substratum concentration to allow enzyme to reach $\frac{V_{max}}{2}$.

4°) Solid-liquid phase extraction thermodynamics

It can be imagined the extraction of cyanide in solid phase by means of water obeys instantaneously a distribution of this cyanide between solid phase and water phase, distribution characterized by an equilibrium constant [19,20,21]



In this case of extraction the equilibrium is however displaced towards the liquid phase (water). For each phase the balance sheet can be written:

a. Liquid phase

$$C_{HCN_{liq.\ phase}} = X_{HCN_{free\ liq.\ phase}}$$

Where C_{HCN} is the formality of HCN and X_{HCN} is the current concentration of HCN. Given the HCN acidity constant ($5 \cdot 10^{-10}$) is weak its dissociated fraction can be neglected.

b. Solid phase

When the extractant solvent (water) is put in contact with the solid phase two phenomena are bred:

1. Solubilisation

During this step one part of HCN remains unsolubilized that means under the bound form and one part of HCN is solubilized that means under free form near to migrate in water phase because we think that the solubilisation is time depend phenomenon. The solubilized form has been called free species, that means detached from its stub.

2. Transfer (diffusion)

The solubilized form (free species) is transferred to water. Thus the extraction implies the solubilisation and the transfer. The balance sheet in this solid phase can be written:

$$C_{HCN_{solid\ phase}} = X_{HCN_{bound\ solid\ phase}} + X_{HCN_{free\ solid\ phase}}$$

There is difference between apparent equilibrium constant (K_{app}) and the true equilibrium constant (K_{true}). It should be noted that in solid phase there is unsolubilized HCN (bound) and solubilized HCN not yet transferred. So it can be imagined a relationship between K_{app} which is the ratio of the no identical species and K_{true} which is the ratio of the free species. Note that we are working in dilute medium.

Indeed

$$K_{true} = \frac{X_{HCN_{free\ liq.\ phase}}}{X_{HCN_{free\ solid\ phase}}}$$

$$K_{app} = \frac{C_{HCN_{liq.\ phase}}}{C_{HCN_{solid\ phase}}} = \frac{Ca}{C_o - Ca}$$

Where $Ca = C_{HCN_{liq.\ phase}}$

C_o is the initial concentration obtained from the tray of the curve giving $C_{HCN_{liq.\ phase}}$ versus time.

$$K_{app} = \frac{X_{HCN_{free\ liq.\ phase}}}{X_{HCN_{bound\ solid\ phase}} + X_{HCN_{free\ solid\ phase}}}$$

$$\frac{1}{K_{app}} = \frac{X_{HCN_{bound\ solid\ phase}}}{X_{HCN_{free\ liq.\ phase}}} + \frac{X_{HCN_{free\ solid\ phase}}}{X_{HCN_{free\ liq.\ phase}}}$$

$$\frac{1}{K_{app}} = \frac{1}{K_{true}} + X_{HCN_{bound\ solid\ phase}} \frac{1}{X_{HCN_{free\ liq.\ phase}}}$$

$$\frac{1}{K_{app}} = \frac{1}{K_{true}} + X_{HCN \text{ bound solid phase}} \frac{1}{X_{HCN \text{ free liq. phase}}}$$

$$\frac{C_o - Ca}{Ca} = \frac{1}{K_{true}} + X_{HCN \text{ bound solid phase}} \frac{1}{Ca}$$

$$\frac{C_o}{Ca} = 1 + \frac{1}{K_{true}} + X_{HCN \text{ bound solid phase}} \frac{1}{Ca}$$

$$\text{The intercept of } \frac{C_o}{Ca} \text{ versus } \frac{1}{Ca} \text{ gives } \frac{1}{K_{true}} + 1.$$

$$K_{true} = \frac{1}{b-1} \text{ Where } b \text{ is the intercept}$$

$$K_{true} = e^{-\Delta G^\circ / RT}$$

$$\ln K_{true} = -\frac{\Delta G^\circ}{RT}$$

$$\Delta G^\circ = -RT \ln K_{true}$$

3. Results and Discussion

The cassava roots generally contain dried materials (35%), glucides (89%), lipides (1%), protides (2,5%), fibres (4,5%), ash (3%), calcium (0,1%), phosphorus (0,1%), iron (0,003%), natrium (0,003%), potassium (1%), starch and linamarin [22]. The percentage of each compound varies with the nature of the soil.

Neither the proteins hydrolysis such as albumines, globulines, glutelines, prolamines and protamines which are all of them the globular proteins solubles in water nor starch hydrolysis giving rise to amylose (soluble) and amylopectine (unsoluble) can not be foreseen, because the above mentioned proteins are absents in cassava roots and the starch hydrolysis happens only when the hot water is used [23]. Let us recall that in our case however the distillate water has been used at 25°C on fresh cassava and during experimentation all the solutions have been covered to avoid the influence of the air CO₂.

The phosphoric acid formation under our experimental conditions is improbable. The minerals elements present in cassava roots do not give the contribution to the formation of the observed acidity, otherwise the increase of pH would have been observed. Hence the evolution of the acidity in water with time is exclusively due to the specific enzymatic hydrolysis of the linamarin, compound giving the specific reaction with linamarase and hydroxynitrilase [22,23,24].

So in the following table it can be seen the decrease of pH with time.


Table 1: Measured and calculated parameters as a function of time at 25 °C

Time	p^H	$C_{HCN} = S = Ca$	$C_{CN^-} = X$	1/S	v_a	v_x	1/ v_a for Ca
h	-	M	M	M ⁻¹	Mh ⁻¹	Mh ⁻¹	M ⁻¹ h
0	6,89 ± 0,001	-	-	-	-	-	-
2	6,63 ± 0,001	0,00011	2,34E-07	9090,90909	5,5E-05	1,17E-07	0,182 E+05
4	6,35 ± 0,002	0,00041	4,46E-07	2439,02439	15,0E-05	1,06E-07	0,067 E+05
6	6,19 ± 0,001	0,00085	6,45 E-07	1176,47059	22,0E-05	0,99E-07	0,045E+05
8	5,89 ± 0,001	0,00339	12,88E-07	294,98525	127,0E-05	3,22E-07	0,008E+05
10	5,71 ± 0,001	0,00776	19,49E-07	128,86598	218,5E-05	3,31E-07	0,005E+05
12	5,56 ± 0,001	0,01549	27,54E-07	64,55779	386,5E-05	4,03E-07	0,003E+05
14	5,32 ± 0,002	0,04677	47,86E-07	21,38123	1564,0E-05	10,16E-07	0,001E+05
16	5,09 ± 0,001	0,13489	81,28E-07	7,41345	4406E-05	16,71E-07	0,00023E+05
18	4,83 ± 0,001	0,44668	148,00E-07	2,23874	15589,5E-05	33,36E-07	0,000064E+05
20	4,58 ± 0,002	1,41254	263,02E-07	0,70794	48293,0E-05	57,51 E-07	0,00002E+05
22	4,35 ± 0,001	4,07380	446,68E-07	0,24547	1,33063	91,83E-07	0,752
24	4,23 ± 0,002	7,07945	588,84E-07	0,14125	1,50283	71,08E-07	0,665
26	4,13 ± 0,001	11,22018	741,31E-07	0,08913	2,07037	76,24E-07	0,483
28	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
30	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
32	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
34	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
36	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
38	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
40	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
42	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
44	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
46	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305
48	4,03 ± 0,001	17,78279	933,25E-07	0,05623	3,28131	95,97E-07	0,305

226 In this table 1 the time is expressed in hours (h), the mean values of pH are used, M is the
 227 molarity of HCN and CN^- , v_a is the velocity of HCN formation ($\frac{\Delta C_{\text{HCN}}}{\Delta t}$) and v_x is the
 228 velocity of HCN dissociation ($\frac{\Delta C_{\text{CN}^-}}{\Delta t}$) or the velocity of CN^- formation

229


230 The table 1 gives the values of pH of solution as a function of time and the calculated values
 231 of $[\text{HCN}]$, $[\text{CN}^-]$, $\frac{1}{Ca}$, the velocity of HCN formation (v_a), velocity of CN^- formation (v_x) are
 232 indicated. When velocity of HCN formation (v_a) is compared to velocity of CN^- formation
 233 which is also the velocity of HCN dissociation (v_x) it can be seen that v_a is higher than v_x as
 234 it is plotted in figure 2.



235

236

Figure 2a. v_x versus v_a




237

238

Figure 2b. v_a versus v_x

239 This shows that HCN is effectively a weak acid, it is weakly dissociated. This behavior can
 240 also be observed in figure 3 where C_{CN^-} is plotted versus C_{HCN} .

241



242

243

Figure 3. C_{CN^-} versus C_{HCN}

244


245

246

Also both the velocity of HCN formation (v_a) and the velocity of its dissociation (v_x) increase with time and afterwards (28h) remain constant showing the roll of hydroxynitrilase as it can be seen in figure 4.

248

249



250

251

Figure 4a. v_a versus time

252





Figure 4b. v_x versus time

Note that the change of pH is not observed after 28hours for this variety of cassava (Sassou) where the extraction of HCN at 25°C is fast. The enzymes have selectivity and proficiency to accelerate the velocity of the formation reaction of a complex enzyme-substratum [4,17].



where E is enzyme and S is substratum, ES is a complex enzyme-substratum and P is the product of reaction. The formation of the complex ES gives rise to a transition state for the reaction having a low activation energetic barrier. The reaction needs less external energy contribution to happen. The critical first stage in a reaction with enzymatic catalysis is the complex formation ES. The complex ES is the reactant in the conversion step of the substratum in product, its concentration determines the reaction velocity which is maximum when the all enzyme molecules are complexed by substratum.

This situation corresponds to the substratum high concentrations where the enzyme is a so called saturated with the substratum. For an enzyme given concentration, the figure of initial velocity of reaction (V_a) versus substratum initial concentration $[S]$ shows a typical curve of saturation kinetic where a tray is observed at high $[S]$ when velocity $V = V_{max}$. The approximation has been applied as it can be seen in figure 5; the curve shape predicted by the theory has been observed. The hydroxynitrilase obeys Michaelis relation.

Hans Lineweaver-Dean Burk method has been used $\frac{1}{V_a} = \frac{1}{V_{max}} + \frac{K_M}{V_{max}} \frac{1}{[S]}$ as it is indicated in figure 5 where it can be observed that

$$K_M = 0,015 = 1,5 \cdot 10^{-2}; V_{max} = 0,0072; -\frac{1}{K_M} = -67,457; R^2 = 0,972.$$




Figure 5. $\frac{1}{V_a}$ versus $\frac{1}{[S]}$

The ulterior experiments will show us the choice between the two-two sequential reaction mechanism, two-two ping-pong reaction mechanism or a reaction of enzyme with one substratum. The values of K_M is very variable, it depends on enzyme nature, substratum nature, temperature and pH.

K_M is not only a measure of the catalytic efficiency but also the enzyme affinity for its substratum. It has been reported [16] that the enzymes have properties sensitive to pH. Many proteins are active in narrow zone of pH generally comprised between 5 and 9 because pH is acting on different factors [15] :

- 1) Substratum-enzyme bond
- 2) Catalytic activity of enzyme
- 3) Ionisation of substratum
- 4) The variations of proteins structure (generally important only at utmost pH).




Figure 6. V_a versus pH

It has been reported that the initial velocities of many enzymatic reactions as a function of pH give rise to bell shaped curves. These curves are the reflex of the ionization state certain residues of amino acid which are in specific ionization state for the activity of the enzyme [15]. In the above mentioned case however (figure 6) this phenomenon is not entire because only the half bell shaped curve is observed in solid-liquid phase extraction in the interval of the studied pH. It should be noted that the determination of enzymatic reactions mechanisms from only kinetic data are uncertain. It is necessary to resort to other experimental approaches should confirmation is needed. Experiments are continuing in laboratory and we are hopeful that in nearest future the real mechanism will be elucidated. When the HCN concentration is followed as a function of time a characteristic curve is obtained showing a tray corresponding to its extractive initial concentration (figure 7).





Figure 7. C_{HCN} versus time

317 It is observed the same phenomenon for C_{CN^-} concentration as it is shown in figure 8.
318



319

320 **Figure 8. C_{CN^-} versus time**

321

322 Concerning the thermodynamics study, table 2 gives the parameters allowing to calculated
323 K_{true} and ΔG° .

324

325 **Table 2. Values of C_0 (HCN), C_a , $\frac{C_0}{C_a}$, $\frac{1}{C_a}$**

326

C_a	$\frac{1}{C_a}$	C_0	$\frac{C_0}{C_a}$
0	0	0	0
0,00011	9090,90909	17,78279	161661,72730
0,00041	2439,02439	17,78279	43372,65854
0,00085	1176,47059	17,78279	20920,92941
0,00339	294,98525	17,78279	5245,66077
0,00776	128,86598	17,78279	2291,59665
0,01549	64,55778	17,78279	1148,01743
0,04677	21,38123	17,78279	380,21788
0,13489	7,41343	17,78279	131,83179
0,44668	2,23874	17,78279	39,81103
1,41254	0,70794	17,78279	12,58923
4,07380	0,24547	17,78279	4,36516
7,07945	0,14125	17,78279	2,51189
11,22018	0,08913	17,78279	1,58489

327

328 The figure 9 corresponding to the table 2 gives the value of the intercept, **b**.

329




Figure 9. C_0/Ca versus $1/Ca$

According to the values of the intercept K_{true} and ΔG° are calculated.

Indeed:

$$K_{true} = \frac{1}{b-1} = \frac{1}{3,16242 \cdot 10^{-6} - 1} = -146\,250,574 \, i^2$$

with i = imaginary number.

$$\Delta G^\circ = -RT \ln K_{true} = (-29,43 - 4,95 \ln i) \, \text{kJ mol}^{-1}.$$

It can be observed the existence of the negative equilibrium constant which is of course the imaginary constant meaning that the exchange of cyanide between the solid phase and water phase is so slow that no sharing equilibrium can be set up permanently between the content of those two phases.

The extraction of cassava cyanide by means of water is an irreversible phenomenon towards the water, it takes a few hours.

4. CONCLUSION

KUNYIMA procedure is a simple method proposed to deal with the enzymatic kinetic of the cyanide elimination in cassava using solely pH-meter in order to size a cassava processing reactor. It is aimed to build a fast and efficient method, economically payable (time is money) which will be a fast procedure of elimination of cyanide with the production of cassava without cyanide according to the order of magnitude of the kinetic constants values found.

The conception of cassava cyanide removal reactor depends on kinetic parameters [25,26]. It is in this logic KUNYIMA procedure should be uptaken.

KUNYIMA procedure will allow to distinguish the two-two sequential reaction mechanism from the two-two ping-pong reaction mechanism or the reaction of enzyme with one substratum [15].

KUNYIMA procedure needs to take into account thermodynamics information according which the HCN extraction from cassava roots to water at 25°C is an irreversible phenomenon towards the water. It will not be surprising to find the same observations on the other species at the other temperatures as it will be revealed in the futures papers.

Ultimately it can be said that linamarin and hydroxynitrilase are undoubtedly Michaelis enzymes [15].

REFERENCES

[1] Ernesto, M; Cardoso, AP; Nisala, D; Mirione, E. Massaza, F; Cliff, J; Haque, MR; Bradbury, Jtt. Persistent konzo and cyanogens toxicity from cassava in northern Mozambique, ACTA TROPICAL, 82(3): 357-362 Juin 2002.

[2] Agbor Egbe, T., I. Lape Mbome., The effects of processing techniques in reducing cyanogen levels during the production of some Cameroonian cassava foods, Journal Elsevier of Food Composition and Analysis 19, pp. 354–363, 2006.

[3] E.Atibu Kazinguvu, NM Mbuyamba, B.Kunyima, A.B.Kunyima, Kinetic study on elimination of cyanide from cassava; Annales Africaines de Médecine, Vol.1, N°1, Septembre 2007.

[4]Borel., Maquatre., Le Peuch., Randorx., Gillery., Bellon., Monboisse., Biochimie Dynamique, De Boeck et Larcier s.a. Université, Bruxelles, 1997.

[5] Felix Zelder., Lucas Tivana., Corrin-based chemosensors for the assured detection of endogenous cyanide, Journal Royal Society of Chemistry, Vol 13, N°1, pp 1–316, 2015.

[6] A.J. André Criqui., Influence de la liaison des hormones thyroïdiennes aux protéines plasmatiques sur l'homéostasie de la fonction thyroïdienne, Université Paul-Sabatier de Toulouse, Thèse de doctorat, 2006.

[7] FSANZ., Cyanogenic glycosides in cassava and bamboo shoots. A Human Health Risk Assessment, Technical report Series N°. 28, 2004.

[8] O. R. Ezeigbo, M. U. Ekaiko and Z. O. Ibegbulem, Effect of Cooking Time on Starch and Cyanide Contents of Freshly Harvested Cassava Tubers Used for Tapioca Production, Sciencedomain international., British Biotechnology Journal 8(4): 1-6, n°. BBJ.16944, 2015.

[9]FSANZ., Final assessment report proposal P1002. Hydrocyanic acid in ready-to-eat cassava chips, First review report. Report Number 2-09, 2009.

- 394 [10] Younoussa Diallo, Momar Talla Gueye, Cheikh Ndiaye, Mama Sakho, Amadou Kane,
395 Jean Paul Barthelemy, Georges Lognay., A new Method for the Determination of cyanide
396 Ions and theis quantification in some Cassava Varieties, American Journal of Analytical
397 Chemistry, 5, 181-187, Senegal, 2014.
- 398 [11] S. C. Kobawila, D. Louembe, S. Keleke, J. Hounhouigan and C. Gamba., Reduction of
399 the Cyanide Content during Fermentation of Cassava Roots and Leaves to Produce Biked
400 and Ntobambodi, Two Food Products from Congo, African Journal of Biotechnology, Vol. 4,
401 No. 7, pp. 689-696, 2005.
- 402 [12] Kaneko, J.J. ,Thyroid function In : Clinical biochemistry of domestic animals, 5th edition
403 San Diego,Academic Press, 571-588, 1997.
- 404 [13] L.Pauwels, Nzayilu N'ti : Guide des arbres et arbustes de la Région de Kinshasa –
405 Brazzaville, 1993.
- 406 [14] Skoog , West, Holler , Croush, Fundamentals of Analytical Chemistry, 9^e edition, De
407 Boeck, 2014 .
- 408 [15] Petre Atkins et Loretta Jones., Chimie : molécules, matières, métamorphoses, 3^{ème} éd.
409 Américaine, 1998.
- 410 [16] Voet D. et Voet J. G., Biochimie, 2^{ème} édition. De Boeck Université, pp. 472-491, 1998.
- 411 [17] Soustelle M., Mécanisme Réactionnel et cinétique chimique, Hermès Science
412 Publications, Lavoisier, 2011.
- 413 [18] Falade K.O. and Akingbala J.O., Utilization of cassava for food, Food Rev. Int., 27(1),
414 51-83, 2010.
- 415 [19] J.C. Jungers, L.Sajus, I.Aguirre, D. Decroocq, L'analyse Cinétique de la transformation
416 chimique, edition TECHNIP, France, 1967.
- 417 [20] Pierre Infelta et Michael Graetzel., Thermodynamique : Principes et Applications ;
418 Edition Brown Walker Press, Florida, 2006.
- 419 [21] A.B.Kunyima, LJ Muswema, BM Kunyima, Elimination of cyanide from cassava roots
420 grown in Congo, Case of bitter variety NGAMANZA: kinetic and thermodynamic studies;
421 Annales Africaines de Médecine, Vol.1, N°2, Mars 2008.
- 422 [22] Silvestre P., Arraudeau, M., Le manioc. Collection : Techniques Agricoles et
423 Productions Tropicales, XXXII. ACCT. Maisonneuve et Larose (Eds) 1983.
- 424 [23]Jonathan Clayden, Nick Greeres, Stuart Warren, Chimie Organique, 2^e édition, De
425 Boeck, 2013.
- 426 [24] Eric Giraud, Contribution à l'étude physiologique et enzymologique d'une nouvelle
427 souche de *lactobacillus plantarum* amylytique isolée du manioc fermenté, Thèse,
428 Université de Provence Aix-Marseille 1, Mars 1993.

- 429 [25] M.Bahroun Sami, Modélisation et approche thermodynamique pour la commande des
430 réacteurs chimiques catalytiques triphaniques continus et discontinus, Thèse, Université
431 Claude Bernard, Lyon 1, 2010.
- 432 [26] P.Trambouze, Jean Paul Euzen , Les réacteurs chimiques : De la conception à la mise en
433 œuvre, édition TECHNIP, France, 2002.