

# **Original Research Article**

## **TAXONOMIC COMPENDIUM OF PLANT COMMUNITY ON SHANKARACHARYA RESERVE FOREST**

### **Abstract**

The present investigation was carried out to study the taxonomic compendium of plant community on Shankaracharya reserve forest. Locally also known as Sulaiman Teng, the study site is situated between an altitude of 1575 to 1967 m amsl and a latitude of 34°04'35.56" N and 74°51'08.63" E longitude. The hillock over looks famous Dal lake and presents a beautiful look of Srinagar city on the eastern side of Jammu and Kashmir. The survey was conducted and plant samples were collected for identification. The taxonomic compendium of the plant community identified in Shankaracharya Reserve forest includes 72 Genera (69 Angiosperms and 3 Gymnosperms) and 84 species (78 Angiosperms and 6 Gymnosperms) out of which 54 were herbs and grasses, 12 were shrubs and 18 were trees. Monocots and dicots contributed nearly 91.2% of the total angiosperm population indicating that the forest grove is rich in pioneer and mid successional species composition. The study area hosts a remarkable floristic richness with majority of taxa belonging to family Fabaceae followed by Rosaceae, Poaceae and Pinaceae.

**Key words:** taxonomic, compendium community, Shankaracharya

### **Introduction**

The knowledge of patterns of variation in forest structure over time and space have always served as the basis of formulating forest management strategies that seek to sustain a broad array of forest goods and services [1]. The concept of sacred groves is prevalent in most parts of the world and Hughes and [2] have presented a detailed overview of their distribution around the globe mainly in Asia, Africa, Australia, Europe and America. A report of [3] has described sacred groves in Ghana, Senegal and Sumatra. In India, they occur in Western Ghats, Madhya Pradesh, Maharashtra, Meghalaya, Karnataka, etc., and found in variety of habitats ranging from scrub forests of Thar Desert, to rain forests in Western Ghats and temperate forests of Himachal Pradesh and Jammu and Kashmir in the North to deciduous forests of Kerala in the South. Estimates suggest that there might be between 1,00,000 to 1,50,000 sacred groves around the country [4]. Further, all forms of vegetation in the sacred groves are supposed to be under the protection of the reigning deity of that grove, and the removal of even a small twig is taboo [5]. SHANKARACHARYA RESERVE FOREST hillock also known as SULIMAN TENG or TAKHT-I- SULIMAN in Kashmir is one of such groves famous for religious tourism all over the country. This sacred groves is one of the first instances of traditional conservation in Kashmir; however, due to increasing loss in earlier times and biodiversity loss these habitats demand new conservation approaches

39 which will enable fair share of the wider values of conservation to the local communities and  
40 positive inherited attitudes towards conservation goals. Nature worship has been a key force  
41 of shaping the human attitudes towards conservation and sustainable utilization of natural  
42 resources. Such traditional practices have been invariably operating in different parts of India  
43 and Shankaracharya is one of the living examples of least studied plantation forest which can  
44 now be regarded as the remnant primary forest left untouched by the local inhabitants for  
45 conservation of biodiversity.

46 The vegetation of Kashmir forests has been occasionally explored in the past by few  
47 botanists. The prominent among them are [6 and 7]. The early twentieth century saw the  
48 work of [8 and 9] and who published their work in the form of manuals and floras. Their  
49 work was followed by that of [10 and 11]. [12] from Botanical Survey of India, surveyed  
50 many places in the valley with a view to document a comprehensive list of flora of this  
51 region. In all these studies, the vegetation characteristic of Shankaracharya hillock; which is  
52 largely characterized by a typical topographic physiognomy, is poorly cited. This hillock has  
53 a long history of degradation due to exploitative deforestation for timber and other NTFP's.  
54 However, following its massive re-vegetation programme in 1940 and 1970 with many  
55 species including exotic tropical pines viz., *Pinus helpensis*, *Pinus roxburghii* and *Pinus*  
56 *canarensis*, the hillock has been designated as a reserve forest. Despite a recorded history of  
57 more than seventy years since reforestation of this hillock was taken up, only one or two  
58 comprehensive attempts have been made to conduct a systematic study of the vegetation  
59 composition and its structure. The present study was; therefore, undertaken to study the  
60 floristic and successional status of this unique man made reserved forest grove's which now  
61 comprises many plant species of both tropical and temperate origin. The information  
62 generated from this study helped to classify this reserved forest hillock on the basis of  
63 floristic composition.

## 64 **Materials and methods**

### 65 **Study site**

66 The present investigations were carried out on Shankaracharya hillock which stands  
67 declared as a reserve forest under Section 11 of the Forest Act of 1987 and a Game Sanctuary  
68 under Section 5 of the Game Prevention Act of 1998 [13]. The hillock is maintained by the  
69 Forest Department, Government of Jammu and Kashmir for religious, aesthetic and

70 recreational purposes. The study site is located between 34°04' 35.56" to 34°05' 25.08" N  
 71 latitude and 74°50' 03.16" to 74°51' 08.63" E longitude (Table 1). Spread over an area of  
 72 141.93 ha, this forest grove lies in South-Eastern side of Srinagar and presents a marvelous  
 73 view of world famous Dal Lake and other adjoining environs. The altitude of the study site  
 74 varies from 1575 to 1967 m asl (Table 1) with entire hillock serving as a catchment area to  
 75 Dal Lake.

76 **Table 1: Geographical attributes and Lithology of Shankaracharya Reserve Forest**

| Altitude        | Aspect     | Longitude        | Latitude       | Lithology  | Slope      |
|-----------------|------------|------------------|----------------|------------|------------|
| 1575-1705 m asl | North West | 34° 04' 35.36" N | 74°50'03.132"E | Deep loam  | Steep      |
|                 | North East | 34°05'25.38" N   | 74°50'03.145"E | Fine loam  | Steep      |
|                 | South East | 34°05' 25.396" N | 74°50'03.148"E | Silty loam | Very steep |
| 1705-1835 m asl | North West | 34 °04' 35.44" N | 74°50'03.160"E | Deep loam  | Steep      |
|                 | North East | 34°05'25.46" N   | 74°50'03.155"E | Fine loam  | Very steep |
|                 | South East | 34° 05' 25.490"N | 74°50'03.158"E | Silty loam | Steep      |
| 1835-1967 m asl | North West | 34°04'35.508" N  | 74°50'03.16"E  | Deep loam  | Very steep |
|                 | North East | 34°04'35.563"N   | 74°50'03.16"   | Fine loam  | Very steep |
|                 | South East | 34°05'25.083"    | 74°51' 08.63"E | Silty loam | Very steep |

## 77 **Plant sample collection and Identification**

78 Significant taxonomic reservoirs [14] were used for plant identification in this study.  
79 Specimens of some of the species particularly difficult to identify at the species level were  
80 got identified by expert taxonomists in Division of Environmental Science, SKUAST of  
81 Kashmir, Shalimar, Srinagar and Department of Botany, University of Kashmir. For this  
82 purpose herbarium specimens with some distinguishing characteristics of general appearance  
83 of vegetative or reproductive material were prepared and these served as a crude basis for  
84 their classification.

## 85 **Results and discussion**

86 The vegetation survey conducted indicated that the selected reserve forest on  
87 Shankaracharya hill is rich in plant composition, supporting many plant species, of diverse  
88 taxonomic and ecological significance. Floristic composition and successional status of  
89 Shankaracharya groove sacred groves are small patches of man-modified conserved  
90 landscapes or lingering samples of natural vegetation with the highest levels of biological  
91 diversity. Besides centre of high species richness [15] these areas act as a gene pool and  
92 provide refuge to a large number of endemic, endangered and threatened species [16] and  
93 render many ecological services such as regulation of perennial water, maintaining local  
94 micro-environment and sustenance of biogeochemical cycles [17]. Among the 13,720 sacred  
95 groves in India, Shankaracharya is a reserve forest being maintained for aesthetic and  
96 recreational purposes [17]. The grove is very rich in species richness as depicted by plant  
97 community organizational analysis; of the study area, which revealed presence of 84 plant  
98 species belonging to 72 genera and 40 families. The data presented in Table 2 reveals that  
99 this reserve forest hosts 40 families (38 Angiosperms and 2 Gymnosperms), 72 Genera (69  
100 Angiosperms and 3 Gymnosperms) and 84 species (78 Angiosperms and 6 Gymnosperms)  
101 out of which 56 were herbs, 12 were shrubs, and 18 were trees (Fig. 1). Dicotyledons  
102 contributed nearly 91.2% of the total angiosperms recovered in the area. Fabaceae was the  
103 dominant family with (ten) 10 species followed by Rosaceae (8) Poaceae (7), Pinaceae (5),  
104 Asteraceae (5) Malvaceae (4) Lamiaceae (3), Convolvulaceae (3), Salicaceae (3),  
105 Cannabaceae (2), Caryophyllaceae (2), Polygonaceae (2), Apiaceae (2) and Plantaginaceae  
106 (2). Families with representation of only one species included Moraceae, Juglandaceae,  
107 Sapindaceae, Chenopodiaceae, Cupressaceae, Rhamnaceae, Berberidaceae, Oleaceae,  
108 Thymelaeaceae, Scrophulariaceae, Zygophyllaceae, Iridaceae, Caprifoliaceae,  
109 Ophioglossaceae, Violaceae, Urticeae, Labiatae, Boraginaceae, Campanulaceae,

Apocynaceae, Scrophulariaceae, Cupuliferae, Amaranthaceae and Hypericaceae. The survey of Shankaracharya forest hillock conducted by [18] reveals the presence of 49 families with dominance of dicotyledenous (85%). Out of the 86 plant species recorded in the study area, 18 were trees, 12 were shrubs and 56 were herbaceous (Table 2). This species representation pattern of herbs > shrubs > trees has also been reported by [19 and 20] in undisturbed patches of Branwar temperate coniferous forests of Kashmir and temperate broadleaved Rhododendron forest of Western Arunachal Pradesh respectively. The tree plantations thus established on Shankaracharya hillock (long devoid of native plant cover) have acted as successional catalysts, facilitating re-colonisation of many woody and non woody native flora. [21 and 22] have reported that planting of trees encouraged the invasion of local species and thereby resulting in enhanced richness of shrubs and herbaceous species. [23] reported that reforestation by diverse type of plant communities along the environmental gradient facilitated improvement in local habitat. The re-vegetation effort thus profoundly determine the patterns of community structure and its distribution.

The classification of the flora; recorded on Shankaracharya hillock, in [24] biological spectrum reveals that this forest groove is mainly shaped as annual seed bearing therophytic vegetation (65%) which complete their life cycle in one year and pass on unfavourable season by overwintering. The second dominant category of flora was that of phanerophytes (34.5%) which includes shrub and tree species whose perennating buds are borne on aerial shoot reaching a height of at least 25 cm or more above the ground surface. These results show that the flora at Shankaracharya Forest Reserve is predominantly therophytic to climate. [25] have reported almost similar pattern of biological spectrum for Kandi forest range in Kashmir respectively. [26] stated that while higher number of therophytes indicates the shortage of moisture, the less number of phanerophytes specifies that the selected forest is the best representative site for open physiognomy where the latter lifeform spectrum decreases with increasing altitude. The phanerophytes viz., *Cupressus torulosa*, *Cedrus deodara* and *Pinus wallichiana* form a dominant community with prominent association of *Tribulus terretris*, *Cannabis sativus*, *Linaria dalmatica*, *Taraxacum officinale*, *Medicago sativa*, *Cuscuta europaea*, *Artemisia obsinthum*, *Iris nepalensis*, *Lonicera quinquelocularis*, *Rumex acetos*, *Rumex orientalis*, *Lespedeza cuneata*, *Ophioglossum vulgatum*, *Cichorium intybus*, *Viola odorata*, *Panicum crusgalli*, *Foeniculum vulgare*, *Cynodon dactylon*, *Pegasus hermala*, *Chenopodium album*, *Cumimum cyminum*, *Utricadioca*, *Salvia moorcraftian*, *Trifolium repens*, *Nepeta cataria*, *Organum vulgare* and

143 *Rosa webbiana*, *Zizyphus vulgaris*, *Rubus fruticosus*, *Indigofera gerardiana*, *Rosa*  
144 *moschata*, *Berberis lycium*, *Jasminum humile* and *Daphne oleoides* through out the  
145 altitudinal range.

## 146 **Conclusion**

147 Measuring the success of ecological restoration is not as straightforward and simple as  
148 it appears. The overall results of this study can be concluded as the vegetation analysis  
149 conducted indicates presence of 40 families, 38 angiosperms and 2 gymnosperms represented  
150 by 54 grasses and herbs, 12 shrubs and 18 tree species. The extent of shrubs and grasses and  
151 herbs has greatly expanded due to improvement in site conditions over a period of induced  
152 secondary succession. Thus the reforestation of Shankaracharya forest has influenced the  
153 microclimate of the area as depicted by invasion of new species of shrubs and herbs on the  
154 hillock from the adjoining areas.

## 155 **References**

- 156 1. Spies TA. Forest Structure: A key to the ecosystem. Northwestscience.72. Special Issue  
157 No.2, USA. Structure and biomass in the Yucatan Peninsula, Mexico: effects of forest  
158 disturbance. *Forest Ecol. Manag.* 1998; **247** : 80-90.
- 159 2. Hughes JD, Chandran MD. Sacred groves around the earth: An overview, **In** : *Conserving*  
160 *the Sacred for Biodiversity Management* (eds. P.S., Ramakrishnan, KG., Saxena and  
161 Chandrashekara, U.M.) Oxford and IBH Publishing Co., New Delhi. 1997; 69-46.
- 162 3. MAB. Sacred places and vernacular conservation. M.A.B. 23. UNESCO, Man and  
163 Biosphere Programme, Paris. 1995; 20-21.
- 164 4. Malhotra KC, Gokhle Y, Chatterjee S, Srivastava S. *Cultural and Ecological Dimensions of*  
165 *Sacred Groves in India*, INSA, New Delhi. 2000.
- 166 5. Khan MS, Aamil M, Zaidi A. Associative effect of *Brady rhizobium* sp (vigna) and  
167 phosphate solubilizing bacteria on moongbean (*Vigna radiata* L.) wilczek, *Bio.*  
168 *Journal.* 1997; **10** : 101-106.
- 169 6. Jacquemont V. Letters from India, describing a journey in the British Dominions of India,  
170 Tibet, Lahor and Kashmir during the years 1828-1831, undertaken by order of the  
171 Freanch Government. London. 1835; 2.

- 172 7. Thomson T. Western Himalaya and Tibet: A narrative of journey through the mountains of  
173 Northern India. 1848.
- 174 8. Conventry BO. *Wild Flowers of Kashmir*, Series I-III. Raithby, Lawrence and Co., Ltd,  
175 London. concentration for 32 neotropical tree species. *Canadian Journal of Forest*  
176 *Research*. 1925;**33**: 1039-1045.
- 177 9. Mukerjee SK. A revision of Labiatae of Indian Empire. *Rec. Bot. Sum. India*. 1940; **14**: 1.
- 178 10. Pennell FW. . The scrophulariaceae of the Western Himalayas. Monog. *Acad. nat. Sci.*  
179 *Phinland*. 1943; 5.
- 180 11. Ludlow F. The Primulas of Kashmir. *J.R. Hort.* 1951; **76** : 191-206.
- 181 12. Rao APVP, Naidu MVS, Ramavatharam N, Rao GR. Characterization, classification and  
182 evaluation of soils of different land forms in Ramachandrapuram Mandal of Chitpor  
183 district in Andhra Pradesh for sustainable land use planning. *Journal of the India*  
184 *Society of Soil Science*. 2008; **50**(1) : 23-33.
- 185 13. Sidiq R. *Forest Records*, of J and K; 1987.
- 186 14. Willis JH. A Handbook to Plants in Victoria. Vol. I. 2<sup>nd</sup> Edn. (Melbourne Univ. Press:  
187 Melbourne.) Specimens examined: Jammu & Kashmir: Ladakh, 11000ft., July, 1905,  
188 A. Meebold 3315 (CAL); Suru Valley, 14500 ft., 27.6.1928. B.B.Osmaston 1970; 204  
189 (K).
- 190 15. Upadhaya K, Pandey HN, Law PS, Tripathi RS. *Biodiversity Conserv.* 2003; **12**: 583-597.
- 191 16. Jamir SA, Pandey HN. *Biodiversity Conservation*. 2003; **12**: 1497-1510.
- 192 17. Dar GH, Bhagat RC, Khan MA. *Biodiversity of the Kashmir Himalaya*, Valley Book  
193 House, Srinagar. 2002.
- 194 18. Paul TM. Studies on woody ornamental plants for land scape use. Ph.D thesis submitted to  
195 Sher-e-Kashmir University of Agricultural Science and Technology, Srinagar.  
196 2001;20-31.
- 197 19. Dar IY. Edaphic factors and plant community organization in Branwar forest of Kashmir  
198 Himalaya. M.Sc Thesis Submitted to University of Kashmir Srinagar. 2011. 30-36.

- 199 20. Paul A. Studies on diversity and regeneration ecology of Rhododendrons in Arunachal  
200 Pradesh. Ph.D. thesis, Assam University, Silchar, Assam, India. 2008.
- 201 21. Verma RK, Kapoor KS, Rawat RS, Subramani SP, Kumar S. Analysis of plant diversity  
202 in degraded and plantation forests in Kunihar Forest Division of Himachal Pradesh.  
203 *Indian Journal of Forestry* 2004; **27**(2) : 223-227.
- 204 22. Verma RK, Kapoor KS, Rawat RS, Subramani SP, Kumar S. Analysis of plant diversity  
205 in degraded and plantation forests in Kunihar Forest Division of Himachal Pradesh.  
206 *Indian Journal of Forestry* 2005; **28**(1) : 11-16.
- 207 23. Dar GH, Christensen KI.. Habitat diversity and zonality of vegetation in Sind valley,  
208 Kashmir Himalaya. *Nature and Biosphere*. 1999; **4**(1&2) : 49-71.
- 209 24. Raunkier C. The Life Forms of Plants and Statistical Plant Geography, being the collected  
210 papers of C. Raunkier (Ed. Frank N. Egerton), Oxford University Press, Oxford.  
211 Reprinted 1978. Ayer Co Pub., in the "History of Ecology Series". 1943.
- 212 25. Hillal AR. Plant diversity and regeneration status in Kandi range of Kamraj Forrest  
213 Division of Kashmir. Ph.D thesis submitted to Sher-e-Kashmir University of  
214 Agricultural Sciences & Technology of Kashmir, Shalimar, Srinagar; 2014
- 215 26. Asri Y. Plant diversity in Touran Biosphere Rservoir, Publishing Research Institute of  
216 Forests and Rangelands, Tehran, Iran. 2003; **305** : 306.

217

218

219 **Table 2: Floristic list and plant life form spectra along an altitudinal gradient in Shankaracharya Reserve Forest**

| S. No. | Family       | Species                    | Common name/Local name        | Life form | Altitude (m) |           |           |
|--------|--------------|----------------------------|-------------------------------|-----------|--------------|-----------|-----------|
|        |              |                            |                               |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 1.     | Rosaceae     | <i>Pyrus cumminis</i>      | Pear/Tang                     | T         | +            | +         | -         |
|        |              | <i>Prunus cerasifera</i>   | Plum/Gurdhoal                 | T         | +            | -         | -         |
|        |              | <i>Prunus armenica</i>     | Apricort/Cheer                | T         | +            | -         | -         |
|        |              | <i>Crataegus oxycantha</i> | Hawthorn/Ring                 | S         | +            | -         | -         |
|        |              | <i>Rosa webbiana</i>       | Wild rose/Arwal               | S         | +            | -         | +         |
|        |              | <i>Rubus fruticosus</i>    | Black berry/Daen Chanch       | S         | +            | +         | -         |
|        |              | <i>Rubus pungens</i>       | Rubus oldhamii/Rang ratch     | S         | -            | +         | -         |
|        |              | <i>Rosa moschata</i>       | Rose hip                      | S         | +            | -         | +         |
| 2.     | Salicaceae   | <i>Populus alba</i>        | Silver poplar (Dodhi fras)    | T         | +            | -         | -         |
|        |              | <i>Populus nigra</i>       | Black poplar (Bati fras)      | T         | +            | -         | -         |
|        |              | <i>Salix fragilis</i>      | Brittle willow                | T         | +            | -         | -         |
| 3.     | Moraceae     | <i>Morus alba</i>          | White mulberry/Toot           | T         | +            | +         | -         |
| 4.     | Juglandaceae | <i>Juglans regia</i>       | Walnut/Doon                   | T         | +            | -         | -         |
| 5.     | Cannabaceae  | <i>Celtis australis</i>    | Nettle tree/Brimji            | T         | +            | -         | +         |
|        |              | <i>Cannabis sativa</i>     | Hemp/Bhang                    | H         | +            | +         | +         |
| 6.     | Sapindaceae  | <i>Aesculus indica</i>     | Indian horse chestnut/Handoon | T         | +            | +         | -         |

Contd...

220

221

222

| S. No. | Family       | Species                     | Common name/Local name  | Life form | Altitude (m) |           |           |
|--------|--------------|-----------------------------|-------------------------|-----------|--------------|-----------|-----------|
|        |              |                             |                         |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 7.     | Fabaceae     | <i>Robinia pseudoacacia</i> | Black locust/Kikar      | T         | +            | +         | -         |
|        |              | <i>Quercus ilex</i>         | Holm oak                | T         | -            | +         | +         |
|        |              | <i>Indigo gerardiana</i>    | Himalayan indigo/Neel   | S         | +            | +         | -         |
|        |              | <i>Lespedeza cuneata</i>    | Chinese bush clover     | H         | +            | -         | +         |
|        |              | <i>Medicago sativa</i>      | Lucerne/Poshi gassi     | H         | +            | +         | -         |
|        |              | <i>Trifolium repens</i>     | White Clover/Batak nur  | H         | +            | -         | -         |
|        |              | <i>Trifolium fragiferum</i> | Clover/Batak laut       | H         | +            | +         | +         |
|        |              | <i>Melilotus alba</i>       | Sweet clover            | H         | -            | +         | -         |
|        |              | <i>Trigonella emodi</i>     | Himalayan Fenugreek     | H         | -            | -         | +         |
|        |              | <i>Cytisus scoparius</i>    | Common broom            | S         | +            | +         | -         |
| 8.     | Pinaceae     | <i>Pinus helpensis</i>      | Aleppo Pine             | T         | -            | +         | -         |
|        |              | <i>Cedrus deodara</i>       | Himalayan cedar/Deodar  | T         | -            | +         | +         |
|        |              | <i>Pinus roxburghii</i>     | Chir pine/Chir          | T         | -            | +         | -         |
|        |              | <i>Pinus canariensis</i>    | Canary Island Pine      | T         | -            | +         | -         |
|        |              | <i>Pinus wallichiana</i>    | Blue pine/Kail          | T         | -            | -         | +         |
| 9.     | Cupressaceae | <i>Cupressus torulosa</i>   | Bhutan cypress/Sarvikul | T         | +            | +         | -         |

Contd...

223

224

225

| S. No. | Family          | Species                          | Common name/Local name             | Life form | Altitude (m) |           |           |
|--------|-----------------|----------------------------------|------------------------------------|-----------|--------------|-----------|-----------|
|        |                 |                                  |                                    |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 10.    | Rhamnaceae      | <i>Ziziphus vulgaris</i>         | Zizyphus/Bre                       | S         | -            | -         | +         |
| 11.    | Berberidaceae   | <i>Berberis lycium</i>           | Indian barberry/Kawdach            | S         | +            | +         | -         |
| 12.    | Oleaceae        | <i>Jasminium humile</i>          | Yellow Jasmine                     | S         | +            | +         | +         |
| 13.    | Polypodiaceae   | <i>Adiantum capillusveneris</i>  | Adiantum/Gew theer                 | H         | -            | -         | +         |
| 14.    | Thymelaeaceae   | <i>Daphne oleoides</i>           | Dafne spatolata                    | S         | -            | +         | +         |
| 15.    | Zygophyllaceae  | <i>Tribulus terretris</i>        | Puncture Vine/Mister kund          | H         | +            | +         | +         |
| 16.    | Caryophyllaceae | <i>Dianthus angulatus</i>        | Himalayan Pinks                    | H         | +            | +         | -         |
|        |                 | <i>Phytolacca acinosa</i>        | Indian Poke                        | H         | -            | -         | +         |
| 17.    | Convolvulaceae  | <i>Cuscuta europaea</i>          | Devil's hair/Wozul kukli poot      | H         | +            | +         | -         |
|        |                 | <i>Cuscuta cuspidata</i>         | Golden thread/Kokil pot            | H         | -            | +         | -         |
|        |                 | <i>Convolus arvensis</i>         | Bindweed/Soi posh                  | H         | -            | +         | -         |
| 18.    | Iridaceae       | <i>Iris nepalensis</i>           | Graceful himalayan iris/Mazar mond | H         | +            | +         | +         |
| 19.    | Caprifoliaceae  | <i>Lonicera quinquelocularis</i> | Translucent Honey suckle/Pakhur    | H         | +            | +         | +         |
| 20.    | Polygonaceae    | <i>Rumex acetosa</i>             | Sheep's sorrel/Choki chen          | H         | +            | +         | +         |

226

Contd...

227

228

| S. No. | Family          | Species                      | Common name/Local name   | Life form | Altitude (m) |           |           |
|--------|-----------------|------------------------------|--------------------------|-----------|--------------|-----------|-----------|
|        |                 |                              |                          |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 21.    | Ophioglossaceae | <i>Rumex orientalis</i>      | Spinach dock/Jungli abuj | H         | +            | -         | +         |
| 22.    | Asteraceae      | <i>Ophioglossum vulgatum</i> | Adder's Tongue/Chonchur  | H         | +            | +         | +         |
|        |                 | <i>Cichorium intybus</i>     | Chicory/Handi posh       | H         | +            | +         | -         |
|        |                 | <i>Taraxacum officinale</i>  | Dandelion/Hand           | H         | +            | +         | +         |
|        |                 | <i>Artemisia absinthium</i>  | Worm wood/Tethwan        | H         | +            | +         | +         |
|        |                 | <i>Tragopogon pratensis</i>  | Meadow Salsify           | H         | -            | +         | +         |
| 23.    | Violaceae       | <i>Cardus nutans</i>         | Musk thistle             | H         | -            | +         | +         |
| 24.    | Apiaceae        | <i>Viola odorata</i>         | Wood violet/Bunafshah    | H         | +            | +         | +         |
|        |                 | <i>Foeniculum vulgare</i>    | Common fennel/Baidanii   | H         | +            | +         | -         |
|        |                 | <i>Cuminum cyminum</i>       | Cumin/Zur                | H         | +            | +         | -         |
| 25.    | Poaceae         | <i>Cynodon dactylon</i>      | Dūrvā grass/Dramun       | H         | +            | +         | +         |
|        |                 | <i>Panicum crusgalli</i>     | Cockspur/Hama            | H         | +            | +         | -         |
|        |                 | <i>Stipa sibirica</i>        | Stipa sibirica           | H         | +            | -         | -         |
|        |                 | <i>Eragrostis nigra</i>      | love grass               | H         | -            | -         | +         |
|        |                 | <i>Hordeum murinum</i>       | False barley/Pingi       | H         | -            | -         | +         |

229

Contd...

230

231

| S. No. | Family         | Species                     | Common name/Local name    | Life form | Altitude (m) |           |           |
|--------|----------------|-----------------------------|---------------------------|-----------|--------------|-----------|-----------|
|        |                |                             |                           |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 26.    | Chenopodiaceae | <i>Sorghum halepense</i>    | Aleppo grass              | H         | -            | +         | +         |
|        |                | <i>Poa angustifolia</i>     | Meadow-grass              | H         | -            | +         | +         |
|        |                | <i>Chenopodium album</i>    | Lamb's quarters/Lachij    | H         | +            | +         | -         |
| 27.    | Urticeae       | <i>Urtica dioca</i>         | Nettle/Soi                | H         | +            | +         | -         |
| 28.    | Labiatae       | <i>Salvia moorcroftiana</i> | Kashmir Salvia/Sholer     | H         | +            | +         | +         |
| 29     | Lamiaceae      | <i>Thymus serpyllum</i>     | Breckland thyme           | H         | +            | +         | +         |
|        |                | <i>Origanum vulgare</i>     | Oregano                   | H         | +            | -         | -         |
|        |                | <i>Nepeta cataria</i>       | Cat mint/Gandi soi        | H         | +            | +         | -         |
| 30.    | Boraginaceae   | <i>Lycopsis arvensis</i>    | Small bugloss/Handi gaasi | H         | -            | +         | -         |
| 31.    | Campanulaceae  | <i>Campanula colorata</i>   | Bell flower/Chari hakh    | H         | -            | +         | -         |
| 32.    | Malvaceae      | <i>Malva sylvestris</i>     | Blue Mallow               | H         | -            | +         | -         |
|        |                | <i>Malva album</i>          | Purple Mallow             | H         | -            | +         | +         |
|        |                | <i>Malva rotundifolia</i>   | Dwarf mallow/Sochal       | H         | -            | -         | +         |
|        |                | <i>Peganum harmala</i>      | Syrian Rue/Isband         | H         | +            | -         | -         |

232

Contd...

233

234

| S. No. | Family           | Species                            | Common name/Local name            | Life form | Altitude (m) |           |           |
|--------|------------------|------------------------------------|-----------------------------------|-----------|--------------|-----------|-----------|
|        |                  |                                    |                                   |           | 1575-1705    | 1705-1835 | 1835-1967 |
| 33.    | Apocynaceae      | <i>Vinca major</i>                 | Bigleaf periwinkle                | H         | -            | +         | +         |
| 34.    | Scrophulariaceae | <i>Linaria dalmatica</i>           | Balkan toadflax                   | H         | +            | +         | -         |
| 35.    | Plantaginaceae   | <i>Kickxia subsessilis</i>         | Branched Cancerwort               | H         | -            | +         | +         |
|        |                  | <i>Plantago lanceolata</i>         | Ribwort Plantain/Veuth gulla      | H         | -            | -         | +         |
| 36.    | Oxalidaceae      | <i>Oxalis corniculata</i>          | Creeping wood sorrel (Khati buti) | H         | -            | -         | +         |
|        |                  | <i>Oxalis</i> spp                  |                                   | H         | -            | -         | +         |
| 37.    | Hypericaceae     | <i>Hypericum perforatum</i>        | Amber                             | H         | -            | -         | +         |
| 38.    | Hamamelidaceae   | <i>Parrotiopsis jacquemontiana</i> | Parrotia/Hatab                    | S         | +            | -         | -         |
| 39.    | Cupuliferae      | <i>Quercus ilex</i>                | Holm oak                          | T         | -            | +         | +         |
| 40.    | Amaranthaceae    | <i>Amaranthus</i> spp              | Pigweed                           | H         | -            | +         | -         |

235 T = Tree, S = Shrub, H = Grasses and Herbs, + = Present, - = Absent




**Fig. 1: Community structure of Shankaracharya Reserve Forest**