

1 **Effect of arbuscular mycorrhizal fungi on the growth and development of**
2 *Citrus aurantium* L.

3
4 **Abstract**

5 *Citrus aurantium* L. is the most used rootstock in Morocco. The use of
6 arbuscular mycorrhizal fungi can favor the good growth of this species. A study
7 was conducted under the greenhouse to assess its reaction to mycorrhizal
8 colonization. *Citrus aurantium* L. plants were inoculated at the stage of 2 leaves,
9 with a local composite inoculum of arbuscular mycorrhizal fungi, originating
10 from the rhizosphere of many species of Citrus. The growth and mycorrhizal
11 parameters were observed seven months after inoculation. The results obtained
12 showed that the inoculated plants had a growth significantly higher than the
13 control ones, with a gain of 90.1%. Inoculation of the plants is also effective on
14 the leaves emission, branches formation and growth of collar diameter, gains
15 were respectively 266, 300 and 300%. The fresh weight of root and vegetative
16 mass also showed higher values on the inoculated plants, with a gain of 478%
17 and 425% respectively. The average frequency and intensity of root colonization
18 by mycorrhizal fungi are 93.33% and 50.6% respectively. The number of spores
19 formed at the rhizosphere of plants inoculated is 132 / 100 g of soil and they are
20 represented by 23 species belonging to the genera *Glomus* (8 species),
21 *Acauolospora* (9 species), *Scutellospora* (3 species), *Pacispora* (2 species) and
22 *Gigaspora* (1 species) .The beneficial effects of mycorrhizal *Citrus aurantium*
23 L. are discussed in this study.

24 Keywords: Morocco, *Citrus aurantium* L., rootstock, mycorrhizal fungi (AMF),
25 growth, biomass.

Introduction

In Morocco, the citrus fruit sector has a great socio-economic importance. It represents a source of major currency estimated at 3 billion DH a year (MAPM, 2014), contributes substantially to improving farmers' incomes and generates over than 21 million working days per year (MAPM, 2014). However, the Moroccan citrus industry knows several problems related to diseases, pests (Vanderweyenne, 1974; Abbassi, 1990; Benyahia, 1998 et 2007; Smaili et al., 2013; Boudoudou et al., 2015) and various abiotic environmental stresses, case of salinity, low temperatures and periods of drought (Benyahia, 2007; Castle, 1993).

To cope with these biotic and abiotic constraints, the use of door resistant grafts remains the most convenient and the most recommended way. The bitter orange is the most used rootstock in Morocco, due to its capacity to adapt to different types of soil, its resistance to limestone, its good affinity for grafting with almost all varieties of citrus and its good resistance to salinity and to *Phytophthora* sp. diseases, as well as the good quality of fruits (Vanderweyen, 1982; Nadori et al., 1988; Castle et al., 1993; Benyahia , 1998 , 2007, 2004; Boudoudou et al., 2015).

Strengthening the resistance of the bitter orange is possible by the use of biotechnological techniques in the nursery which allows having *Citrus aurantium* L. seedlings more robust and also resistant to pathogenic organisms and water stress after transplantation. Several studies have indeed shown in other plant species that the mycorrhizal symbiosis allow more efficient uptake of water and nutrients, especially the phosphorus (Nouaim et Chaussod, 1996; Giri et al., 2003; Citeresi et al., 2008; Simard, 2014) and reduce the adverse effects of the biotic and abiotic environment constraints (Evelin et al., 2009; Porcel et al., 2006).

In this sense and to exploit the beneficial effects of endomycorrhizae, it is advisable to use in the citrus orchards vigorous rootstocks, previously mycorrhizal in nurseries. Thus, this work aims to study the effects of arbuscular mycorrhizal fungi (AMF) on growth parameters and root development of *Citrus aurantium* L. (bitter orange).

Material and methods

Plant material

59 *Citrus aurantium* L. seeds were disinfected with sodium hypochlorite; Seedlings are grown in
60 plastic pots filled with a mixture of peat and sterile sand. Sterilization is performed in an oven
61 at 250° C for 2 hours to remove the soil micro flora. All pots were placed in a greenhouse
62 until the stage of two true leaves, and regularly watered with tap water.

63 **Inoculum production**

64 Barley (*Hordeum vulgare*), mycotrophic plant, was chosen for the production of a composite
65 arbuscular mycorrhizal fungi inoculum. Barley grains were disinfected with sodium
66 hypochlorite at 5% for 2 minutes, and put to germinate in plastic bowls filled with a mixture
67 of disinfected sand and soil from the rhizosphere of a large number of species of the genus
68 Citrus. The rhizospheric soil used contains 70 number of endomycorrhizal species.

69 After four weeks of culture, the frequency and intensity of mycorrhizal barley roots were
70 estimated using the method of Hyman and Philips (1970) and these mycorrhizal roots were
71 used as endomycorrhizal inoculum.

72 **Inoculation with mycorrhizal**

73 The plants of *Citrus aurantium* L. were transplanted in pots filled with disinfected sand of
74 Mamora and containing fragments of barley mycorrhizal roots.

75 The pots are then placed in the greenhouse; Control and mycorrhizal plants were regularly
76 watered with distilled water.

77 **Experimental set**

78 The experimental protocol is carried out randomly; two batches of plants were performed
79 with five plants for each batch.

80 Lot 1: Control plants (T) have not undergone any treatment

81 Lot 2: Plants inoculated with mycorrhizal fungi (Myco).

82 The pots were then placed in a greenhouse for seven months, June to December, at a
83 temperature varying from 21 to 27 °C.

84 After seven months of culture, the plants of *Citrus aurantium* L. were cut at the root collar.

85 The roots were washed with tap water and dried on paper towels overnight under ambient
86 laboratory conditions. The height of the vegetative part was measured. The fresh weight of

the aerial part and root biomass were measured using a numerical scale. The diameter of the rod was measured with caliper scales and the number of branches on the vegetative part was counted.

Mycorrhizal rates inside roots

The roots observation was prepared according to the method of Philips and Haymann (1970). They were first washed with water; the finest roots were then cut into a length of 1 cm then immersed in a solution of 10% KOH (potassium hydroxide) and placed in the water bath at 90° C for one hour to eliminate cytoplasmic contents. At the end of this period, roots were rinsed and transferred in a solution of H₂O₂ (hydrogen peroxide) for 20 min at 90° C in the water bath until the roots became white. Roots were then rinsed, after this; they were dyed with cresyl blue, at 90° C for 15 min.

Evaluation of mycorrhizal rate

After eight months of culture, evaluation of mycorrhizal parameters was conducted by observing thirty root fragments of about 1 cm, randomly chosen to quantify mycorrhizae (Trouvelot et al., 1986; Amir et Renard, 2003; Kormanik et Mc Graw, 1982). The mycorrhizal frequency (F%), the mycorrhizal intensity (M%), the arbuscular content (TA) and the vesicular content (TV) of AMF inside the root bark are measured by assigning a mycorrhizal index ranging from 0 to 5 (Derkowska et al., 2008).

Spore's extraction

The wet sieving method described by Gerdemann and Nicholson (1963) was adopted to extract the spores from soil of *Citrus aurantium* L., the samples of rhizospheric soil were taken at random from each pot. Species identification is performed basing on macroscopic and microscopic observation of morphological characters and referring to the key determination defined by Schenk and Perez (1990) and the website of INVAM.

Statistical Analysis

Statistical analyzes were performed by analysis of variance by the ANOVA test at the 5% level using the STATISTICA software.

Results

After seven months of growth, inoculation of seedlings of *Citrus aurantium* L. with endomycorrhizal composite inoculums marked a positive effect (Figure 1) on all measured parameters. (Table 1) A clear improvement in the height of the inoculated plants was recorded with a gain of more than 90.1%. Similarly, inoculating the plants is also effective on the leaves show, the formation of branches and growth of the diameter of the collar, the gain is respectively 266%, 300% and 300% (Figure 2).

Mycorrhiza was also beneficial for the development of the root system and the vegetative mass. After seven months of growth, the weight of the root and the aerial mass of inoculated plants were larger than those of the control, respectively with an increase of 478% and 425%.

The mycorrhizal frequency is about 93.3% and the mycorrhizal intensity is of 50.6%.(Figure 3). Moreover, different structures characterizing arbuscular endomycorrhizae were observed in inoculated roots: arbuscules, vesicles and intra and extracellular hyphae (Figure 4).

The arbuscular and vesicular contents of inoculated roots is about 40.83% and 3.36% respectively % (Figure 3). However, the roots of non-inoculated plants are not mycorrhizal.

The number of spores is in the order of 132 spores /100g of soil. % (Figure 5). Preliminary identifications (Table 2) have allowed to note that isolated spores belong to 23 species (Figure 6): *Pacispora boliviana*, *Acaulospora colossica*, *Acaulospora* sp1, *Acaulospora koskei*, *Acaulospora* sp2, *Acaulospora laevis*, *Pacispora robiginia*, *Glomus aggregatum*, *Acaulospora* sp3, *Acaulospora foveta*, *Scutellospora* sp1, , *Scutellospora castanea*, *Glomus deserticola*, *Gigaspora* sp, *Glomus spinuliferum*, *Glomus fecundisporum*, *Glomus intraradices*, *Scutellospora* sp2, *Glomus boreale*, *Glomus ambisporum*, *Glomus fecundisporum* , *Acaulospora gedanensis*, *Acaulospora nicolsonii*.

According to the classification of Oehl and Sieverding (2003), these species are distributed in 5 genera (*Glomus*, *Gigaspora*, *Acaulospora*, *Pacispora*, *Scutellospora*) and 5 families (*Glomaceae*, *Gigasporaceae*, et *Acaulosporaceae*, *Pacisporaceae* , *Scutellosporaceae*,) and 2 orders (*Glomerales*, *Diversisporales*).

Glomus intraradices, *Glomus deserticola*, *Acaulospora foveta*, *Glomus aggregatum* are respectively the most dominant species in the rhizosphere of the *Citrus aurantium* L. inoculated plants, their frequency of occurrence (Figure 7) vary between 6.81 % and 11.36%.

Similarly, the isolated species are represented by 38% of *Glomus*, 37% of *Acaulospora*, 13% of *Scutellospora*, 8% and 4% of *Pacispora* and *Gigaspora* (Figure 8).

Discussion

The surveys carried out in the rhizosphere of the inoculated *Citrus aurantium* L. plants showed that these roots contain endomycorrhizal structures: Vesicles, arbuscules, internal and external hyphae and endophytes. The presence of these structures characterizing endomycorrhizae allows to classify *Citrus aurantium* L. as a mycotrophic species (Artib et al., 2016; Wang, 2012; Camprubi et Calvet, 1995).

The plants inoculated with the composite endomycorrhizal inoculum were clearly better compared to the control plants in terms of leave's number and stem diameter, which confirms

the results of Lahlali (2001), plant heights, the same result was obtained by Syversten et Graham 1999 and Menge J.A 1978, The ability of AMF to stimulate growth is confirmed by the positive influence on aerial and root biomasses, which is in accordance with results obtained by (Farih *et al.*, 1988, Nouaim et Chaussod, 1996; Giri *et al.*, 2003; Citernesi *et al.*, 2008 ; Mattew *et al.*,1999 ;Gonzalez et ferrera, 2001).

The influence of these fungi on the root development is perhaps due to increased absorption of phosphorus that may cause proliferation and cell elongation (Black, 1965). This root development is due to the formation of a greater number of rootlets, which confirms that the AM fungi increase the rooting zone (Boureima *et al.*, 2008).

The beneficial effect of endomycorrhizal fungi on the growth of *Citrus aurantium* L. has been demonstrated; this is in agreement with the works of Mwangi *et al.* (2009) and Chliyah *et al.* (2014) they also demonstrated that the inoculation of tomato plants with AM fungi stimulated the weight and length of the aerial and root parts of these plants.

The AMF have a positive effect on the growth and development of the roots of *Lycium europaeum* plants (Touati *et al.*, 2015), the olive tree (Chliyah *et al.*, 2014 ; Porras Soriano *et al.*, 2005 ; Estaun *et al.*, 2003 ; Ganz *et al.*, 2002 ; Porras Soriano *et al.*, 2009 ; Sidhoum *et al.*, 2013), date palm tree (Sghir *et al.*, 2014 ; Smith et Read, 1997 ; Janos 1977), *Retama monosperma* (Hatimi *et al.*,1997), *Argania spinosa* L.(Bousselmame *et al.*,2002) ,and Carob tree (Talbi *et al.*,2016).

Growth of *Citrus aurantium* L. is dependent on mycorrhiza (Nemec, 1991, Farih *et al.*, 1988), The growth observed in *Citrus aurantium* L. could be explained by the ability of these fungi to colonize roots more rapidly and by improving the nutritional intake of plants, especially by increasing the minerals absorption (Eden, 1965; Smith *et al.*, 1986; George *et al.*, 1992; Toro *et al.*, 1998), and the capacity of plants to explore more space in soil (Joner *et al.*, 2000).

Indeed, endomycorrhizae allow the transport of the little mobile nutrients in the soil to the plant, especially, phosphorus (Rhodes & Gerdemann, 1975; Bolan, 1991; Gianinazzi & Schüepp, 1994; Smith & Read, 1997; Duponnois *et al.*, 2005; Lambers *et al.*, 2008 ; Strullu, 1991; Harrison, 1999), and improve drought tolerance (Ruiz-Lozano et Azcón, 1995), and induce in them morphological and physiological transformations to tolerate the environmental constraints (Porcel *et al.*, 2006).

In this work, 23 species of (*Glomerales*, *Gigasporales*, *Acaulosporales*, *Diversisporales*) were detected in the rhizosphere of *Citrus aurantium* L. treated with endomycorrhizal composite inoculum constituted by these genera, however, 2 species of *Glomerales* have been reported in the rhizosphere of *Citrus aurantium* L. grown in the field, Artib *et al.* (2016) and Wang (2012) reported only 12 species in the citrus rhizosphere in southern China. In the

rhizosphere of other plant species, Abbas *et al.* (2006) reported 6 species of arbuscular mycorrhizal fungi (AMF) in seven Moroccan Tetraclines. Tellal *et al.* (2008) noted 10 species in the rhizosphere of *Casuarina cunninghamiana* and *Casuarina glauca* developing in 15 sites and two nurseries in Morocco. Bouamri *et al.* (2006) and Sghir *et al.* (2015) revealed 15 and 9 species in the Tafilalt and Zagora palm rhizosphere. In Central Europe, Oehl *et al.* (2003) identified 12 species in the rhizosphere of the vine. In Jordan, Mohammad *et al.* (2003), Kachkouch *et al.* (2014) isolated 6 species and 5 genera respectively in the rhizosphere of the olive tree. Talbi *et al.*, 2015 reported 31 species in the Carob tree rhizosphere.

The count of the spores of the mycorrhizal fungi showed a dominance of the genus *Glomus*. This dominance was also found in the citrus rhizosphere (Wang, 2012) and in the rhizosphere of *Citrus aurantium* L. Artib *et al.* (2016), the olive-tree rhizosphere (Kachkouch *et al.*, 2012 ;2015), Oleaster (Sghir *et al.*, 2013), date palm (Sghir *et al.*, 2015).

The root mycorrhizal frequency is very high in inoculated *Citrus aurantium* L. (93.33%) with an important spore number of 132/100g of soil. However, previous results showed a mycorrhization frequency of (73%) and a number of spores of 48 per 100 g of soil (Artib *et al.*, 2016).

Plants products in nurseries are provided with a few vigorous root systems and slightly structured and cannot, therefore, withstand biotic and abiotic stresses they face after transplantation. From this study, and after the production of a complex of native fungi, composed of several species, it can be concluded that the plants of *Citrus aurantium* L. inoculated with AMF present another way of promoting the growth of this plant which can be used as biological means to tolerate stresses. Indeed, its establishment is interesting for nurseries, since in the USA, inoculation with AMF of Citrus grown in nurseries has become a common practice (Menge *et al.*, 1978).

Conclusion

From the whole of this study, it is clear that the endomycorhization (AMF) of plants of *Citrus aurantium* L. has a beneficial effect on the growth parameters. The positive effects obtained with the rootstock following the inoculation can therefore be explained by the great exploitation of the soil by the mycelium, This effect is even important for *Citrus aurantium*

L. a perennial species of great economic and nutritional importance, usually cultivated in soils, with mineral potential, often low, and found in forms little assimilable by the roots.

The rootstock inoculation by endomycorrhizal fungi has several advantages: Substantially shortens the development time of the plants, significantly reduces the application of chemical fertilizers, and has the advantages of providing quality products.

The increased performance of mycorrhizal plants in digesting food from the soil and resisting environmental stresses gives fungi symbiotes a role as bio fertilizers and crop protection agents. At the same time, adequate management of mycorrhizae in agricultural environments would extend the quality and sustainability of soils, while protecting the environment in the long term and reducing production costs.

References

- **Abbas Y., Ducouso M., 2006.** Diversity of arbuscular mycorrhizal fungi in *Tetraclinis articulata* (Vahl) Masters Woodlands in Morocco. Ann. For. Sci, 63 : 285– 291.
- **Abbassi M., 1990.** Contribution à la lutte biologique contre les homoptères ravageurs des Citrus au Maroc. Thèse de Doctorat d'Etat. Faculté des Sciences Agronomiques de Gembloux, Belgique , p 185.
- **Amir H., Renard A., 2003.** Etude microbiologique générale de quelques sols de forêts sclérophylles de Nouvelle-Calédonies : Statuts des mycorrhizes à arbuscules. PCFS-UNC N° 54/2003/CP , p 22.
- **Artib M., Chliyah M., Touati J., Talbi Z., Selmaoui K., Ouazzani Touhami A., Benkirane R., and Douira A., 2016.** Study of Arbuscular mycorrhizal fungi diversity in the rhizosphere of citrus grown in Morocco. IJAPBC, 5(3): 2277 – 4688.
- **Benyahia H., Ait Haddou M. M., Jriifi A., Lamsettef Y., 2004.** Effet de la salinité de l'eau d'irrigation sur la colonisation des racines des porte-greffes d'agrumes par *Phytophthora parasitica*. Fruits, 59: 101–108.
- **Benyahia H., 1998.** Effet de la salinité sur le développement de la pourriture racinaire des agrumes. Thèse de 3 ème cycle, Université. CADI AYAD, Faculté des sciences Semlalia, Marrakech , p 160.
- **Benyahia H., 2007.** Amélioration de la résistance des porte-greffes d'agrumes vis-à-vis des contraintes biotiques et abiotiques. Faculté des Sciences Dhar El Mahraz - Fès - Doctorat National.
- **Black C. A., 1965.** Methods of soil analysis. Am. Soc. Agron, Wisconsin, USA, 9: 1114-1162.

- 254 • **Bolan N.S., 1991.** A critical review on the role of mycorrhizal fungi in the uptake of
255 phosphorus by plants. *Plant and Soil*, 134: 189-207.
- 256 • **Bouamri B., Dalpé Y., 2006.** Arbuscular Mycorrhizal fungi species associated with
257 rhizosphere of *Phoenix dactylifera* l. in Morocco. *African Journal of Biotechnology*,
258 5(6) : 510-516.
- 259 • **Bouamri B., Dalpé Y., 2006.** Arbuscular Mycorrhizal fungi species associated with
260 rhizosphere of *Phoenix dactylifera* l. in Morocco. *African Journal of Biotechnology*.
261 5(6): 510-516.
- 262 • **Boudoudou D., Fadli A., Talha A., Bourachdi Y., Douira A., Benyahia H., 2015.**
263 Effect of seasonal and citrus rootstocks on inoculum density of *Phytophthora* spp. in
264 Citrus orchard in a heavy soil of the Gharb region of Morocco. *Biolife*, 3(2):367-377.
- 265 • **Boureima S., Diouf M., Diop T.A., Diatta M., Leye E.M., Ndiaye F., Seck D.,**
266 **2008.** Effects of arbuscular mycorrhizal inoculation on the growth and the
267 development of sesame (*Sesamum indicum* L.). *African Journal of Agricultural*
268 *Research*, 3 (3): 234-238.
- 269 • **Bousselmame F., Kenny L., Achouri M., 2002.** Effet des mycorhizes à vésicules et
270 arbuscules sur la croissance et la nutrition de l'arganier (*Argania spinosa* L.). *Actes*
271 *Inst. Agron Vet*, 22 (4) : 193-198.
- 272 • **Camprubí A., Calvet C., Estaun V., 1995.** Growth enhancement of Citrus reshni
273 after inoculation with *Glomus mosseae* and *Trichoderma aureoviride* and associated
274 effects on microbial populations and enzyme activity in potting mixtures. *Plant and*
275 *soil*, 173: 233- 238.
- 276 • **Castle W.S., Tucker D.P.H., Kiezdorn A.H., Youtsey C.O., 1993.** Rootstocks for
277 Florida Citrus. University of Florida Institute of Food and Agricultural Sciences John
278 T. Woeste. Dean for Extension.
- 279 • **Chliyeh M., Ouazzani Chahdi A., Selmaoui K., Ouazzani Touhami A., Filali-**
280 **Maltouf A ., El Modafar C., Moukhli A., Oukabli A ., Benkirane R., Douira A.,**
281 **2014.** Effect of *Trichoderma harzianum* and arbuscular mycorrhizal fungi against
282 *Verticillium* wilt of tomato. *International Journal of Recent Scientific Research*, 5 (2):
283 449-459.
- 284 • **Citernes A. S., Vitagliano C., 2008.** Plant growth root system morphology of *olea*
285 *europaea* L. Rooted cuttings as influenced by arbuscular mycorrhizas. *S. J. Horticult*
286 *Sci. Biotechnol*, 73(5): 647-654.
- 287 • **Citernes A.S., Vitagliano C., 2008.** Plant growth root system morphology of *olea*
288 *europaea* L. Rooted cuttings as influenced by arbuscular mycorrhizas. *S. J. Horticult*
289 *Sci. Biotechnol*, 73(5): 647-654.

- 290 • **Derkowska E., Sas Paszt L., Sumorok B., Szwonek E., Gluszek S., 2008.** The
291 influence of mycorrhization and organic mulches on mycorrhizal frequency in apple
292 and strawberry roots. *Journal of Fruit and Ornamental Plant Research*, 16: 227-242.
- 293 • **Duponnois R., Colombet A., 2005.** The mycorrhizal fungus *Glomus intraradices* and
294 rock phosphate amendment influence plant growth and microbial activity in the
295 rhizosphere of *Acacia holosericea*. *Soil Biol. Biochem*, 37: 1460- 1468.
- 296 • **Eden T., 1965.** Tea.2nd ed. Longmans, London, p 205.
- 297 • **Estaun V., Camprubi A., Calvet C., Pinochet J., 2003.** Nursery and field response
298 of olive trees inoculated with two arbuscular mycorrhizal fungi, *Glomus intraradices*
299 and *Glomus mosseae*. *J Am Soc Hortic Sci*, 128: 767-775.
- 300 • **Evelin H., Kapoor R., Giri B., 2009.** Arbuscular mycorrhizal fungi in alleviation of
301 salt stress. *Annals of Botany*, 104: 1263–1280.
- 302 • **Farih A., Nadori E.B., Nhami A., Boukhriss H., Walidi L.D., 1988.** Réponse à la
303 mycorrhization de 4 portes greffes d'agrumes élevés dans trois substrats différents. *Al*
304 *Awamia*, 64:55-62.
- 305 • **Ganz T.R., Kailis S.G., Abbot L.K., 2002** .Mycorrhizal colonization and its effect
306 on growth, P uptake and tissue phenolic content in the European olive (*Olea europaea*
307 L.). *Adv Hortic Sci*, 16 (3-4): 109-116.
- 308 • **George E., Haussler K.U., Vetterlein D., Gorgus E., Marschner H., 1992.** Water
309 and nutrient translocation-by hyphae of *Glomus mosseae*. *Can. J. Bot*, 70: 2130-2137.
- 310 • **Gerdemann J .W., Nicholson T.H., 1963.** Spores for mycorrhizal endogone species
311 extracted from soil by wet sieving and decanting. *Trans. Br. Mycol. Soc*, 46: 235-244.
- 312 • **Gianinazzi S., Schüepp H., 1994.** Impact of arbuscular mycorrhizas on sustainable
313 agriculture and natural ecosystems. Birkhäuser, Basel.
- 314 • **Giri B., Kapoor R., 2003.** Influence of arbuscular mycorrhizal fungi and salinity on
315 growth, biomass, and mineral nutrition of *Acacia auriculiformis*. *Biology and Fertility*
316 *of soils*, 38: 170-175
- 317 • **Giri B., Kapoor R., 2003.** Influence of arbuscular mycorrhizal fungi and salinity on
318 growth, biomass, and mineral nutrition of *Acacia auriculiformis*. *Biology and Fertility*
319 *of soils*, 38: 170-175.
- 320 • **Gonzalez C., Ferrera C., 2001.** Roca fosforica y *Glomus* sp. En el crecimiento de
321 naranjo agrico rock phosphate and *Glomus* sp.On sour orange growth .*Terra*,
322 18(4):362-366.
- 323 • **Harrison M.J., 1999.** Interfaces and nutrient transport in plant / fungal symbioses.
324 *Journal of Experimental Botany*, 50: 1013-1022.
- 325 • **Hatimi A., Achouri M., Oihabi A., 1997.** Endomycorhization de légumineuses
326 fixatrices des dunes: Croissance et nutrition phosphatée. *Cahiers «Sécheresse»*, 8(2):
327 99-102.
- 328 • **Janos D. P., 1977.** Vesicular-arbuscular mycorrhizae affect the growth of *Bactris*
329 *gasipaes*. *Principes*, 21: 12–18.

- 330 • **Joner E. J., Johansen A., 2000.** Phosphatase activity of external hyphae of two
331 arbuscular mycorrhizal fungi. *Mycol Res*, 104: 81-86.
- 332 • **Kachkouch W., Ouazzani Touhami A., Filali Maltouf A., El Modafar C.,**
333 **Moukhli A., Oukabli A., Benkirane R., and Douira A., 2013.** Arbuscular
334 mycorrhizal fungi species associated with rhizosphere of *Olea europaea* L. in
335 Morocco. *Journal of Animal & Plant Sciences*, 15(3): 2275-2287.
- 336 • **Kormanik P.P., Mc Graw A.C., 1982.** Quantification of vesicular- arbuscular
337 mycorrhizae in plant roots. In *Methods and principles of Mycorrhizal Research*. N.C.
338 editor. APS Press, p 37.
- 339 • **Lahlali R., 2001.** Amplification en aeroponie de trois espèces mycorrhiziennes et
340 mycorrhization de certaines variétés de portes greffes d'agrumes. Mémoire de Troisième
341 cycle en agronomie. ENA. Meknes.
- 342 • **Lambers H., Raven J.A., Shaver G.R., Smith S.E., 2008.** Plant nutrient-acquisition
343 strategies change with soil age. *Trends in ecology & evolution*, 23(2):95-103.
- 344 • **MAPM: (Ministère de l'agriculture et de la pêche maritime), 2014.** Fellah-
345 trade. <http://www.fellah-trade.com/fr/info-filiere/chiffres-cles/fruits-et-legumes>
346 (*consulté* le 25/08/2016).
- 347 • **Matthew W., Fidelibus ., Chris A., Martin., 1999.** Growth of *Citrus volkameriana*
348 inoculated with AM fungi in moist or periodically dries soils. University of Arizona
349 college of Agriculture Citrus Research Report.
- 350 • **Menge J .A., Johnson L.V., Platt R.G., 1978.** Mycorrhizal dependency of several
351 citrus cultivars under three nutrient regimes. *Departments Plant Pathology and Plant*
352 *Science*, University of California, Riverside, California 92521, U.S.A. *New Phytol*,
353 81:553-559.
- 354 • **Menge J. A., Johnson L.V., Platt R.G., 1978.** Mycorrhizal dependency of several
355 citrus cultivars under three nutrient regimes. *Departments Plant Pathology and Plant*
356 *Science*, University of California, Riverside, California 92521, U.S.A. *New Phytol*,
357 81:553-559.
- 358 • **Mohammad M.J., Hamad S.R., and Malkawi H.I., 2003.** Population of arbuscular
359 mycorrhizal fungi in semi-arid environment of Jordan as influenced by biotic and
360 abiotic factors. *J. Arid Environ*, 53: 409-417.
- 361 • **Mwangi M .W., Monda E. O., Okoth S. A., Jefwa J .M., 2011.** Effect of
362 *Trichoderma harzianum* and arbuscular mycorrhizal fungi on growth in tomato
363 (*Lycopersicon esculentum* mill) seedlings, napier (*Pennisetum purpureum* l) and tea
364 (*Camellia sinensis* l) cuttings. *Tropical and Subtropical Agroecosystems*, 11: 423 –
365 429.
- 366 • **Nadori E.B., Ouammou M., Kayaf M., 1988.** Comportement de trois clones de
367 clémentinier sur différents porte-greffes. *Al Awamia*, 64: 44-54.

- 368 • **Nemec S., 1991.** Effects of soil –phosphorus and *Glomus intraradices* on growth,
 369 non structural carbohydrates, and photosynthetic activity of *Citrus aurantium*. Plant
 370 and soil, 128(2):257-263.
- 371 • **Nouaim R., Chaussod R., 1996.** Rôle des mycorhizes dans l'alimentation hydrique
 372 des plante, notamment des ligneux en zone arides .Options Méditerranéennes, 20: 9-
 373 26
- 374 • **Oehl F., Sieverding E., Ineichen K., Mader P., Boller T., Wiemken A., 2003.**
 375 Impact of land use intensity on the species diversity of arbuscular mycorrhizal fungi
 376 in agroecosystems of central Europe. Applied and Environmental Microbiology, 69:
 377 2816-2824.
- 378 • **Philips J. M., Hayman D. S., 1970.** Improved procedures for clearing root and
 379 staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of
 380 infection. Trans. Br. Mycol. Soc, 55 158-161.
- 381 • **Porcel R., Aroca R., Azcon R., Ruiz-Lozano J.M., 2006.** PIP aquaporin gene
 382 expression in arbuscular mycorrhizal Glycine max and Lactuca sativa plants in
 383 relation to drought stress tolerance. Plant Mol. Biol, 60: 389-404.
- 384 • **Porras Soriano A., Sorano Marintin M.L., Porras Piedra A., Azcon P., 2009.**
 385 Arbuscular mycorrhizal fungi increased growth, nutrient uptake and tolerance to
 386 salinity in olive trees under nursery conditions. Journal of Plant Physiology, 166:
 387 1350-1359.
- 388 • **Rhodes L.H., Gerdemann J.W., 1975.** Phosphate uptake zone of mycorrhizal and
 389 non- mycorrhizal onions. New phytologist, 75.555.
- 390 • **Ruiz Lozano J .M., Azcón R., 1995.** Hyphal contribution to water uptake in
 391 mycorrhizal plants as affected by the fungal species and water status. Physiol.
 392 Plantarum, 95: 472–478.
- 393 • **Schenck N C., Pérez Y., 1990.** Manual for the identification of VA mycorrhizal
 394 fungi. Synergistic Publications. Gainesville. Florida, p 286 .
- 395 • **Sghir F., Chliyah M ., Touati J., Mouria B ., Ouazzani Touhami A ., Filali**
 396 **Maltouf A., El Modafar C ., Moukhli A ., Benkirane R., Douira A., 2014.** Effect
 397 of a dual inoculation with endomycorrhizae and *Trichoderma harzianum* on the
 398 growth of date palm seedlings. Int. J. Pure App. Biosci, (6): 12-26.
- 399 • **Sghir F., Chliyah M., Kachkouch W., Khouader M., Ouazzani Touhami A.,**
 400 **Benkirane R and Douira A., 2013.** Mycorrhizal status of *Olea europaea* spp. oleaster
 401 in Morocco. Journal of Applied Biosciences, 61: 4478 – 4489.
- 402 • **Sghir F., Touati J., Mouria B., Ouazzani Touhami A., Filali Maltouf A., El**
 403 **Modafar C., Moukhli A., Benkirane R and Douira A., 2016.** Variation in
 404 pathogenicity of *fusarium oxysporum* f. sp. albedinis on two cultures associated with
 405 date palm of moroccan oasis, 2(3):56-68.

- **Sidhoum W., Fortas Z., 2013.** Effect of Arbuscular mycorrhizal fungi on growth of semi-woody olive cuttings of the variety "Sigoise" in Algeria. American Journal of Research Communication, 1(11): 244-257
- **Simard F., 2014.** Stimulation de la synthèse des composés nutraceutiques et aromatiques dans les fines herbes et les légumes par les champignons mycorhiziens à arbuscules. Université Laval. Québec, Canada.
- **Smaili M.C., Abbassi M., Boutaleb J.A., Blenzar A., 2013.** Richesse spécifique des ennemis naturels associés aux vergers d'agrumes au Maroc: Intérêt et implication pour la lutte biologique OEPP, 43 :155–166.
- **Smith S. E., John J .B., Smith F .A., Bromley J. L., 1986.** Effect of mycorrhizal infection on plant growth, nitrogen and phosphorous nutrition in glass house grown *Allium cepa* L. *New Phytol*, 103: 359- 373.
- **Smith S.E., Reed D.J., 1997.** Mycorrhizal symbiosis. London Academic Press.
- **Strullu D. G., 1991.** Mycorrhizes et développement des plantes. In les mycorrhizes des arbres et plantes cultivées. Tec et Doc. Lavoisier. Paris, p79-80.
- **Syverstsen J.P., Graham J.H., 1999.** Phosphorus supply and arbuscular mycorrhizas increase growth and net gas exchange of Citrus spp. growth at elevated (CO₂). Plant and Soil, 208: 209-219.
- **Talbi Z ., EL Asri A., Touati J ., Chliyah M ., Ait Aguil F ., Selmaoui k ., Sghir F ., Ouazzani Touhami A., Benkirane R and Douira A., 2015.** Morphological characterization and diversity of endomycorrhizae in the rhizosphere of Carob tree (*Ceratonia siliqua*) in Morocco. *Biolife*, 3(1):196-211.
- **Talbi Z., Chliyah M., Mouria B., El Asri A., Ait Aguil F., Ouazzani Touhami A., Benkirane R and Douira A ., 2016.** Effect of double inoculation with endomycorrhizae and *Trichoderma harzianum* on the growth of carob plants .IJAPBC, 5(1): 2277 – 4688.
- **Tellal M., 2008.** Contribution à l'étude de la symbiose Casuarina-microorganismes et son importance sur la production de plants en pépinière et la fertilité du sol. Thèse de Doctorat. Université Ibn Tofail, Fac. des Sciences, Kénitra, Maroc, 135p.
- **Toro M., Azcón R., Barea J .M., 1998.** The use of isotopic dilution techniques to evaluate the interactive effects of *Rhizobium* genotype, mycorrhizal fungi, phosphate-solubilizing rhizobacteria and rock phosphate on nitrogen and phosphorus acquisition by *Medicago sativa*. *New Phytol*, 138: 265–273.
- **Touati J., Chliyah M., Ouazzani Touhami A., Benkirane R., Douira A., 2015.** Effect of arbuscular mycorrhizal fungi on plant growth and root development of the boxthorn tree (*Lycium Europaeum*) under a greenhouse conditions. *Int. J. Pure App. Biosci*, 2 (6): 84-91.
- **Trouvelot A., Kough J .L., 1986.** Mesure du taux de mycorhization VA d'un système racinaire. Recherche de méthodes d'estimation ayant une signification fonctionnelle. In Physiological and genetical aspects of mycorrhizae. Gianinazzi-Pearson V. et Gianinazzi S. (Eds.), INRA édition, Paris, pp. 217-221.

- **Vanderweyen A., 1982** .Contribution à l'étude de la gommose à Phytophthora des agrumes au Maroc, Université. Nancy, Thèse, Nancy, France, 1982 ; p 160.
- **Wang P., Zhang J.J., Shu B., 2012.** Arbuscular mycorrhizal fungi associated with citrus orchards under different type of soil management southern china.Plant Soil Environ, (7):302-308.

Agronomic parameters	Inoculated plants	Control plants
Root fresh weight (g)	2.43	0.42
Fresh weight of the vegetative parts (g)	2.1	0.40
Height (cm)	21.3	11.2
Diameter of the rod (cm)	0.4	0.1
Number of leaves	22	6
Number of branches	3	0
Table. 1: Agronomic parameters of <i>Citrus aurantium</i> L. after 7 months of inoculation with endomycorrhizal fungi.		




Figure 1. Effect of endomycorrhizae on the growth of aerial and root parts of *Citrus aurantium* L. plants. Controls (A), Mycorrhizal plants (B).

462

463

464

465

466

467

468

469

470




Figure 2. Effect of mycorrhizal inoculation on RW: root weight, H: height, D: diameter, LN: leaf number, NB: number of branches, VW: vegetative weight




Figure 3. Evaluation of the mycorrhizal rate of the inoculated roots of *Citrus aurantium* L. mycorrhizal frequency (MF), mycorrhizal intensity (MI), arbuscular content (AC) and vesicular content (VC).




Figure 4: Roots of mycorrhizal *Citrus aurantium* L. arbuscules (a); Extra (eh) and intra-radicular (ih) hyphae, spores (s); Vesicles (v) and endophytes (en) (G × 400).

Number	Name	Form	Color	size	Wall thickness	Spore surface	Hyphe length
1	<i>P.boliviana</i>	globular	orange	70	2.5	granular	50
2	<i>A. colossica</i>	globular	mauve green	77.5	2.5	granular	-
3	<i>Acaulospora</i> sp1	globular	yellow	120	2.5	granular	-
4	<i>A. koskei</i>	globular	beige	107.5	7.5	smooth	-
5	<i>Acaulospora</i> sp2	globular	Yellow beige	105	7.5	smooth	-
6	<i>A. laevis</i>	globular	brown yellow	75	7.5	smooth	-
7	<i>P. robiginia</i>	globular	brown yellow	95	2.5	smooth	-
8	<i>G. aggregatum</i>	globular	brown yellow	80	7.5	smooth	-
9	<i>Acaulospora</i> sp 3	globular	yellow	72.5	2.5	smooth	-
10	<i>A.nicolsonii</i>	globular	yellow	112.5	7.5	smooth	83
11	<i>A. foveta</i>	globular	brown	50	12.5	granular	-
12	<i>Scutellospora</i> sp1	globular	beige	75	2.5	smooth	96.5
13	<i>G. deserticola</i>	globular	clear yellow	250	12.5	smooth	-

14	<i>S. castanea</i>	globular	Yellow brown	50	2.5	granular	-
15	<i>G. deserticola</i>	oval	clear yellow	112.5	7.5	smooth	50
16	<i>Gigaspora sp</i>	globular	clear Green	37.5	2.5	granular	-
17	<i>G. spinuliferum</i>	globular	Yellow darkened	112.5	7.5	smooth	50
18	<i>G.fecundisporum</i>	globular	brown yellow	200	10	smooth	-
19	<i>G. intraradices</i>	oval	Yellow darkened	60	7.5	smooth	-
20	<i>G. deserticola</i>	globular	Yellow orange	200	7.5	smooth	-
21	<i>G. intraradices</i>	globular	Yellow darkened	75	2.5	smooth	-
22	<i>Scutellospora sp2</i>	globular	brown	107.5	7.5	smooth	-
23	<i>G. boreale</i>	oval	brown yellow	62.8	2.5	smooth	23.6
24	<i>G. ambisporum</i>	globular	brown	70	7.5	smooth	83.2
25	<i>Glomus sp</i>	globular	clear yellow	75	7.5	smooth	-
26	<i>G. spinuliferum</i>	globular	Yellow darkened	112.5	7.5	smooth	-
27	<i>A. gedanensis</i>	globular	brown beige	115	7.5	smooth	-

Table 2. Identification of mycorrhizal fungi isolated from the rhizosphere of *Citrus aurantium*




Figure. 5: Average spore density of AM fungi of mycorrhizal and control plants














Fig 6: Endomycorrhizal fungi isolated from the rhizosphere of mycorrhizal *Citrus aurantium* L.



515



516

517 **Figure 7:** Isolation frequency of mycorrhizal species from the rhizosphere of *Citrus aurantium*

518


519

520

521

522

523



524

525

Fig 8: Distribution of the isolated genera from the rhizosphere of *Citrus aurantium* L.