

Original Research Article

**FIELD EVALUATION OF FOLIAR BLEND MICRONUTRIENT
FERTILIZER ON COTTON (*Gossypium sp*) PRODUCTION IN
KATSINA STATE, NIGERIA**

ABSTRACT

During the 2013 and 2014 cropping season, field experiments were conducted to evaluate the effect of combined application of NPK fertilizer and FoliarBlend micronutrient application on yield and growth of Cotton at *Funtua* (longitude 07° 19'E, latitude 11° 32'N), in Katsina State, Nigeria. This study was done as a randomized complete block design with three replications. The treatments were T1 (500mls of FoliarBlend in 20 litres of water); T2 (20kgNPK/ha + 500mls of FoliarBlend in 20 litres of water); T3 (40kgNPK/ha + 500mls of FoliarBlend in 20 litres of water); T4 (60kgNPK/ha + 500mls of FoliarBlend in 20 litres of water) and control. Soil samples were collected for the determination of chemical and physical properties before planting. Data on growth parameters (plant height, number of leaves, number of nodes, & number of branches) and yield parameters (number of bolls & yield) were also collected and subjected to 2 ways Analyses of Variance (ANOVA) and means separated with Least Significant Difference(LSD) at P=0.05.

The study revealed that during wet season of 2013 cropping system, the results from the field evaluations showed that combined application rate of 40kgNPK/ha + 500mls of FoliarBlend in 20litres of water produced the highest number of leaves (36), plant height (76cm), number of branches (21), and number of nodes (30), compared to the least value of 30, 59.6cm, 18, and 21 respectively obtained in the control at *funtua* location.

The study also revealed that application rate of 60kg NPK/ha + 500mls of FoliarBlend in 20 litres of water had the highest number of bolls (19) and yield (5.2 t/ha) compared to the least values of 13, and 2.13 t/ha obtained respectively in the control at *funtua* site.

Similarly in the 2014 dry season, the same trend as in the 2013 was obtained with the application rate of 60kg NPK/ha + 500mls of FoliarBlend in 20 litres of water, given the highest number of bolls (24) and yield (6.1 t/ha) compared to the least values of 15 and 2.75 t/ha obtained respectively in the control at *funtua* site.

Considering the overall result, it can be concluded that combined application rate of 40kgNPK/ha + 500mls of FoliarBlend in 20litres performed best with the highest values on the growth parameters, and while 60kgNPK/ha + 500mls of FoliarBlend in 20litres gave the highest yield and yield parameters than the sole application of either NPK or FoliarBlend.

Key words: NPK 15:15:15 fertilizer, FoliarBlend micronutrients, yield, cotton, Nigeria

Introduction:

Cotton (*Gossypium spp*) belongs to the family Malvaceae and it is usually considered to be an annual crop but can also be left in the field to continue to grow into a shrub. Cotton is a major cash crop and a veritable source of foreign exchange in the Northern part of Nigeria prior to the discovery of crude oil in commercial quantity (Ogunlela and Abed, 1982). It is the second most important commercial arable crop in the country ranking next to groundnut. It is produced almost entirely within the area bound by latitudes 7⁰ and 14⁰ N which comprises the country's savannah zone. The production of cotton in Nigeria was greater than all the other fibre crops like kenaf, jute, etc, as a result of the use of cotton lint in the textile industries and cotton seeds for manufacture of vegetable oil, while the residue is used to produce cotton seed cake used in feeding livestock.

An estimated 0.6 – 0.8 million hectares of land is put to cotton production annually in Nigeria. The major cotton producing areas in the country are Katsina, Kaduna, Kano, Sokoto, Plateau, Bauchi, Borno, Adamawa, Zamfara, Kebbi and Jigawa States. The Gongola valley (that cuts across Adamawa and Taraba states) and the Plateau in the north-eastern zone constitute some of the major cotton producing areas in Nigeria. Production of cotton depends on various factors ranging from weather, cotton price, soil and nutrient availability, pests & diseases and problems of the textile industries. Pest and diseases control is highly necessary to enhance good field performance of cotton. However, the increased in the prices of farm inputs especially fertilizer and poor pricing of cotton lint as a result of the collapse of textile industries in Nigeria brought about the decrease in production of cotton in Nigeria (Prentice, 1992). However, the present Government is trying to resuscitate the textile industries in the country; as such, mass production of cotton is imperative for the survival of these industries.

Judicious application of fertilizers may boost cotton production several folds (100 to 300%), provided the crop is planted early. Fertilizer trials have shown that the application of Nitrogen and Phosphorus fertilizers can increase cotton yields, provided that the crop is protected against insect pests (Faulkner, 1974). Cotton showed tremendous response to nitrogenous fertilizers in all soil types, but its response to phosphate fertilizer is erratic and variable in most soils (Malik *et al* 1996). However, nitrogen and phosphorus are highly mobile elements in the plant (Agbede, 2009). The phosphorus requirements of cotton are considered very low because of its deep root system and indeterminate growth habit (Muhammad, *et al* 2001). Also, the Information on foliar micronutrient fertilizer recommendation on cotton production in Nigeria is lacking. The aim of this study is therefore to develop the appropriate fertilizer formulation (inorganic and micronutrient) that is specific to cotton in soils of the cotton belt of North Western Nigeria.

Materials and Methods:

Characterization of the Study Area:

The study was conducted in three (3) locations within Katsina state namely: Funtua, Bakori and Faskari. The study was conducted during wet seasons of 2013 and dry- season of 2014 at Funtua (longitude 07° 19'E, latitude 11° 32'N), *Bakori* (longitude 07° 26'E, latitude 11° 33'N) and *Faskari* (longitude 07° 02'E, latitude 11° 43'N) in Katsina State, Nigeria.

The daytime temperature is 28-30 degrees. Climate of this area is arid with dry and hot summer. The mean annual rainfall is 1101.25mm and the growing period of 200 days. Rainfall pattern is bimodal with the major rainy season from May to October. The soils of the study area are classified mainly as Ferric luvisols according to FAO classification system (FAO, 1988) or Alfisols in USDA soil Taxonomy system (Soil Survey Staff, 1992). They are generally low in CEC owing to predominance of kaolinite in their clay fraction, low organic matter and inherent fertility. The soil characteristic of the experimental site is sandy loam in texture (Table 1).

Field Trial

Soil auger was used to collect surface soil samples (0 -15cm) from the experimental site before planting and processed for routine soil analysis. Field preparation was done mechanically with one ploughing and harrowing, this was followed by manual ridging and mapping out of the plots. The plot size measuring 5m x 5m = 25m² with alley of 1m left in between plots was used. The experimental design was randomized complete block design with three replications. The treatments were T1 (500mls of FoliarBlend in 20 litres of water); T2 (20kgNPK/ha + 500mls of FoliarBlend in 20 litres of water); T3 (40kgNPK/ha + 500mls of FoliarBlend in 20 litres of water); T4 (60kgNPK/ha + 500mls of FoliarBlend in 20 litres of water) and T5 (control).

An improved variety of cotton (*Gossypium sp*) SAMCORT 1 was planted on 90cm ridges , with intra- row spacing of 45cm. The seed rate was 4seeds per hole, giving a population density of 54,000 plants/ha and later thinned to two plants per stand. The inorganic fertilizer (NPK 15-15-15) was applied at 3WAP as basal application (by placing in holes 5cm and 8cm away from the plant) immediately after thinning and then top dress with Urea at 8 weeks after planting. The micronutrient was applied as 500 mls of foliar blend fertilizer in 20 litres of water by spraying evenly around the base of the crop and upward, to cover the underside of the leaf. Three applications of the FoliarBlend micronutrient fertilizer were carried out at vegetative stage immediately after NPK fertilizer application; flowering initiation and boll formation respectively. All agronomical practices were duly observed.

The growth parameters measured included plant height; number of leaves, number of nodes and number of branches, while the yield parameters were number of bolls & yield. The data were subjected to 2 ways Analyses of Variance (ANOVA) and means separated with Least Significant Difference(LSD) at P=0.05.

Results & Discussions:

Pre-Planting Soil Analysis

Results of physical and chemical properties of soils of Funtua are shown in Table 1.

The textural class of the soil is sandy loam with sand values of 580 g kg⁻¹; silt (220g kg⁻¹) and clay (150g kg⁻¹) (Table 1).The sandy nature of this soil is attributed to the granitic and gneissic origin of the parent materials. This contributes to low cationic retention and water holding capacity. The pH (water) is 6.0 suggesting that the soils may not have acidity problems and therefore will not require liming. The organic carbon content is low, with a value of 4.20 g kg⁻¹. While the total nitrogen and available phosphorus values were far less than the moderate fertility class. The Exchangeable cations (Ca, Mg, K and Na) values are moderate (Table 1).

Cation exchange capacity is low. In soils of low cation exchangeable capacity (CEC), cationic holding capacity will be low. However, because of the higher percent base saturation of the soils of this study area, nutrients are readily available to plants and small doses of fertilizer could produce significant crop responses (Table 1).

Analysis of FoliarBlend Fertilizers

The FoliarBlend was subjected to laboratory analysis to determine the concentration of some essential relevant nutrients and elements considered toxic to crops, man and environment. The results as stated below show that the chemical properties of the product met the specifications and the minimum acceptable investigational allowance for organic fertilizers.

Analysis of FoliarBlend as validated by the Department of Soil Science, Faculty of Agriculture/Institute for Agricultural Research, Ahmadu Bello University Zaria, Nigeria, February, 2012 is shown in table 2.

Growth Response of Cotton to FoliarBlend micronutrient application at Funtua site in Katsina state

The growth and yield results obtained for the 2013 rain-fed and 2014 irrigated seasons are presented in Figure 1, 2, Tables 2&3

At 40kgNPK/ha + 500mls of FoliarBlend in 20 litres of water, the plant height (76cm), number of branches (21), number of nodes (30) and number of leaves (36) per plant performed better than the control. This was followed by treatment of 20kgNPK/ha + FoliarBlend and FoliarBlend only respectively (Fig1). During the irrigated season, all treated plants performed better than the control in terms of plant height, number of branches, number of leaves, number of nodes and the highest value was obtained in 40kg NPKha⁻¹ and FoliarBlend (Fig2). This was strongly followed by application by the 20kg NPKha⁻¹ and FoliarBlend and FoliarBlend only respectively. The least value was obtained in Control (Fig2). This result is in line with the findings of Makhdum et al, (2000).

The application of 60kgNPKha⁻¹ + FoliarBlend gave the highest number of bolls (20/plant), and total yield (5.2 t/ha) at rain-fed season when compared to the other treatment levels, while the lowest value was obtained at control plots in the parameters mentioned above (Table). This was followed by the application of 40kgNPKha⁻¹ + FoliarBlend and 20kgNPKha⁻¹ +FoliarBlend respectively. The same trend was obtained during the irrigated season (Table 3)

CONCLUSIONS

The growth parameters of variety SAMCORT 1 were enhanced positively by the application of 40KgNPKha⁻¹ + FoliarBlend during the both seasons (rain-fed and irrigated) in Funtua site and the other 2 locations (Bakori and Fasikari) where the study were also carried out. While, application of 60kgNPKha⁻¹ + FoliarBlend to cotton on the field showed that under prevailing climatic and field conditions with good management practices, the yield of cotton can be greatly improved.

160 Conclusively, under the prevailing poor climatic and soil conditions, fertilization at 40Kg
 161 NPKha⁻¹ (15:15:15) + FoliarBlend is adequate for optimum growth, while 60kgNPKha⁻¹ +
 162 FoliarBlend gave the optimum yield of cotton (*Gossypium sp.*)

163 REFERENCES:

164 Agbede, O. O.(2009). *Understanding Soil and Plant Nutrition*: Salmon Press and coy Keffi,
 165 Nigeria. Pp147-156

166
 167 Chude, V.O. (1998) Understanding Nigerian soils and their fertility management for
 168 sustainable agriculture, Paper delivered at inaugural Lecture atAhmadu Bello University Zaria

169
 170 Halevy, J. (1979) “Fertilizer requirements for high cotton yields”, *Proc. 14th colloq.*,”Soils in
 171 Mediterranean type climates and their yield potential”, International Potash Institute (IPI), Berne,
 172 Switzerland, pp. 359-365.

173
 174 Idem, N.U. A.,(1999) Cotton Production in Nigeria: Baraka press and publishers Ltd, Kaduna.
 175 Pp45-54

176
 177 Malik, M.N.A., Chaudhry, F.I. and Makhdum, M.I. (1996) “Investigation on phosphorus
 178 availability and seed cotton yield in silt loam soils”, *J. An. Plant Sci.*, 6(12), 21-23.

179
 180 Makhdum, M.I., Malik, M.N.A., Chaudhry, F.I. and Shabab-ud-din (2000) “Response of cotton
 181 to balanced fertilization in soils of variable texture”, *The Pakistan Cottons*, 44, 49-55.

182
 183 Muhammad I. M., Muhammad N. A., Malik, S., and Fazal I. C., (2001) Effects of phosphorus
 184 fertilizer on growth, yield and fibre quality of two cotton cultivars Pakistan J. of Research
 185 (science) 12:140-146

186
 187 Ogunlela, V.B.G., and Abed, S.M (1982) Growth Response, Yield and Components of Upland
 188 Cotton (*Gossypium hirsutum L*) as affected by rates and time of N fertilization in the Nigerian
 189 Savanna. *J. of Fertilizer Research*, 3:399-409

190
 191 Prentice, A. N (1992) Cotton with special Reference to Africa. 1st Edition Longman,
 192 Lagos. Pp23-28

193 **Table 1: Physical and Chemical Properties of soils of Funtua site, Katsina State.**

PROPERTIES	VALUES
Sand (g/kg)	580
Silt (g/kg)	220
Clay (g/kg)	150
pH(H2O)	6.0
pH(Kcl)	5.3

194	Organic carbon (g/kg)	4.2
	Nitrogen (g/kg)	0.053
195	Avail. Phosphorus (mg/kg)	0.90
196	<i>Exchangeable Bases(meq /100g)</i>	
197	Ca	3.2
	Mg	1.1
198	K	0.12
199	Na	0.42
	Al	0.3
200	CEC	5.3
201	BS (%)	69.62
202	<i>Extractable Micronutrients (mgkg⁻¹)</i>	
203	Zn	1.37
	Mn	9.73
204	Fe	5.40
205	B	0.27
206		

207 **Table 2: Analysis of FoliarBlend Product**

208	Boron (B).....	0.002%
209	Cobalt (Co).....	0.002%
210	Iron (Fe).....	0.10%
211	Manganese.....	0.10%
212	Molybdenum (Mo).....	0.002%
213	Soluble Zinc (zn).....	0.05%

214 Derived from: Boric Acid, Cobalt MEA, Ferrous Sulphate, Manganese Sulphate,
 215 Ammonium Molybdate, Zinc Sulphate. The above laboratory data clearly confirms that
 216 FoliarBlend is rich sources of plant essential micronutrients for plant growth.

217




Fig 1: The growth response of the cotton plant to FoliarBlend micronutrient fertilizer in Funtua site, Katsina state under 2013 rain-fed season




Fig 2: The growth response of the cotton plant to FoliarBlend micronutrient fertilizer in Funtua site, Katsina state under 2014 irrigated season

227 **Table 3: Effect of various treatment combinations of FoliarBlend on the yield of cotton**
 228 **(t/ha) at funtua under rain-fed 2013**

TREATMENT	Yield t/ha	Number of bolls
FoliarBlend Only	2.50ba	16
20kgNPK/ha + FoliarBlend	2.50ba	17
40kgNPK/ha + FoliarBlend	4.50c	18
60kgNPK/ha + FoliarBlend	5.20d	20
Control	2.13a	13

229
 230 **TABLE 4: effect of various treatment combinations of FoliarBlend on the number of bolls**
 231 **and yield of cotton (t/ha) at funtua under irrigation 2014**

TREATMENT	Yield t/ha	Number of Bolls
FoliarBlend Only	2.90ba	17ba
20kgNPK/ha + FoliarBlend	3.20b	18b
40kgNPK/ha + FoliarBlend	5.50c	21c
60kgNPK/ha + FoliarBlend	6.10d	24d
Control	2.75a	15a

232
 233