UNDER PEER REVIEW

EGG QUALITY CHARACTERISTICS OF JAPANESE QUAILS *(Coturnix coturnix japonica)* FED VARYING LEVELS OF FERMENTED TARO COCOYAM *(Colocasia esculenta var esculenta)*MEAL.

Abstract.

A study was conducted to investigate the effect of FTCM on the egg quality parameters of Japanese quails .Two hundred and twenty five Japanese quails were randomly allotted to five dietary treatments (1-V) of 36 hens and 9 cockerels each. Each treatment was replicated thrice with 12 hens and three cockerels per replicate. In each of the five diets, FTCM was used to replace maize at 0%, 25%, 50%, 75% and 100% for treatments i, ii, iii, iv and v respectively. Quails were fed experimental diets over a period of 70 days. Feed intake was measured daily and quails were weighed weekly .Eggs were collected on weekly basis for egg quality analysis. Results showed that, mean weekly egg weight, shell weight, yolk weight, albumen weight, yolk index and egg breaking strength of quails fed treatments i and ii were significantly($P<0.05$) higher than those fed other treatments. Quails fed diets i, ii and iii recorded highest values of shell thickness. However, there were no significant ($P>0.05$) differences in relative yolk and relative albumen weights across treatments. The results suggest that FTCM could replace maize up to 100% without adverse effect, but for premium egg quality, replacement should not exceed 25%.

KEY WORDS: Quails, FTCM, Maize and Egg quality

Introduction

Poultry production offers the greatest potential for bridging the gap of protein deficiency existing in Nigeria. The poor performance of non- ruminants (poultry, rabbits and pigs) is due to high cost of conventional ingredients such as maize, sorghum , millet, soybeans , groundnut etc. These conventional feedstuffs are expensive and are gravely faced by the problems of seasonality. This all year round unavailability has led to stiff competition between man and livestock industries as demand outstrips supplies. The need to search for alternative feed stuffs that are cheap and readily available cannot be over emphasized. A lot of Researchers have been able to use supplemental levels of non- conventional feed stuffs to reduce cost of production such as: Cocoyam [1 ; 2] , Cassava [3] , Rubber seed [4] , Mango kernel [5; 6] , Sweet orange peels [7] and *Prosopis* [8].Taro cocoyam is one of the most widely spread root crop grown in humid tropics. About 60% of the world production is grown in Africa and 40% in Asia [9]. Fresh taro root is high in energy and contains about 25% starch and 1% protein .Taro has highest contents of P, Mg and Zn of all root/tuber crops and rich in all vitamins except for nicotinic acid [10]. Protein of taro cocoyam is well supplied with the essential amino acid but somehow low in histidine and lysine [11]. Cocoyam is cheaper and yields better than maize [12] and as such stands a better chance of replacing maize. However, the presence of anti- nutrients constitute a problem. Appropriate processing technique such as boiling , fermentation , toasting, sun drying

UNDER PEER REVIEW

etc. could take care of its limitation [13] .This study was designed to investigate the effect of fermented taro cocoyam on egg quality parameters of Japanese quails.

Materials and Methods

The study was carried out in Cross River University of Technology Teaching and Research Farms Calabar, Calabar- Nigeria.

Experimental diets

Unpeeled taro cocoyam corms were bought from Bendeghe village, Etung, Cross River State, Nigeria. Corms were peeled and chopped into sizeable chips of about 14g each .These chips were submerged in a big plastic pot-like container with a tight lid and fermented in water at room temperature (28^0c to 29^0c) for 48hrs. Sun- drying lasted for a week (7days) and this reduced the moisture content to less than 10% for longer keeping. The 48hrs fermented taro cocoyam corms (FTCC) and other ingredients were milled separately and used to formulate experimental diets. The diets had varying protein (%) and energy (ME/Kcal/kg) levels of 21.20, 20.89 , 20.78 , 20.68 , 20.59 , and 2690.30 , 2679.58 , 2593.20 , 2562.20 and2552.90 for protein and energy respectively. These levels were for treatments 1, 2, 3, 4 and5 respectively. The diets were analyzed according to the methods of [14]. Energy was calculated according to the methods of [15]. The percentage composition of the experimental diets is presented in Table one.

Table 1: Percentage composition of diet with fermented taro cocoyam (*Colocasia esculenta var esculenta*) meal

for laying Japanese quails (*Coturnix coturnix japonica)*.

Levels of inclusion

Ingredient	0%	25%	50%	75%	100%
Maize	46.00	34.50	23.00	11.50	0.00
Cocoyam	0.00	11.50	23.00	34.50	46.00
Soybean	20.80	21.00	19.70	19.90	20.00
Fish meal	4.20	4.60	5.80	6.00	6.00
P.K.C	10.90	10.00	10.00	10.00	10.00
Wheat offal	10.00	10.00	9.75	9.10	8.30
Bone meal	7.00	7.00	7.00	7.00	7.00
Salt					
Palm oil					
Vita/ min premix					

UNDER PEER REVIEW

	0.50	0.50	0.50	0.50	0.50
	0.50	0.50	0.50	0.50	0.50
	0.10	0.40	7.50	1.00	1.20
Total	100.00	100.00	100.00	100.00	100.00
Analyzed nutrients: Crude protein (%)	21.20	20.89	20.78	20.63	20.59
ME/Kcals/Kg	2690.30	2679.58	2593.20	2562.20	2552.90

Experimental birds, design and management

The birds (quails) used for the experiment were a total of twelve (12) hens and three (3) cockerels per replicate. In summary, a total of 225 quails (180 hens and 45 cockerels) were studied over a period of ten weeks (70 days). Birds were randomly allotted to five dietary treatments 1 , 2 ,3 , 4 and 5 formulated with 48hrs fermented taro cocoyam meal (FTCM) at 0% , 25% , 50% ,75% and 100% replacement levels respectively (Table 1). Quails were housed in three tier cages of dimension 0.75m2 x0. 38m^2.The cages were made up of wood and wire mesh. Same quantity of feed was weighed and served to the quails on weekly basis, the feed was increased by 50g weekly. Fresh, clean water was supplied *ad libitum* daily. Adequate bio-security and bio- sanitary measures were strictly adhered to. Records of feed intake were kept daily, initial body weight was taken on commencement and subsequently on weekly basis. Total number of eggs laid per replicate group were collected twice daily (8am and 4pm) and recorded. At the end of every week fresh eggs were collected for egg quality analysis.

Determination of egg quality parameters

Nine fresh eggs (large, medium and small) were separated from each treatment (3 eggs from each replicate) and used at weekly interval to determine egg quality parameters throughout the experimental period. The eggs were weighed using a sensitive electronic weighing balance (JJ560 USA) citizen scale to the nearest 0.01g. Each egg was weighed and broken around the equator with care being taken to keep the yolk intact. Yolk was separated from the albumen using an egg separator and weighed separately. Relative yolk and relative albumen weights were

UNDER PEER REVIEW

calculated in percentage by relating the yolk or albumen weights measured to the nearest gram to the whole weight of the particular egg and multiply by 100.

Relative yolk weight =Yolk/Egg wt. x 100

Relative albumen weight=Albumen/Egg wt. x 100

Egg shells were sun- dried for three days and their weights taken with the aid of a sensitive electronic balance. Shell thickness was measured using micrometer screw gauge. The measurement were taken from the pointed, middle and broad part of the egg and the mean thus obtained and measured to the nearest 0.01mm. Yolk height was taken with the aid of a Spherometer at the heighest point of its surface and yolk width was measured at long and short axis using vernier caliper. The values obtained were used to calculate yolk index. Egg breaking strength (EBS) was calculated using the formular suggested by [16] and represented as;

Yolk Index = mean height/mean width

EBS =50.86 x (EW) 0.915, where EW =Egg weight.

Data obtained were subjected to analysis of variance using the completely randomized design (CRD) as described by [17]. Least significant difference method was used to separate means that differed significantly [17].

Results and Discussion

The egg weights of quails fed control diets and 25% inclusion levels of cocoyam had significantly ($p<0.05$) better egg weights compared to quails on other dietary treatments. [18] indicated that egg mass or weight can be used as a criteria in assessment of nutritional status. Similar results were reported in which egg weights increased significantly ($p<0.05$) at higher levels of dietary protein inclusion [19 and 20].The weight ranged between 9.40 to10.80g.This range was higher than the 8.92 to 10.13g reported by [20] and the 7.89 to 9.30g reported by [21]. [22] indicated that a good quail egg should weigh about 9.30g and above. Quails fed various diets met this requirement. The egg shell weights of quails placed on treatments i and ii were significantly ($p<0.05$) heavier than those of other treatments. It was observed that beyond 25% inclusion levels , the egg shell weights decreased with increased supplementation. This could be

UNDER PEER REVIEW

attributed to decrease in dietary protein across the treatments. The egg shell weights ranged from 1.12 to 1.30g. This range was higher than 0.90-0.93g reported by [19] and the 0.91 to 0.95 reported by [23]. It could be as a result of the varying Ca:P ratio in the experimental diets of the various experiments. Result showed that shell thickness of quails fed control diets and 25% inclusion levels of cocoyam had significantly ($p<0.05$) thicker shells than quails fed other diets. Shell thickness ranged from 0.203 to 0.217mm. The result was lower than that reported by [19](0.24-0.26mm). The result contradicts with the report of [24] who asserted that, shell thickness declines with age due to increase in egg size which forces the constant amount of shell to spread thinner and thus reducing the thickness and quality. But was in agreement with the report of [25] who asserted that, the amount of shell materials deposited per egg does not change with age and size. Otherwise, treatments V and IV should have had significantly ($P<0.05$) thicker shells according to [24] . Egg breaking strength followed similar trend with quails placed on diets I and II having significantly ($P<0.05$) higher values compared with others. The result ranged from 442.1 to 502.77. This implies that eggs of quails placed on treatments I and II will not easily get cracked or broken when compared with eggs of other treatments. Generally, egg breaking strength (EBS) of quails is lower than that of chickens which could be as high as 2322±55.60 [26 ; 27]. By implication, the higher the EBS the better the egg quality. Yolk weight and albumen weights of quails fed control and 25% inclusion levels of cocoyam had significantly ($P<0.05$) higher weights than those of other treatments. Yolk and albumen weights ranged from 3.28 to 3.60g and 5.00 to 5.91g respectively. The result of the yolk weight was in agreement with the report of [25] who noted that, albumen weight (AW) is negatively correlated with the yolk weight (YW).This infers that, eggs with higher values of AW will have lower yolk weights. The values recorded for AW in this finding were different from those recorded by [23] (4.42 to 4.60g).The result was in agreement with the reports of [28] who indicated that AW was highly correlated with egg weight. It was observed that treatments I and II with highest values of egg weights (EW) had highest values of albumen weights (AW). The AW is sufficient to measure egg quality [29]. There were no significant ($P>0.05$) differences recorded in relative yolk weights (RYW) and relative albumen weights (RAW) across the treatments. RYW and RAW ranged from 33.33 to 34.89 and52.75 to 54.72 respectively. The results of RYW and RAW are in agreement with the observation of [28] who observed that, as the egg size (ES) increases, the RYW decreases and the RAW increases. This result also agrees with the findings of [25] who

UNDER PEER REVIEW

asserted that although the age of a layer has positive relationship with ES, it exerts a negative influence on the yolk properties. Result showed significant (P<0.05) difference in yolk index (YI) with treatment I and II having highest values. The YI ranged from 0.468 to 0.489. This range varied from the 0.35 reported by [19]. YI is the measure of the standing up quality of the yolk. Average values for fresh eggs usually fall between 0.40 to 0.42 [29]. The authors further stated that, the YI of 0.25 or lower indicates the weakness of the yolk and extreme difficulties in handling it for measurement without breaking. In this findings, one could deduce that the YI of all the treatments were within the range and could be handled for measurement without scattering easily.

Table 2: Summarized table of egg quality characteristics of laying quails fed vary levels of fermented taro cocoyam meal.

Treatments levels(%)

Parameter	0%	25%	50%	75%	100%
External characteristics					
Egg weight (g)	8.636 ± 0.41^{a}	8.633±0.40 ^{a}	8.128±0.32^{b}	7.757±0.41^{c}	7.516±0.52c
Egg shell weight (g)	1.29±0.08^{a}	1.30±0.08^{a}	1.19±0.09 ^{b}	1.15±0.08^{c}	1.12±0.10 ^{d}
Egg breaking strength	502.36±19.11^{a}	502.77±18.58^{a}	472.68±14.87^{b}	451.23±19.00^{c}	442.21±23.77 ^{c}
Shell thickness(mm)	0.216±0.004^{a}	0.216±0.005 ^{a}	0.217±0.005^{a}	0.203±0.006^{b}	0.205±0.006^{b}
Internal characteristics					
Yolk weight (g)	3.60±0.10^{a}	3.45±0.07^{b}	3.34±0.07^{c}	3.28±0.10^{c}	3.28± 0.10^{c}
Albumen weight (g)	5.90±0.25^{a}	5.91±0.24^{a}	5.56±0.16^{b}	5.21±0.62^{c}	5.00±0.35^{c}
Relative yolk weight (%)	33.47±0.43	33.40±0.41	34.05±0.49	34.73±1.00	35.32±1.29

UNDER PEER REVIEW

Relative albumen weight (%)	54.61±0.39	54.65±0.35	54.73±0.34	53.52±0.84	52.75±1.19
Yolk index	0.488±0.06[a]	0.489±0.06[a]	0.475±0.005[b]	0.472±0.002[b]	0.468±0.002[b]

Different superscripts (a, b, c and d) within the row indicates significant ($p<0.05$) differences.

Conclusion

The results of this study suggests that, fermented taro cocoyam meal could replace maize up to 100% without compromising egg quality but for premium egg quality, replacement should not exceed 25%.

References

Agwunobi LN, Agwukan PO, Cora OO, Isika MA. Studies on the use of *Colocasia esculenta*(taro cocoyam) in the diets of weaned pigs. *Tropical Animal health and production*. 2002; 34(3):243-247.

Okon BI, Obi MB, Ayuk AA. Performance of quails (*Coturnix coturnix japonica*) fed gradual levels of boiled sun-dried tarococoyam(*Colocasia esculenta var esculenta*) as a replacement of maize. *Agricultural Journal*. 2007; 2(6):654-657.

Ogbonna JU. Nutritive value of cassava peels meal for poultry. *A Review of the proceeding of the 2nd annual conference of Animal Science Association of Nigeria. 1997; Lagos, sept*.

UNDER PEER REVIEW

Agwunobi LN, Ina-ibar OB. Processed rubber seed meal in the diets of growing Japanese quail (*Coturnix coturnix japonica*).*Proceedings of the 32nd Annual conference of the Nigerian society for Animal production (NSAN). 2007;Calabar, march,Pp.386-388.*

Abang FB, Ejugwu G, Aker DT. Nutrient profile of Japanese quail(*Coturnix coturnix japonica*) breast muscles and egg (whole) fed sun-dried mango(*Mangifera indica*) kernel meal. I.O.S.R *Journal of Agriculture and Veterinary Science* (IOSR-JAVS). 2015; 3(8):37-40.

Diarra SS, Usman BA, Igwebuike JU. Replacement value of boiled mango kernel in broiler finisher diets. *ARPRO Journal of Agriculture and Biological Science. 2008; 5:47-52.*

Oluremi OIA, Ojighen VO, Ejembi EH. The nutritive potentials of sweet orange (*citrus sinensis*) rind in broiler production. *International Journal of Poultry Science. 2006;*5(7):613-617.

Yusuf ND, Ogali DM,Hassan DI, Musa MM, Doma UD. Effect of decorticated fermented *Prosopis* seed meal(*Prosopis africana*) on growth performance of broiler chickens. *International Journal of Poultry Science*. 2008; 7(11):105-1057.ISSN.

Viotti V. Taro production in hawaiian. Island. *Honolulu advertiser. 2004;* March.

Purseglove JW. *Tropical Crops Monocotyledons*.John willey and sons. 1992; New York.

Ohtsuka R, Kawabe T, Inaoka T, Suzuki T, Hong F, Akinchi OT, Sugaharu T. Composition of local and purchased food consumed by the Gidra in lowland papua. *Ecology of Feed and Nutrition*. 1984; 15:159-169.

Food and Agricultural Organization(FAO). Food and Agricultural Organization production year book. Food and Agricultural Organization(FAO) of the United Nations. 2004; Rome, Italy.

Obi MB. Improving the nutritional value of taro cocoyam(*Colocasia esculenta var.esculenta*) through processing for the feeding of growing and laying quails. Ph.D. thesis submitted to Department of Animal Science University of Calabar. 2010; Pp210.

Association of Official Analytical Chemist . *Official methods of analysis* (17th edition). 2000; Washington D.C.

Carpenter KJ, Clegg KM. The metabolisable energy of poultry feeding stuffs in relation to their chemical composition. *Journal of science and Agriculture. 1956; 7:45-51.*

Arad z, Marder J. Difference in egg shell quality among the Sinai bedouing fowls, the commercial white leghorn and their cross breeds. *British Poultry Science*. 1982; 23:107-112.

Steel RGD, Torrie JH. *Principles and proceedings of statistics,Mc Graw-hill book.1980; Co.Inc.Newyork*.

UNDER PEER REVIEW

Lin, H., Buyse, J., Sheng,Q.K., Xie,Y.M. and Song,J.L.(2004). Effects of ascorbic acid supplementation on the immune function and laying performance of heat-stressed laying hens. *Journal of Agriculture and Environmental*. 1:103-107.

Bawa,G.S. Lombin,H., Karson,P.,Musa, U., Pay,E. and Shamak,D. (2011). Response of Japanese Breeder quails to varying Dietary Protein Levels in the Tropics. *Nigerian Journal of Animal Production. 38:43-54*.

Babangida, S.and Ubosi,C.O.(2006). Effects of varying dietary protein levels on the performance of laying Japanese quails (*Cortunix coturnix japonica*) in a semi-arid environment. *Nigeria Journal of Animal Production*, 33(1):45-52.

Jika, C.P.(2015). Egg quality parameters of quails fed graded levels of *Prosopsis africana* seed coat meal as a replacement for maize. B. Agric. Project submitted to the Department of Animal Production, University of Agriculture, Makurdi.Pp5.

Yamane etal

Sachdev, A.K., Ahya, S.D and Cropals,R. (1975). The ability of hens to maintain calcium deposition in the egg shell and egg yolk as the hen ages. *Poultry Science*.54:1729-1723-1729.

Roland (Sr), D.A., Putnem, C.E. and Hilburn,R.L.(1978). The influence of age on ability to hen to maintain egg shell calcification when stressed with inadequate dietary calcium. *Poultry Science*.57:1616-1621.

Essien, A I, Ekanem EO. Effect of the age of layers, rate of egg produce and time of oviposition on some initial egg shell quality indices in a humid tropical environment. *Beitrage zur Tropischen land wirl*. *Shaft veterinary medicine*.1990; 28:125-131.

Oluwofeso, O and Omisami,OO. (2008). Utilization of cassava peels treated with yeast culture by egg laying chickens. *Proceedings of the 33rd annual conference of the Nigeria society for Animal production. Ogun,2008. Pp492-495*.

Woolford, R. (1985). Nutrition and egg quality. *Poultry International* (May edition). 46 – 48.

Fletcher OI, Brittan WM, Rahn AO, Savage SL. The influence of layer flock age on egg component yield and solid content. *Poultry Science*. 1980; 60(80):983-987.

Card, I E and Neshiem, MC(1972). *Poultry Production* 11th edition. Philadelphialea and Febiger.

Abang,F.B, Ejugwu, G. and Aker, D.T.(2015). Nutrient profile of Japanese quail(*Coturnix coturnix japonica*) breast muscles and egg (whole) fed sun-dried mango(*Mangifera indica*) kernel meal. I.O.S.R *Journal of Agriculture and Veterinary Science*(IOSR-JAVS). 3(8):37-40

UNDER PEER REVIEW

Agwunobi, L.N., Agwukan, P.O.,Cora,O.O. and Isika M.A.(2002). Studies on the use of *Colocasia esculenta*(taro cocoyam) in the diets of weaned pigs. *Tropical Animal health and production*. 34(3):243-247.

Agwunobi,L.N. and Ina-ibar, O.B.(2007). Processed rubber seed meal in the diets of growing Japanese quail (*Coturnix coturnix japonica*).*Proceedings of the 32nd Annual conference of the Nigerian society for Animal production (NSAN).Calabar, march,2007.386-388.*

Arad,z. and marder,J. (1982). Difference in egg shell quality among the Sinai bedouing fowls, the commercial white leghorn and their cross breeds. *British Poultry Science*.23:107-112.

Association of Official Analytical Chemist (2000). *Official methods of analysis* (17th edition). Washington D.C.

Babangida, S.and Ubosi,C.O.(2006). Effects of varying dietary protein levels on the performance of laying Japanese quails (*Cortunix coturnix japonica*) in a semi-arid environment.*Nigeria Journal of Animal Production*, 33(1):45-52.

Bawa,G.S. Lombin,H., Karson,P.,Musa, U., Pay,E. and Shamak,D. Response of Japanese Breeder quails to varying Dietary Protein Levels in the Tropics. *Nigerian Journal of Animal Production.* 2011. *38:43-54*.

Card, I.E and Neshiem,M.C.(1972). *Poultry Production* 11th edition.Philadelphia lea and Febiger.

Carpenter, K.J and Clegg, K.M.(1956). The metabolisable energy of poultry feeding stuffs in relation to their chemical composition. *Journal of science and Agriculture.7:45-51.*

Diarra,S.S, Usman, B.A. and Igwebuike,J.U. (2008). Replacement value of boiled mango kernel in broiler finisher diets. *ARPRO Journal of Agriculture and Biological Science. 5:47-52.*

Essien AI, Ekanem EO. Effect of the age of layers, rate of egg produce and time of oviposition on some initial egg shell quality indices in a humid tropical environment. *Beitrage zur Tropischen land wirl*. *Shaft veterinary medicine*.1990; 28:125-131.

Fletcher OI, Brittan WM, Rahn AO, Savage SL. The influence of layer flock age on egg component yield and solid content. *Poultry Science*. 1980; 60(80):983-987.

Food and Agricultural Organization(FAO),(2004). Food and Agricultural Organization production year book. Food and Agricultural Organization(FAO) of the United Nations. Rome, Italy.

***Jika, C.P.(2015). Egg quality parameters of quails fed graded levels of *Prosopsis africana* seed coat meal as a replacement for maize. B. agric, project submitted to the Department of Animal Production, University of Agriculture, Makurdi.Pp53.

UNDER PEER REVIEW

Lin, H., Buyse, J., Sheng,Q.K., Xie,Y.M. and Song,J.L.(2004). Effects of ascorbic acid supplementation on the immune function and laying performance of heat-stressed laying hens. *Journal of Agriculture and Environmental*. 1:103-107.

Obi,M.B.(2010). Improving the nutritional value of tarococoyam(*Colocasia esculenta var.esculenta*) through processing for the feeding of growing and laying quails. P.hD. thesis submitted to Department of Animal Science University of Calabar, Calabar.Pp210

Ogbonna,J.U. (1997). Nutritive value of cassava peels meal for poultry. *A Review of the proceeding of the 2nd annual conference of Animal Science Association of Nigeria. Lagos, sept 1999*.

Ohtsuka,R., Kawabe,T., Inaoka,T., Suzuki, F., Hong.O.T., Akinchi and Sugaharu,T.(1984). Composition of local and purchased food consumed by the Gidra in lowland papua. *Ecology of Feed and Nutrition*. 15:159-169.

Okon,B.I, Obi,M.B. and Ayuk,A.A.(2007). Performance of quails(*Coturnix coturnix japonica*) fed gradual levels of boiled sun-dried tarococoyam(*Colocasia esculenta var esculenta*) as a replacement of maize. *Agricultural Journal*.2(6):654-657.

Oluremi,O.I.A., Ojighen,V.O. and Ejembi, E.H. (2006). The nutritive potentials of sweet orange (*citrus sinensis*) rind in broiler production. *International Journal of Poultry Science* 5(7):613-617.

Oluwofeso, O. and Omisami,O.O. (2008). Utilization of cassava peels treated with yeast culture by egg laying chickens. *Proceedings of the 33rd annual conference of the Nigeria society for Animal production. Ogun,2008,Pp492-495*.

Oluyemi J.A. and Robert, F.A. (2000) Poultry production in the warm climate. Macmillian publisher. L.t d.London.

Purseglove, J.W.(1992). *Tropical Crops Monocotyledons*.John willey and sons. New York.

Roland (Sr), D.A., Putnem, C.E. and Hilburn,R.L.(1978). The influence of age on ability to hen to maintain egg shell calcification when stressed with inadequate dietary calcium. *Poultry Science*.57:1616-1621.

Sachdev, A.K., Ahya, S.D and Cropals,R. (1975). The ability of hens to maintain calcium deposition in the egg shell and egg yolk as the hen ages. *Poultry Science*.54:1729-1723-1729.

Steel,R.G.D. and Torrie,J.H.(1980). *Principles and proceedings of statistics,Mc Graw-hill book Co.Inc.Newyork*.

Viotti,V.(2004). Taro production in hawaiian. Island. *Honolulu advertiser*,march,6,2004.

UNDER PEER REVIEW

Woolford, R.(1985). Nutrition and egg quality. *Poultry International* (May edition). 46-48.

Yusuf, N.D., Ogali, D.M.,Hassan,D.I., Musa,M.M., and Doma, U.D.(2008). Effect of decorticated fermented *Prosopis* seed meal(*Prosopis africana*) on growth performance of broiler chickens. *International Journal of Poultry Science*.7(11):105-1057.ISSN.