

Original Research Article

EVALUATION AND AGRONOMIC POTENTIAL OF ANDROGENIC PEPPER GENOTYPES DERIVED FROM PIRAN (*Capsicum annuum* L. cv. Piran)

ABSTRACT

Aims: The aim of this study was to evaluate and to estimate the agronomic potential of two androgenic genotypes via important plant, fruit and production traits compared to the mother genotype Piran.

Study design: Random block design in four replications.

Place and Duration of Study: Strumica, Faculty of Agriculture, Goce Delcev University-Stip, Republic of Macedonia, four-years experiment.

Methodology: Evaluation of the androgenic genotypes was performed according to morphological traits given in Descriptor for *Capsicum* spp. The analysis of plant traits was conducted on 20 randomly chosen plants (5 plants per replication) in maximum growth stage from for each androgenic genotype and the control Piran. The analysis of fruit traits was conducted on 40 randomly chosen fruits (10 fruits per replication) in ripe stage from for each androgenic genotype and the control.

Results: Androgenic genotype P3 was characterized as tall pepper type with medium-thick stem and large leaves. According to the fruit weight (41.45 g) and the fruit index higher than 4, the genotype is classified as long, horn-shaped, middle-size fruit pepper. The fruit flesh of 77.05% and the pericarp thickness 0.26 cm characterize this fruit as medium-fleshy. Androgenic genotype P4 was characterized as very tall pepper type with medium-thick stem and medium-large leaves. The fruit weight was 42.84 g, contains averagely 7.37% dry matter, 76.71% fruit flesh and 0.25 cm pericarp thickness. P4 fruit is long, horn-shaped, medium-size and medium-fleshy. Fruits of both genotypes are suitable for fresh and preserved consumption.

Conclusion: Analysis of morphological and productivity traits of androgenic genotypes provides valuable information for future breeding. Androgenic genotype P3 differ from the initial genotype for four traits, while the P4 genotype for six traits. The results of this research prove the advantage of pepper androgenesis as a tool for enrichment of biodiversity and fast development of valuable pepper genotypes for future breeding.

Keywords: androgenesis, pepper, Piran, androgenic genotype P3, androgenic genotype P4, morphological traits, characterization

1. INTRODUCTION Introduction week need to be improved. Moving back and forth from ideas, REVISED....

Pepper (*Capsicum annuum* L.) is an important crop that can be improved by utilization of the methods of classical genetics, but also by using plant tissue culture and plant biotechnology tools which can shorten breeding process of many crops [1-2].

Androgenesis is one of the methods utilized for creation of haploid and spontaneous dihaploid plants in *in-vitro* culture [3-5]. Gained haploids and dihaploids have certain genetic potentials which are quickly phenotypically expressed because they origin from a haploid cell [6-7]. These androgenic plants, before being included in breeding process, must be studied for their growth and development stages, their genetic and morphological characteristics compared to the mother genotype which give information about similarities and differences between androgenic and mother genotypes [8-10].

The aim of this study was to evaluate and to estimate the agronomic potential of two androgenic genotypes P3 and P4 derived from initial genotype Piran via important morphological and production traits.

2. MATERIAL AND METHODS

The method of androgenesis method was successfully applied for development of fertile androgenic plants *in vitro* conditions from the Macedonian variety Piran. Detailed description of the methodology and the process of creation of androgenic genotypes is presented in different publications [11-14]. In this research plants and fruits of two androgenic genotypes P3 and P4 created from the variety Piran were tested in greenhouse conditions at the Faculty of Agriculture, Goce Delcev University – Strumica, Republic of Macedonia, during four-years experiment. The variety Piran, used as mother genotype in anther culture, was used as control (standard) during evaluation of the androgenic genotypes P3 and P4. It was obtained from the gene bank of Faculty of Agriculture, Goce Delce University – Stip [15]. Piran is Macedonian variety which can be cultivated in open field and under protected conditions. The average plant height is 80 cm, fruits are large with average weight of 100-120 g, yellow in unripe stage and red when ripe. Piran average yield is 60-80 t/ha [14,16].

2.1 Research design and dynamics

During the first experimental year seeds obtained from fertile androgenic plants were utilized as starting material in the experiment. Seeds from selected typical fruit from each androgenic genotype of the previous experimental year were used in the subsequent experimental years. Each experimental year the seeds of studied genotypes were sown in the beginning of April in non-heated greenhouse. At the end of May, the seedlings were transplanted into pots (d=22 cm) filled with mixture of soil, sand and peat. The experiment was design in four replications, in random block design. From each androgenic genotype and control genotype Piran, 40 plantlets (10 per replication) were transplanted in plastic pots, which were placed in the greenhouse with 30cm/100cm distance scheme. Plants were cultivated according to standard cultivation practices under protected conditions. Irrigation and nutrition were performed as fertigation with drip system. Selection of parameters and evaluation of genotypes was performed according to recommendations given in Descriptor for Capsicum sp. [17]. The analysis of plant traits was conducted on 20 randomly chosen plants (5 plants per replication) in maximum growth stage from for each androgenic genotype and the control Piran. During the research the following plant traits were analyzed:

- plant height (cm)
- plant height to the first branching (cm)
- stem thickness (cm)
- internode length (cm)

The analysis of fruit traits was conducted on 40 randomly chosen fruits (10 per replication) in ripe stage from for each androgenic genotype and the control. During the research the following fruit traits were analyzed:

- fruit length (cm)
- fruit width (cm)
- fruit index (fruit shape)
- fruit weight (g)
- number of fruit locules
- fruit flesh percentage (%)
- pericarp thickness (cm)
- dry matter in fresh fruit (%)

The analysis of productivity traits was conducted on whole experiment level. During the research the following productivity traits were analyzed:

- number of fruits per plant
- yield per plant (g)
- total yield (kg/m²)

A classification of plant, leaf, fruits and productivity of androgenic genotypes was made based on the results of studied morphological and production traits according to Jankuloski (1983, 1997). It is the most suitable and acceptable classification of peppers by morphological traits which are breed and cultivated in the Western Balkans [18-19].

2.2 Biometric analysis of characteristics

Statistical analysis of variance is applied for the evaluation of the experiment in general and each of the characteristics among all tested genotypes (One-Way ANOVA test).

For evaluation of the difference between examined genotypes Duncan's multiple range test is utilized for each trait in each experimental year on 0.05% and for the four-year average on level 0.05% and 0.01%.

The statistical analyzation of the results was performed by IBM SPSP Statistics Software 19.0.

3. RESULTS AND DISCUSSION

3.1 Morphological traits of androgenic plants

Morphological traits of pepper plants as total plant height and plant height to the first branching, leaf shape, stem thickness, fruit size and shape are traits which give basic marks of a pepper variety. However, they are dependent on many external factors as temperature, air humidity and cultivation practices [1, 14]. The mean values of morphological traits of plants in maximum growth stage during the four-years research of androgenic lines P3, P4 and the mother genotype Piran are presented in Table 1.

The average plant height was 87.61 cm (Piran), 88.55 cm (P3 genotype) and 93.68 cm (P4 genotype). In I and II research year, the plant height shows significant difference between P4 genotype and the control, while in IV year, the plant height of P3 and P4 is significantly different as compared to the control. The average height to the first branching of plant is 39.48 cm (P3) and 39.48 (P4) and it is significantly different as compared to the control genotype (34.50 cm). For the first three experimental years, there is a significant difference between the mean values for P3 and P4 genotypes and the control Piran. Other researchers also reported differences in plant height between androgenic lines and initial variety [20-22].

Stem thickness of tested genotypes vary from 0.65 cm (P3, I year) to 1.14 cm (P4, I year), where the lowest average value was measured in IV experimental year and the highest in III experimental year. Averagely for the whole experimental period, the highest average value for stem thickness is registered for P4 genotype (1.01 cm) and it is significantly different from the control Piran (0.89 cm).

The mean internode length of studied genotypes has not shown statistically significant difference. Longer internodes were measured during the first two experimental years as compared to internode length in the third and the forth experimental year. Many plants, including pepper, have internodes length which is determined by day-night temperature variations, which is mainly due to increased cell elongation than due to increasing of cell number. Peppers have shorter internodes when difference between day and night temperatures is smaller [23].

Table 1. Morphological characteristics of plants at maximum growth stage of P3 and P4 androgenic genotypes and the control Piran

Experimental year	Genotype	Plant height (cm)	Height to the first branching (cm)	Stem thickness (cm)	Internode length (cm)
—	Piran	85.10b	33.80b	0.88b	9.90a
	P3	90.15b	40.73a	0.96ab	9.42a
	P4	98.15a	41.15a	1.14a	9.18a
	Piran	93.60b	37.20b	0.90b	10.47a
=	P3	99.30b	44.90a	0.95ab	9.50a
	P4	108.20a	45.20a	1.00a	10.10a
	Piran	87.80a	28.10b	0.96a	4.45a
≡	P3	87.50a	32.45a	1.01a	4.78a
	P4	88.50a	34.30a	0.99a	4.92a
≥	Piran	79.83a	41.83a	0.76a	5.81a
	P3	63.44b	33.67b	0.65a	5.51a
	P4	63.00b	34.56b	0.75a	4.67a
Average I-IV	Piran	87.61 ^{a,1}	34.50 ^{b,1}	0.89 ^{b,1}	7.86 ^{a,1}
Average I-IV	P3	88.55 ^{a,1}	38.62 ^{a,2}	0.94 ^{ab,12}	7.59 ^{a,1}
Average I-IV	P4	93.68 ^{a,1}	39.48 ^{a,2}	1.01 ^{a,2}	7.63 ^{a,1}

Mean separation in columns by Duncan's multiple range test. In each column, values followed by the same letter do not differ significantly at $P=0.05$ and values followed by the same number do not differ significantly at $P<0.01$

Figure 1 presents mean values for total plant height and plant height to the first branching during whole experimental in flowering stage and maximum growth stage of plants. The analysis of plant growth of the whole plant shows that the P3 plants, averagely have grown in height for 19.13 cm, P4 plant 21.91 cm, while the control plants 28.31 cm. This finding is an indicator that the mother variety Piran is characterized by the greatest potential of growth and development as compared to the androgenic genotypes. It can be considered as positive, if the aim of breeders is to gain taller plants, or negative if the aim is creation of shorter plants because of certain cultivation measures. On the other hand, the analysis of the plant height to the first branching showed that the average values of this trait between the two growth stages is the same as 3.25 (control Piran), 3.91 cm (P3 genotype) and 3.81 cm (P4 genotype) beside the difference of the total plant height of the genotypes analyzed. The results of total plant height and plant height to the first branching show that plant growth between the flowering phase and maximum growth phase is happening above the first branching, consequently the plant grows in height with development and growth of new branches. The growth in stem base is very small. This characteristic is very important for cultivation of different pepper varieties under protected conditions because a lush growth will need stem and branches staking in order they to sustain fruit load [14]. This is the reason for development of different systems for pepper plant support with temporary or permanent staking construction [24].




Fig. 1. Comparison of plant height and plant height to first branching in flowering stage and maximum growth stage of androgenic genotypes P3 and P4 and the control Piran

Table 2. Length, width and index of leaf of P3 and P4 androgenic genotypes and the control Piran

Experimental year	Genotype	Leaf length (cm)	Leaf width (cm)	Leaf index
—	Piran	11.86a	6.40a	1.86b
	P3	11.94a	5.89b	2.03a
	P4	11.31a	5.96b	2.03a
	Piran	10.21b	5.33a	1.92b
=	P3	11.09a	5.20ab	2.14a
	P4	10.32b	4.86b	2.14a
	Piran	9.24b	5.04a	1.84b
	P3	10.41a	5.20a	2.01a
≡	P4	10.62a	5.09a	2.10a
	Piran	8.68a	4.45ab	1.96b
	P3	9.43a	4.24b	2.26a
	P4	9.25a	4.67a	1.96ab
IV	Piran	10.76 ^{a,1}	5.76 ^{a,2}	1.88 ^{b,1}
	P3	11.15 ^{a,1}	5.41 ^{b,12}	2.07 ^{a,2}
	P4	10.73 ^{a,1}	5.22 ^{b,1}	2.07 ^{a,2}
	Piran	10.76 ^{a,1}	5.76 ^{a,2}	1.88 ^{b,1}

Mean separation in columns by Duncan's multiple range test. In each column, values followed by the same letter do not differ significantly at $P=0.05$ and values followed by the same number do not differ significantly at $P<0.01$

The average leaf length of androgenic genotypes is 11.15 cm (P3) and 10.73 (P4). The only statistically proven difference in leaf length is in II experimental year, when the leaf length of P3 genotypes (11.09 cm) is longer as compared to P4 genotype (11.09 cm) and the control Piran (10.21 cm). The leaves of P3 (5.41 cm) and P4 (5.22 cm) are significantly narrower than the control leaves (5.76 cm) which is proven during the whole experimental period. The average leaf index for androgenic genotypes P3 and P4 is 2.07, which is significantly higher than the control Piran (1.88). Leaves of P3 genotype are classified as large-size leaves, while P4 as medium-size leaves. Although they have the same leaf index, the first one are longer. Shrestha *et al.* (2011) presented results of leaves dimensions of diploid androgenic plants smaller than the mother genotype which is opposite to our results [25].

3.2 Morphological traits of fruits from androgenic plants

Morphological traits of pepper fruit vary in wide ranges, influenced and caused by many factors. The most authors agree that variation degree of morphological trait is genetically dependent and inherited, but they are always influenced by the complex of external factor, cultivation technology and cultivation practices (under protected conditions or open field conditions). When making pepper characterization it is obvious that the most variations are observed in fruits, an organ which determines taxonomy and variety of the species [1, 6, 14, 20, 26].

The average values for fruit length of androgenic genotypes and the control in the experimental period did not show significant difference for $P=0.05$ and $P<0.01$, which indicates that the control and androgenic genotypes are characterized by stability and good equability of fruit length (Table 3). Fruit length have not shown significant difference during I and II experimental year. In III experimental year the fruit length of the control genotype is significantly different compared to P4 androgenic genotype, while in IV experimental year there is significant difference between the control and P4 genotype. The biggest average fruit length is gained in the control fruits in III experimental year (15.41 cm), the smallest in P3 (12.50 cm) in IV. The fruit length during the experimental period vary from 10.46 cm to 14.11 cm. Koleva Gudeva and Trajkova (2012) refer to Piran fruit length from 12.7 cm to 20.7 cm [27].

Table 3. Length, width, index, weight and number of locules of ripen fruits of androgenic lines P3 and P4 and the control Piran

Experimental year	Genotype	Fruit length (cm)	Fruit width (cm)	Fruit index	Fruit weight (g)	Number of fruit locules
—	Piran	13.54a	3.37a	4.04a	32.17b	2.10a
	P3	14.41a	3.40a	4.28a	43.40a	2.00a
	P4	14.70a	3.50a	4.27a	48.00a	2.05a
=	Piran	13.31a	3.40a	3.96a	35.77b	2.30a
	P3	12.97a	3.51a	3.81a	39.59ab	2.55a
	P4	13.82a	3.63a	3.89a	43.79a	2.20a
≡	Piran	12.88b	3.43b	3.79a	37.13b	2.10a
	P3	13.74b	3.66ab	3.79a	44.30a	2.15a
	P4	15.41a	3.78a	4.09a	50.10a	2.20a
≥	Piran	14.12a	3.09b	4.61a	35.06ab	2.30a
	P3	12.50b	3.37a	4.17ab	37.55a	2.07a
	P4	13.75a	3.18a	3.95b	29.48b	2.35a
Average I-IV	Piran	13.46 ^{a,1}	3.32 ^{b,1}	4.10 ^{a,1}	35.03 ^{b,1}	2.20 ^{a,1}
Average I-IV	P3	13.71 ^{a,1}	3.49 ^{a,1}	3.97 ^{a,1}	41.45 ^{a,2}	2.20 ^{a,1}
Average I-IV	P4	14.11 ^{a,1}	3.52 ^{a,1}	4.05 ^{a,1}	42.84 ^{a,2}	2.20 ^{a,1}

Mean separation in columns by Duncan's multiple range test. In each column, values followed by the same letter do not differ significantly at $P=0.05$ and values followed by the same number do not differ significantly at $P<0.01$

The mean values of fruit width show that androgenic genotypes P3 (3.49 cm) and P4 (3.52 cm) are significantly different compared to the control (3.32 cm) for $P<0.05$. The fruit width of the control peppers shows variation of the average during the experimental years. The biggest width was measured for P3 (3.66 cm) in III experimental year, while the smallest for control (3.09 cm) in IV experimental year. The average of fruit index values varies from 3.79 to 4.61. Only in IV experimental year, the fruit index of androgenic genotype P4 (3.95) is significantly different then the control (4.61).

The statistical analysis of fruit index for four-years experimental period has not shown statistical difference on both statistical levels.

The biggest mean fruit weight is recorded for P4 (42.84 g) compared to P3 (41.45 g) and the control (35.03 g). The fruit weight shows significant variation in the three experimental years, when the fruit weight is always statistically heavier compared to the control fruits.

The average number of fruit locules was from 2.0 to 2.55 without statistically significant difference.

The four-year mean values of traits fruit flesh, pericarp thickness and dry matter in fresh fruit for androgenic lines P3, P4 did not differ significantly from the control Piran (Table 4). The highest mean percentage of fruit flesh is 83.54% (P4; I year) and the lowest for 72.26% (P3; III year). The thickest fruit pericarp is recorded in the last experimental year. The percentage of dry matter in fresh fruit varied during the whole experimental period. During II, III and IV experimental year, the values for this characteristic significantly differ among examined genotypes and the content of dry matter is lower for androgenic fruits as compared to the control fruits. Different androgenic genotypes derived from Piran are tested in a plastic tunnel conditions, where it is proven that statistically significant differences of androgenic fruits compared to the control fruits were observed for fruit length and pericarp thickness [27].

Table 4. Fruit flash, pericarp thickness and dry matter of ripen fruits of androgenic lines P3 and P4 and the control Piran

Experimental year	Genotype	Fruit flesh (%)	Pericarp thickness (cm)	Dry matter in fresh fruit (%)
—	Piran	81.83a	0.21a	8.98a
	P3	83.54a	0.22a	9.37a
	P4	81.73a	0.22a	9.31a
=	Piran	78.48a	0.17b	8.05a
	P3	75.86a	0.27a	7.95a
	P4	76.53a	0.29a	7.08b
≡	Piran	73.22a	0.22a	7.45a
	P3	72.26a	0.24a	7.03ab
	P4	75.05a	0.23a	6.58b
≥	Piran	77.05a	0.35a	7.00a
	P3	76.37a	0.30ab	7.10a
	P4	73.54a	0.27b	6.53b
Average I-IV	Piran	77.65 ^{a,1}	0.23 ^{a,1}	7.87 ^{a,1}
Average I-IV	P3	77.05 ^{a,1}	0.26 ^{a,1}	7.91 ^{a,1}
Average I-IV	P4	76.71 ^{a,1}	0.25 ^{a,1}	7.37 ^{b,1}

Mean separation in columns by Duncan's multiple range test. In each column, values followed by the same letter do not differ significantly at $P=0.05$ and values followed by the same number do not differ significantly at $P<0.01$

Shrestha *et al.* (2011) found that diploid androgenic plants are characterized by big size fruit with high individual fruit weight and volume. Also, it was found that variation exist within diploid population might be die to the anther culture and naturally produced variation. Based in analysis of diploid populations as compared to control variety Boogie, researchers selected some elite diploid lines for development of sweet pepper varieties [25]. The most of androgenic lines were characterized with uniformity and higher values of fruit length, fruit weight and wall thickness than mother variety [20]. On contrary, other authors found lower mean values of fruit characteristics in androgenic diploid plants [21]. Nowaczyk *et al.* (2015) have found that seventy-four percent of studied androgenic plants differ from the donor plant phenotype in ripe fruit color, pericarp structure and taste. Other nine diploid androgenic plants showed differences from the mother genotypes in fruit weight, wall thickness and dry matter content [28]. Luitel and Kang (2013) reported higher dry matter content and improvement of other fruit quality parameters in double haploid lines of minipaprika [29].

3.3 Production traits of androgenic genotypes

The results which characterize production trait of pepper genotypes under this study are presented in Table 5. The average number of fruits per plant during the whole experimental period for androgenic genotypes P3 (5.65) and P4 (5.80) is smaller compared to the control genotype Piran (7.67 fruits).

The fruit number per plant for androgenic genotypes vary from 3.25 to 9.68 and only in II experimental year they are significantly higher than the initial genotype Piran. The average yield per plant is 249,81 g (Piran), 222.73 g (P3) and 231.90 g (P4) without significant difference for $P<0.01$. The highest yield per plant is 420.25 g (P4; III year), while the lowest is 97.5 g (P4; II year). The highest yield per plant for all genotypes under study are recorded during III experimental year.

Table 5. Productivity characteristics of androgenic genotypes P3 and P4 and the control Piran

Experimental year	Genotype	Number of fruits per plant	Yield per plant (g)	Total yield (kg/m ²)
—	Piran	8.05a	216.00a	1.350a
	P3	5.83a	218.13a	1.363a
	P4	5.25a	210.25a	1.314a
=	Piran	6.30a	204.5a	1.279a
	P3	4.70b	171.50a	1.072a
	P4	4.35b	166.00a	1.038a
≡	Piran	10.50a	384.00a	2.400a
	P3	8.23a	361.25a	2.258a
	P4	9.68a	420.25a	2.627a
≥	Piran	5.85a	194.75a	1.215a
	P3	3.25a	112.50a	0.703ab
	P4	3.3a	97.50a	0.609b
Average I-IV	Piran	7.67 ^{a,1}	249.81 ^{a,1}	1.561 ^{a,1}
Average I-IV	P3	5.65 ^{a,1}	222.73 ^{a,1}	1.392 ^{a,1}
Average I-IV	P4	5.80 ^{a,1}	231.90 ^{a,1}	1.449 ^{a,1}

Mean separation in columns by Duncan's multiple range test. In each column, values followed by the same letter do not differ significantly at $P=0.05$ and values followed by the same number do not differ significantly at $P<0.01$

The average yield for P3 is 1,392 kg/m² and 1.449 kg/m² for P4, while it is the highest for the control Piran (1,561 kg/m²). During the whole experimental period the yield (1,561 kg/m²). The total yield of P3 vary from 0.609 to 2.258 kg/m², while for P4 from 0.703 kg/m² to 2.627 kg/m². The control Piran gave from 1.215 kg/m² to 2.400 kg/m² during different experimental years.

The total and standard yield of 6 androgenic lines was higher compared to initial variety Hebar and they surpassed the control by standard yield from 11.02% to 42.26% [20]. Olszewska *et al.* (2015) tested eight double haploid lines of pepper for agro-morphological and molecular traits and found two double haploid lines deriving a special attention regarding yielding because they were either at the level of the maternal form or surpassed both parental forms regarding yield level [28].

3.4 Agronomic potential and evaluation of androgenic genotypes

3.4.1 Androgenic genotype P3

Androgenic genotype P3 is created from the Macedonian variety Piran via androgenesis in *in vitro* conditions. In the phase of maximum growth, the plant can reach height of 88.55 cm, while the mean stem height to the first branching is 44.29 cm with stem width of 0.94 cm. These traits classified P3 genotypes as tall pepper with medium-thick steam. The leaves are averagely 11.15 cm long and 5.41 cm width which characterized the genotype with large leaves. According to the fruit weight (41.45 g) and the fruit index higher than 4, the genotype is classified as long, horn-shaped, middle-size peppers. The fruit flesh of 77.05% and the pericarp thickness 0.26 cm characterize this fruit as medium-fleshy and make suitable for utilization as fresh and preserved. Regarding productivity traits, this genotype is characterized by 5.65 fruits per plant averagely weighting 222.73 g and total yield of 1.392 kg/m². According to production traits, in the given agroecological and cultivation conditions, it is a low-yield pepper genotype.



Fig. 2. Plants and fruits of androgenic genotype P3

3.4.2 Androgenic genotype P4

Androgenic genotype P4 is created from the Macedonian variety Piran via androgenesis in *in vitro* conditions. In the phase of maximum growth, the plant can reach height of 93.68 cm, while the mean stem height to the first branching is 38.62 cm which classified it as very tall pepper. The stem width is 1.01 cm. The height and steam thickness classified this genotypes as very tall pepper with medium-thick steam. The leaf index, a ratio of mean leaf length (10.73 cm) and mean leaf width (5.22 cm), is 2.07 which characterize the leaves as medium-large.

The fruit is characterized with weight of 42.84 g, contains averagely 7.37% dry matter, 76.71% fruit flesh and 0.25 cm pericarp thickness. P4 fruit is classified as long, horn-shaped, medium-size and medium-fleshy pepper. The total yield gained in this research is 1.449 kg/m². One plant is averagely loaded with 5.80 fruits weighting 231.90 g. According to production traits, in the given agroecological and cultivation conditions, it is a low-yield pepper genotype.



Fig. 3. Plants and fruits of androgenic genotype P4

4. CONCLUSION Conclude with future strategy

Analysis of morphological and production traits of androgenic genotypes provides valuable information for future breeding. Androgenic genotype P3 differ from the control variety for the traits: height to the first branching, leaf index, fruit length and fruit weight. Androgenic genotype P4 differ from the control variety for the traits: height to the first branching, stem thickness, leaf index, fruit length, fruit weight and fruit dry matter content. The results of this research prove the advantage of pepper androgenesis as a tool for enrichment of biodiversity and fast development of valuable pepper genotypes for future breeding.

REFERENCES cited as per journal guidelines

1. Bosland PW, Votava EJ. Peppers: Vegetable and Spice Capsicums. New York: CABI Publishing; 2000.
2. Koleva Gudeva L, Trajkova F, Zlatkovski V. Biotechnology and biodiversity: aspects of improvement of genetic resources of agriculture crops. Yearbook of Institute of Faculty of Agriculture, Goce Delcev University -Stip. 2008;VII:57 – 66.

- 305 3. Simaroo Segui JM. Androgenesis in Solanaceae. In: Germanà MA, Lambardi M. editors. *In vitro*
 306 embryogenesis in higher plants, methods in molecular biology, vol. 1359. New York: Springer
 307 Science+Business Media; 2016.
- 308 4. Irikova T, Grozeva S, Rodeva V. Anther culture in pepper (*Capsicum annuum* L.) in vitro. *Acta*
 309 *Physiolgiae Plant.* 2011;33:1559-1570.
- 310 5. Rodeva V, Koleva-Gudeva L, Grozeva S., Trajkova F. Obtaining haploids in anther culture of
 311 pepper *Capsicum annuum* L. and their inclusion in the breeding process. Yearbook of Faculty of
 312 Agriculture, Goce Delcev University – Stip. 2007;VII:7-17.
- 313 6. Nowaczyk L, Nowaczyk P, Olszewska D. Genetic analysis of anther culture-derived diploids of
 314 *Capsicum* spp. *The Journal of Horticultural Science and Biotechnology.* 2015;90(6):747-52.
- 315 7. Irikova TP, Kintzios S, Grozeva S, Rodeva V. Pepper (*Capsicum annuum* L.) anther culture:
 316 fundamental research and practical applications. *Turkish Journal of Biology.* 2016;40(4):719-26.
- 317 8. Olszewska D, Niklas-Nowak A, Nowaczyk P. Variation in the quantitative characters of
 318 androgenic pepper lines derived from hybrid *Capsicum frutescens* L. x *C. chinense* JACQ..
 319 *Vegetable Crops Research Bulletin.* 2010;73/1:5–11.
- 320 9. Pauk J, Lantos C, Somogyi G, Vági P, Ábrahám Táborosi Z, Gémes Juhász A, Tímár Z.
 321 Tradition, quality and biotechnology in Hungarian spice pepper (*Capsicum annuum* L.) breeding.
 322 *Acta Agronomica Hungarica.* 2010;58(3):259-266.
- 323 10. Lantos C. In vitro androgenesis induction in wheat (*Triticum aestivum* L.), triticale (X *Triticosecale*
 324 Wittmack), spice pepper (*Capsicum annuum* L.) and integration of the results into breeding.
 325 Thesis of the Ph.D dissertation. Godolo, Hungary; 2009.
- 326 11. Koleva Gudeva L, Trajkova F, Dimeska G, Spasenoski M. Androgenesis efficiency in anther
 327 culture of pepper (*C. annuum* L.). *Acta Hort. (ISHS).* 2009;830:183-190.
- 328 12. Koleva Gudeva L., Trajkova F. Seed production from pepper obtained in *in vitro* anther culture.
 329 Yearbook of Institute of Southern Crops – Strumica 2004/2005. 2005;IV/V:85-93.
- 330 13. Koleva Gudeva L, Trajkova F. Application of androgenesis as method for improvement of
 331 agricultural crops diversity. III Congress of ecologists of Macedonia with international
 332 participation. Book of proceedings. 2008; 284-290. Macedonian.
- 333 14. Trajkova F. Characterization and agronomic evaluation of some pepper lines (*Capsicum annuum*
 334 L.) obtained via androgenesis. Faculty of Agriculture, Ss. Cyril and Methodius University –
 335 Skopje, pp. 149. Macedonian
- 336 15. Koleva Gudeva L, Trajkova F. Genetic resources of *Capsicum* spp. In the genbank of Faculty of
 337 Agriculture at Goce Delcev University – Stip. III Congress of ecologists of Macedonia with
 338 international participation. Book of proceedings. 2008;303-308. Macedonian
- 339 16. Ministry of Agriculture, Forestry and Water management of Republic of Macedonia, Directorate of
 340 Seed and Seedling Material. National variety list of Republic of Macedonia. Skopje; 2008.
 341 Macedonian.
- 342 17. IPGRI, AVRDC and CATIE. Descriptors for *Capsicum* (*Capsicum* spp.). International Plant
 343 Genetic Resources Institute, Rome, Italy; the Asian Vegetable Research and Development
 344 Center, Taipei, and the Centro Agronómico Tropical de Investigación y Enseñanza, Turrialba,
 345 Costa Rica.; 1995.
- 346 18. Jankuloski D. Study of biological, morphological and qualitative traits of important populations of
 347 long peppers in Macedonia. PhD thesis. Skopje: Faculty of Agriculture; 1983. Macedonian.
- 348 19. Jankuloski D. Pepper and eggplant. Skopje: NIP BAS-TRADE; 1997. Macedonian.
- 349 20. Todorova V, Grozeva S, Rodeva V, Masheva S. Breeding evaluation of pepper lines obtained by
 350 in vitro anther culture. *Genetika.* 2013;45(2):601-10.
- 351 21. Luitel BP, Adhikari PB, Shrestha SL, Kang WH. Morphological characterization of anther derived
 352 plants in minipaprika (*Capsicum annuum* L.). *Korean Journal of Breeding Science.* 2012 Dec
 353 1;44(4):450-61.
- 354 22. Grozeva S, Rodeva V, Todorova V, Pundeva R. Obtaining of pepper plants via anther culture.
 355 *Genetic and Breeding.* 2009;38:25-31.
- 356 23. Inthichack P, Nishimura Y, Fukumoto Y. Diurnal temperature alternations on plant growth and
 357 mineral absorption in eggplant, sweet pepper, and tomato. *Horticulture, Environment, and*
 358 *Biotechnology.* 2013;54(1):37-43.
- 359 24. Gvozdenović, Đ. 2010: Pepper. Institute of Field and Vegetable Crops. Novi Sad: Tampograf;
 360 2010. Serbian
- 361 25. Shrestha SL, Luitel BP, Kang WH. Agro-morphological characterization of anther derived plants
 362 in sweet pepper (*Capsicum annuum* L. cv. Boogie). *Horticulture, Environment, and*
 363 *Biotechnology.* 2011;52/2:196-203.

- 364 26. Kisiąła A, Olszewska D, Niklas-Nowak A, Nowaczyk P. Biometrical characteristics of r2
365 generation of anther-derived pepper (*Capsicum* spp.) plants. Acta Agrobotanica. 2011;Vol.
366 64(3):53-58.
- 367 27. Koleva Gudeva L., Trajkova F. Anther culture of pepper: morphological characteristics of fruits of
368 androgenetic pepper lines (*Capsicum annuum* L.). Journal of Research in Agriculture.
369 2012;1(2):136-145.
- 370 28. Olszewska D, Niklas-Nowak A, Kisiąła A, Dzwonkowska M, Nowaczyk P. Agro-morphological
371 and molecular assessment of doubled haploid lines of pepper (*Capsicum annuum* L.). Zeszyty
372 Problemowe Postępów Nauk Rolniczych. 2015;580.
- 373 29. Luitel BP, Kang WH. Assessment of fruit quality variation in doubled haploids of minipaprika
374 (*Capsicum annuum* L.). Horticulture, Environment, and Biotechnology. 2013;54(3):257-65.