

Original Research Article

SERUM BIOCHEMICAL PARAMETERS OF LAYING JAPANESE QUAILS (*Coturnix coturnix japonica*) FED DIETS CONTAINING FERMENTED TARO COCOYAM (*Colocasia esculenta* var . *esculenta*) MEAL

Abstract

The study was conducted to evaluate the effect of fermented taro cocoyam meal on the serum biochemical parameters of laying Japanese quails. Two hundred and twenty five Japanese quails (*Coturnix coturnix japonica*) were randomly allotted to five dietary treatments (I –V) of 36 hens and 9 cockerels each. Each treatment was replicated thrice with 12 hens and 3 cockerels per replicate. In each of the five diets, 48 hours fermented taro cocoyam meal (*Colocasia esculenta* var. *esculenta*) was used to replace maize at 0%, 25%, 50%, 75%, and 100% respectively. Serum Na^+ , K^+ , CL^- , HCO_3^- , Creatinine, ALT, AST and Total bilirubin were determined on the 70th day. The results of serum biochemistry showed that, CL^- , HCO_3^- , Creatinine, ALT and AST were within normal range. However, it was observed that serum K^+ increased above the upper limit whereas serum Na^+ was slightly below the lower limit in all the treatments. Total bilirubin was above normal range beyond 25% replacement of maize in the diets. It was concluded that, maize can be replaced totally by fermented taro cocoyam meal without adverse effect on most serum biochemical parameters of laying Japanese quails. However, for optimum performance supplementation should not exceed 25%.

Keywords: Serum biochemistry, fermented taro cocoyam meal, maize and laying Japanese quails.

Introduction

There has been a call for substantial increase in protein intake of animal origin in developing countries. [1], observed that there is gross shortage of protein of animal origin in the diet of an average Nigeria. Strategies have been suggested to address the problem of shortage of animal protein in developing countries as human population continue to grow. These includes:

increasing the production of short production cycle animals such as poultry(quails), rabbits and pig as well as reduction in the cost of production of livestock .Feeding cost constitutes the largest cost in raising livestock(about 60-70% of total cost of production) . The above can be achieved by the use of less-expensive and non- conventional feed stuffs such as cocoyam, cassava, mango kernel, *Prosopis africana* seed coat etc. Taro cocoyam originated from Asia [2]. It is one of the most widely spread root crop grown throughout the humid tropics; about 60% of world production is grown in Africa and the remaining 40% in Asia and Pacific Island [2]. In Nigeria, the mature corms are roasted, baked or boiled and eaten. The corms of taro cocoyam provides easily digestible starch. Taro cocoyam is low in protein (1.1% fresh weight), has high content of P, Mg, Zn of any root and tuber crop and less in all the vitamins (except for nicotinic acid) and fairly rich in carotene [3]. Some of the non- starchy nutrients such as proteins, minerals and vitamins are concentrated in the outer peels of the corms [4]. Most varieties are acrid and some contain anti-nutritional factors such as Trypsin inhibitors, Phytates, Tannins, Saponins as well as Raphides [5]. Raphides are minutes bundles of crystals of Calcium oxalates(0.1- 0.4% fresh weight) which accounts for the irritating effect of the taro corms. Tannins interfere with protein digestibility, Phytate and Oxalate interferes with minerals availability, Trypsin causes depressive growth and pancreatic hypertrophy and Saponin causes permeability of the intestine. These effects could be removed or reduced by processing methods such as; boiling, fermentation, toasting and sun-drying etc. [6]. Cocoyam corms and its by- products have not received enough attention in terms of utilization as a livestock feed: [7] use taro cocoyam to replace all the maize in diets for broilers. [8] fed fresh(sun-dried) and boiled taro cocoyam meal to weaned pigs and observed that, the weaned pigs fed boiled taro cocoyam meal competed favorably with those fed maize only. A depressive growth was observed beyond 50%

replacement with those fed sun-dried cocoyam meals. [9] fed boiled taro cocoyam meal to growing Japanese quails and observed a depressive growth beyond 50% inclusion. [10] fed fermented taro cocoyam meal to growing Japanese quails and observed no depressive growth at 100% replacement with maize. However, when fermented taro cocoyam meal was tried on laying quails, the time of lay was delayed beyond 50% inclusions. Egg production was significantly ($P<0.05$) reduced beyond 25% replacement with maize [11]. These findings necessitated the need to assay the serum biochemical parameters of these laying quails as results have shown that, there is a positive correlation between body weight at six weeks and egg production in Japanese quails with some blood constituents [12]. Also because during fermentation a number of roles are played by micro-organisms in food processing: it could be positive or negative. Fermentation enhances the nutrients- vitamins and essential amino- acid by improving protein and fiber digestibility [6] whereas, the negative effect includes spoilage of food products and contamination by pathogenic micro-organism [13].

Materials and methods

The study was carried out in Cross River University of Technology Teaching and Research Farm, Calabar, Nigeria. A total of two hundred and twenty five Japanese quails (180 hens(females) and 45 cockerels(males)) of about 7 weeks of age with average weekly weights ranging from 133g to 163g were studied over a period of ten weeks. Quails were randomly allotted to five dietary treatments (I-V) of 36 hens and cockerels each. Each treatment was replicated thrice with 12 hens and 3 cockerels per replicate. In each of the five diets, 48-hours fermented taro cocoyam meal (FTCM) replaced maize at 0%, 25%, 50%, 75% and 100% as treatments I,II,III,IV and V respectively.

Unpeeled taro cocoyam corms were bought from Bendeghe village, Etung, Cross River, Nigeria. Corms were peeled and then chopped into sizeable chips of about 14g each. These chips were put into a big black plastic pot- like container with a tight lid and fermented in water at room temperature (28°C- 29°C) for 48 hours. Sun drying lasted for a week (7 days) and thus reduced

moisture content to less than 10%. The 48 hours fermented taro cocoyam corms and other ingredients were milled separately and used to formulate the experimental diets (Table 1). The experimental diets were analyzed according to the procedure of [14] and metabolizable energy was calculated using the method of [15]. The birds were managed intensively in cages of three tiers. Each tier was separated with wood. Wire mesh was used for the walls and doors to allow adequate ventilation/lighting. The dimension of each tier was 0.75m × 0.38m. Litter materials (wood shavings) were used on the wooden floor. Each tier was equipped with adequate drinker and feeding troughs. A floor space 0.007m to 0.009m per quail was provided. Artificial lighting was provided with the use of one (200 watts) bulb for each tier to ensure adequate feed intake. Feeds were weighed with a micro scale balance of 2kg serving to ensure a uniform amount across treatments. Quail were served 300gms of feed for the first week at about 8am on a daily basis, the quantity was increased by 50gms on weekly intervals. Fresh clean water was supplied ad-lib. Drinkers and feeders were washed and disinfected using izal when appropriate. At the end of the experiment (70th day/10th week), three (3) laying quails per treatment (one from each replicate) were randomly selected and weighed. Quails were slaughtered by cutting their jugular vein with a sharp knife. The blood samples were collected at slaughter into sterile vacutainers and the serum was separated by centrifugation at 750 g for 15 minutes and stored in a deep freezer until use for serum biochemical analysis. Creatinine was analyzed using SIGMA Kits according to [16]. The Standard Flame Photometer (Gallen Clamp) was used to determine HCO₃⁻, Na⁺, K⁺ and CL⁻. Liver function test was also conducted (ALT (IU/L), AST (IU/L)) according to the methods of [17] and [18].

Table1: Percentage composition of experimental diets with fermented taro cocoyam meal as a replacement for maize in the diets of laying Japanese quails.

Levels of inclusion.					
Ingredients	0%	25%	50%	75%	100%
Maize	46.00	34.50	23.00	11.60	0.00
Cocoyam	0.00	11.50	23.00	34.50	46.00
Soybean	20.80	21.00	19.70	19.90	20.00
Fish meal	4.20	4.60	5.80	6.00	6.00
Palm kernel cake	10.90	10.00	10.00	10.00	10.00
Wheat offals	10.00	10.00	9.75	9.10	8.30
Bone meal	7.00	7.00	7.00	7.00	7.00
Salt	0.50	0.50	0.50	0.50	0.50
Palm oil	0.10	0.40	0.75	1.00	1.20

Vit./min.premix	0.50	0.50	0.50	0.50	0.50
Total	100.00	100.00	100	100	100

Calculated nutrient:

Crude protein%	20.35	20.39	20.40	20.35	20.37
ME (MJ/kg)	2.674	2.670	2.585	2.554	2.538

Analyzed nutrients:

Crude protein (%)	21.20	20.89	20.78	20.63	20.59
ME (MJ/kg)	2.690	2.680	2.593	2.562	2.553

Statistical analysis:

The result of serum biochemical parameters were analyzed using the Standard Error of Mean (SEM).

Results and discussion

The results of the serum biochemical composition of laying Japanese quails fed diets containing 48-hour fermented taro cocoyam are presented in Table 2.

A condition of hyponatremia was observed across the treatments (126-133µmol/L) when compared with the established reference interval for 16weeks old female Japanese quails by [19] (135-155µmol/L). This result was in contrast with that of Nwani [B.Agric. Project.Uni Agric Makurdi,Nig, 2014] who reported that the serum sodium (Na⁺) of quails fed sun-dried

mango kernel meal (SMKM) was within normal range across treatments. These low levels could not be attributed to the experimental diets as quails fed control diet also showed similar result. It could be attributed to syndrome of inappropriate anti-diuretic hormone.

The serum potassium (K^+) of quails rose slightly above the upper limit (hyperkalemia) (6.8-7.2 $\mu\text{m/L}$) when compared with the normal reference range by [19] (3.6-5.0 $\mu\text{m/L}$). Serum potassium tells us about the heart condition. It controls the heart beat rate and the blood pressure [20 ; 21] the fact that the diversions from the normal range were not outrageous, the heart would not have manifested any symptom of disorder. Serum chloride was within normal range in all the treatments (98-100 $\mu\text{m/L}$) when compared with the normal reference range by [19] (98-106 $\mu\text{m/L}$).

This finding were different from that of Nwani [B.Agric. Project.Uni Agric Makurdi,Nig, 2014] who observed slight increase in serum chloride above upper limit in quails fed control diet and 50% SMKM. Hyperchloremia is sometimes associated with excessive fluid loss, Such as diarrhea. Hyperchloremia can be symptomatic with signs of weakness and intense thirst [22].

Serum bicarbonate (HCO_3^-) was within normal range (22-24 $\mu\text{m/L}$) as established by [19] (22-30 $\mu\text{m/L}$). This result was in agreement with the report of Nwani [B.Agric. Project.Uni Agric Makurdi,Nig, 2014] who observed that serum HCO_3^- of quails fed SMKM was within normal range. This implies that, the acid-base ratio was balanced. Low levels of serum HCO_3^- results in metabolic acidosis [23]. HCO_3^- of less than 22 $\mu\text{mol/L}$ is compatible with metabolic acidosis.

Serum Creatinine was within normal range (4.0 - 6.2 $\mu\text{m/L}$) comparing with the established reference range by [19] (1.2 – 6.7 $\mu\text{m/L}$).

The report of Creatinine agrees with that of Nwani [B. Agric. Project.Uni Agric Makurdi, Nig, 2014] who had similar results but was in contrast with the reports of [24] who observed values below the lower limit across treatments. Serum AST and ALT were within normal ranges (242 - 538 IU/L; 4.5 - 8.0 IU/L respectively) when compared with the established reference range by [19] and [25] (243-562 IU/L (AST); 4.5-8.5 IU/L (ALT). These results were in agreement with those of Nwani [B.Agric. Project.Uni Agric Makurdi,Nig, 2014] and [24] who fed SMKM to laying quails and Garcinia kola to broilers respectively. This implies that, the liver of quails fed fermented cocoyam were normal and healthy as high values usually represent liver damage. Total bilirubin was slightly above the upper limit (15.17 $\mu\text{m/L}$) beyond 25% replacement of maize in the diet when compared with the normal reference range by [19] (3.6 -14.2 $\mu\text{m/L}$). High

value of bilirubin signifies liver disease such as hepatitis or blockage of tubes (bile ducts) or diagnose conditions that cause increased destruction of red blood cell [25]. The deviations observed could be associated with diagnose conditions that would have caused increased destruction of the red blood cells as the birds were intermittently treated with pentheryl (antibiotic) as serum ALT and AST which are equally used to ascertain the condition of the liver were within normal ranges.

Table 2: Serum biochemical parameters of laying quails fed varying levels of 48 – hours fermented taro cocoyam (*Colocasia esculenta* var. *esculenta*) meal as a replacement for maize.

Levels of inclusion.

Parameter	0%	25%	50%	75%	100%	Mean	SEM
Na ⁺ (μmol/L)	28.0	126.0	132.0	130.0	133.0	109.8	2.86
K ⁺ (μmol/L)	12.80	12.90	13.00	13.10	13.20	13	0.16
CL ⁻ (μm/L)	98.00	96.00	100.00	97.00	98.00	97.8	0.48
HCO ₃ ⁻ (μm/L)	23.0	22.0	24.0	24.0	24.0	23.4	0.89
Creatinine(μm/L)	4.00	6.20	4.00	4.80	4.60	4.72	0.90
ALT(IU/L)	4.50	5.00	4.50	4.80	8.00	5.36	0.67
AST(IU/L)	242.0	253.0	262.0	332.0	538.0	325.4	124
Total bilirubin(μm/L)	13.0	14.0	15.0	16.0	17.0	15	1.58

Conclusion

This study revealed that, maize can be replaced totally by fermented taro cocoyam meal without adverse effect on most serum biochemical parameters of laying Japanese quails. However, for optimum performance supplementation should not exceed 25%.

References

1. Ogbonna JU. "Nutritional value of cassava peels meal for poultry". *A Review of the Proceeding of 2nd Annual Conference of Animal Science Association of Nigeria*, Lagos. 1997.
2. Viotti V. Taro production in Hawaiian Island. *Honolulu Advertiser*; March 16, 2004.
3. Kelsely TL. *Effect of oxalic acid on calcium bio- availability*. In: Kelsely ed. *Nutritional calcium*. Washington, DC: American Chemistry Society; 1985.
4. Ohtsuka R, Kawobe, J, Inaoka T, Suzuk T, Hongo T, Akinchi SugaharaT. Composition of local and purchased feed consumed by the Gilra in low land Papua. *Ecology of Food and Nutrition*. 1984; (15):159-169.
5. Will RBH, Lin JSI, Greenfield H, Bayliss Smith T. Nutrient composition of taro *C. esculenta* cultivars from Papua New Guinea highlands. *Journal of Food Science and Agriculture*. 1983; 34:1137-1142.
6. Obi MB. Improving the nutritional value of taro cocoyam (*Colocasia esculenta* var *esculenta*) through processing for the feeding of growing and laying Japanese quails(*Coturnix coturnix japonica*). *Ph.D Thesis Dissertation*, University of Calabar. 2010.
7. Abdulrashid M, Agwunobi LN, Jokthan GE. Abdul SB. Carcass Quality Characteristics of Broilers Finisher Fed (*Colocasia esculata* var *esculata*). *A Book of Proceedings of the 32nd Annual Conference of the Nigerian Society of Animal Production (NSAP)*. 2007.
8. Agwunobi LN, Agwukan PO, Cora OO Isika MA. Studies on the use *Colocasia esculenta* (taro cocoyam) in the diets of weaned pig. *Tropical Animal Health and Production*. 2002; 34(4):243-247.
9. Okon BI, Obi MB, Ayuk AA. Performance of quails (*Coturnix coturnix japonica*) fed graded levels of boiled sun-dried taro cocoyam (*Colocasia esculenta*) as a replacement of Maize. *Medwell Online Agricultural Journal*. 2007; 2 (6): 654-657.

10. Abang FB, Ayuk AA Okon B I. Growth performance of growing Japanese quails (*Coturnix coturnix japonica*) fed 48-hours fermented taro cocoyam (*Colocasia esculenta* var. *esculenta*) as a replacement for maize. *Indian Journal of Research Paripex*. 2013; Pp: 288 - 289.
11. Obi MB, Ayuk AA Okon BI. Replacement value of 48-hours fermented taro cocoyam(*Colocasia esculenta* var *esculenta*) meal for maize on performance of laying quails (*Coturnix coturnix japonica*).*Proceedings of British Society of Animal Science and the Association of Veterinary Teaching and Research Work, UK*. 2013; Pp: 180-181.
12. El-Daen MB, Kosba MA, Soliman ASA. Studies of some performance and blood constituents traits in Japanese quail. *Egypt Poultry Science*. 2009; (29) 1187-1208.
13. Ojokoh AO. Effect of fermentation on the chemical composition of mango (*Mangifera indica*) peels. *African Journal of Biotechnology*. 2007; 6(6):1979-1981.
14. Association of Official Analytical Chemist. *Official Methods of Analysis 17th ed.* Washington D C. 2000.
15. Pauzenga U. Feeding Parent Stock. *Zootechnica International*, 1985; Pp.22-24.
16. Feteris WA. A serum glucose method without precipitation. *Animal Journal of Medical Technology*. 1965; (31): 17-21
17. Evans GO. Animal Clinical chemistry: *A Prime for Toxicologist*. Taylor and Francise; 1996.
18. Meyer DJ, Cole EH, Rich LT. *Veterinary Laboratory Medicine Interpretation and Diagnosis*. Philadelphia: Sauders WB Company; 1992.
19. Scholtz N, Halls I, Aachowsky G Saverwein H . Serum chemistry reference value in adult Japanese quails (*Coturnix Coturnix Japonica*) including sex- related difference. *In: A Research Note on Health and Diseases. Institute of Animal Science, Physiology and Hygiene Unit. University of Bonn, Katzen burgweg7-9, 53115 Bonn, Germany*.2009.
20. El-boushy AR, Vender Poel FB. Feather meal: A biological waste, its processing and utilization as a feed stuff for poultry. *Biological waste*. 1990; (32)39-74.
21. Diarra SS, Kwarri ID, Asheikh LG, Muhammed G Igwebuikue JU. Growth Performance of Broiler Chicken Fed Raw or Boiled Mango(*Mangifera indica*) Kernel Meal as Source of Dietary

Energy. *Proceedings of the 13th Annual Conference of Animal Science Association of Nigeria*,
Abu-Zaria. 2008; Pp:311-31

22. Cambier C, Detry B, Beerens D. “Effects of hyperchloremia on blood oxygen binding in
healthy calves”. *Journal Applied Physiology*. 1998; 85(4): 1267-1272. PMID976031

23. Kasimatis E, Maksich D, Jassal V, Borgman J M, Oreeopoulus O G. Predictive factors of low
HCO₃⁻ levels in peritoneal dialysis patients. *Medline Clinical Nephrology*. 2005; 63(4):290-6.
PMID1584725.

24. Sobayo RA, Adeyemi OA, Oso AO, Fafiolu AO, Daramoa JO, Sodipe G *et al* .
Haematological, Serum and Carcass Characteristics of Broilers Chicken fed graded levels
of Garcinia kola (bitter kola) used as Phytobiotic. *Nigeria Society for Animal Production*. 2013;
Pp : 48-56.

25. Robin W. *Laboratory test interpretation of bilirubin*. .United States National Library of
Medicine, National Institute of Health; (2013).