

Effect of Hemoglobin and Potassium polymorphism on milk production traits in goats

ABSTRACT:

A study was conducted to determine the effect of hemoglobin (Hb-type) and potassium (k-type) types on milk yield characteristics of Agropastoral goats. Records were obtained from a total of 250 does over a period of one year. The milk yield characteristics were initial milk yield (IY), average daily yield (ADY), initial yield (IY), peak day (PD), peak yield (PY), and lactation length (LL). The data generated from the study were analyzed using General Linear Model (GLM) procedure of SAS and Duncan Multiple Range Test (DMRT) procedure of SAS was used to separate the significant means. Hb-type significantly influenced some milk yield characteristics of the goats. Does with HbAA and HbAB were similar and significantly had ($P < 0.05$) higher ADY and PY than HbBB. But there were no significant ($p > 0.05$) differences in milk yield characteristics when potassium types were considered, however animals of high potassium (HK) types were found to be favoured by natural selection in the region over low potassium (LK). It was concluded that, potassium and hemoglobin types can be explored to help in improving milk production traits of indigenous goats by identifying the hemoglobin and potassium types that are more adaptive to a particular environment and has a comparative advantage in milk yield characteristics.

Key words: Hemoglobin types, Potassium types, Milk production traits, Goats

INTRODUCTION

Nigeria is endowed with large numbers of small ruminants which are yet to be fully explored for milk production. Goats are the largest in number among ruminant livestock, totaling about 53.8 million (FAOSTAT, 2008). Goats are predominantly owned by the rural households and ownership is spread across all human age groups and sexes (Akpa, 1999). They are kept mostly for meat, the milk is rarely used, but the awareness on importance of goat milk is rapidly gaining attention. Apart from the ease of management and high proliferation, goats have high potential for milk production (Adebambo *et al.*, 2011). Genetic variations within population of animals are of great importance to breeders. This is because the amount of genetic variation seen in a population of animals determines the magnitude of genetic improvement that can be achieved within the species (Salako *et al.*, 2007; Akpa *et al.*, 2011). Electrophoresis techniques have been used to detect polymorphism at protein and enzyme loci for measurement of variation (Salako *et al.*, 2007). Information obtained from this type of study have been useful as genetic markers for important economic traits and aids significantly in selection of superior animals for breeding purposes (Salako *et al.*, 2007; Akpa *et al.*, 2011). Blood biochemical polymorphism in domestic animals is used for parentage determination and indirect selection criteria for livestock species (Sobiech *et al.*, 2008). Blood polymorphism studies have been conducted extensively to identify biodiversity among livestock animals. (Gurcan *et al.*, 2010; Shahrababak, *et al.*, 2009; Nedjar *et al.*, 2008; Nihat *et al.*, 2003). Potassium ion is an essential in regulating osmotic balance between cells and the extracellular fluid (Ellory and Tucker, 1970; Campbell, 1987). Guney *et al.*, (2003) explained that potassium concentration in red cells of small ruminant are controlled by two alleles on a single gene, and they are allele K^H which is responsible for high potassium concentration in blood while allele K^L is responsible for low potassium concentration. K^L is

completely dominant over K^H. Evans *et al.* (1956) identified the HbAA and HbBB as single bands with HbAA being the fastest single band towards the anode during electrophoresis while HbBB was the slowest. The HbAB, however, was identified as a double band having both the fast moving AA and slow moving BB. Das *et al.*, (2004) and Salako *et al.*, (2007) also reported similar haemoglobin types, but occurrences for each haemoglobin type vary from location to location.

It has been evidenced by several researchers that haemoglobin and erythrocyte potassium types in domestic animals are associated with some production/reproduction traits such as milk production, body weight, mortality rate, fleece weight, fertility and adaptation capacity (Al-Samarrae and Younis, 2011; Ekmekci and Mert, 2009; Antunovic *et al.*, 2004; Gurcan *et al.*, 2010; Akinyemi and Salako, 2010;). Moreover haemoglobin and potassium can be determined easily at the postnatal period of young animals and these components are not affected by the environmental factors, and can reduce the time of rearing inferior animals before selection. The relationships between blood polymorphism and milk production traits could be used as indirect selection criteria to improve animal production levels. In Nigeria, a lot of researches have been done to evaluate milk yield characteristics of goats in terms of genetic and non-genetic factors, but there are limited information on relationship between milk yield characteristics with haemoglobin and potassium polymorphism. Therefore, the objectives of this study was to investigate the relationship between these haemoglobin and potassium polymorphic genes and milk production traits.

MATERIALS AND METHODS

Study Area

The Study was carried out in two States within the North-West Nigeria. The States were Jigawa and Katsina State. Katsina lies between latitude 13°01' North of the equator and longitude 07°41' East of Greenwich meridian. It is situated at an altitude of 464m (1525ft) above sea level and is located in the Sudan savannah agro-ecological zone. It has two distinct seasons. The wet season which lasts from June and ends September. The dry season is experienced between October and May. Mean annual rainfall is about 780mm. Katsina State is hot for most parts of the year, even during the wet season. Mean annual temperatures range between 19° in the cold dry season to 38°, with the highest temperatures normally experienced in April and May, just before the rains (Ati *et al.*, 2007). Jigawa State falls within latitude 11°-13°N and Longitude 8-10°E. Temperatures can go as low as single digits in the nights in most of the Northern parts during the Harmattan season (December to February), and could also rise as high as 45 degrees during the hot season (April to June). There are two seasons in Jigawa State namely: rainy and dry seasons with the dry season lasting from November to May and the rainy season lasting between June and October, (Magashi, 2011).

The Animals and Management

The goats were under the agropastoral management. The animals were taken out to graze every morning from 8.00 am to 5.00 pm by children and were penned at night in small open sided shades by tethering. Before setting out for grazing every morning, the goats were given water and supplement which included groundnut haulms, bean haulms, bean pods and dry grasses

containing 53.7% and 10.5% TDN and CP respectively. The young goats were left to suckle their dams freely for the first 6 days postpartum to enable them take advantage of colostrums.

The Measurements

A total of 250 does (whose parities were between 1-6 and average body weight of 28.84 kg) were used for this study. All milk measurements commenced from day 7 postpartum and were taken till the milk yield was less than 100ml.

During milking, lactating does were kept calm by providing supplements in a feeder for them. The quantity of milk available per doe per test day was measured using graduated plastic beakers. Milk yield was determined twice a week.

The young ones were separated from their dams at 18.00hours on the evening proceeding the day of milking. On the test day, the two halves of the udder of each lactating doe were hand-milked for all herds from 06.00 to 08.00hours and the milk yield was recorded to the nearest gram. The average of the total volume of milk collected for the two test days was taken as the average daily yield of the doe for that week. Milk recording was terminated at the point where a doe was unable to produce 100ml and above.

Milk yield characteristics were measured as follows:

Average Daily Yield (ADY): - As average of all test day yields within the period milked.

Initial Yield (IY): - As milk yield at day 7 postpartum

Total Yield (TY): - As milk production during the study before the production drops below 100ml.

Peak Yield (PY): - As the yield with the highest test day yield.

Peak day (PD): - As the day with the highest yield.

Lactation Length (LL): - As the period during which the doe was milk.

Blood Collection and Preparation

5ml of blood was collected from each of the sampled animals by jugular venipuncture, using needle and syringe into test tube containing Ethylene Diamine Tetra Acetic acid (EDTA) as anticoagulant and properly labeled and samples were taken to hematological laboratory of Bayero University Kano Teaching Hospital, Kano. The blood samples were then washed with normal saline and haemolysed with distilled water to release the haemoglobin. The supernatant was removed after centrifuging at 3000 rpm for 5 min and the sample haemoglobin stored until ready for electrophoresis. Cellulose acetate paper strip was used to separate the globin fractions. Electrophoresis was carried out in Shandon electrophoresis tank on cellulose acetate strips 34.5 x 150 with 0.26 M Tris buffer (pH 9.1) at the anode and cathode. The strips were run for 5 minutes at a constant voltage of 250v until a clear separation was observed. Interpretations were made based on the relative mobility of the

haemoglobin bands towards the anode. The genotype that migrated faster was labeled HbAA; the slow moving fraction was identified as HbBB. The double band, consisting of both fast and slow band; was labeled HbAB as described by Tella *et al.*, (2000) and Das *et al.*, (2004). The potassium concentrations in the blood were determined by colorimetric method using spectrophotometer. The erythrocyte potassium below or equals to 13.00 mmol/l was labeled low potassium, while the erythrocyte potassium above 13.00 mmol/l was tacked high potassium as describe by Galip and Elmaci (2001).

Statistical Analysis

The data generated from the study were analysed using General Linear Model (GLM) procedure of SAS (SAS, 2000). The model incorporated Hb-type and K-type, as fixed factors while milk yield characteristics were the dependent variables. Duncan Multiple Range Test (DMRT) procedure of SAS was used to separate the significant means. The linear model was as follows:

$$Y_{ij} = \mu + K_i + H_j + E_i$$

Where

Y_{ij} = Measurement on traits

μ = Overall mean

K_j = Effect of potassium types (K = Lk, Hk)

H_j = Effect of hemoglobin types (k = AA, AB, BB)

E_{ij} = Random error effect

Results

Heamoglobin types and their Relationship with Milk Yield Characteristics in goats

The influence of Hb type on milk yield characteristics is presented in Table 1. The results shows that only ADY and PY were significantly ($P < 0.05$) influenced by Hb types. Does with HbAA and HbAB were similar and significantly had ($P < 0.05$) higher ADY and PY than HbBB. All other milk yield characteristics followed the same trend, though not significant.

Table 1: Mean Values ($\pm$ se) of milk yield characteristics of goats according to Heamoglobin Types

Milk yield characteristics	HbAA	HbAB	HbBB	LOS
ADY(g)	253.70 $\pm$ 2.92 ^a	257.00 $\pm$ 2.76 ^a	240.55 $\pm$ 5.55 ^b	*
TY(kg)	30.19 $\pm$ 0.59	30.32 $\pm$ 0.58	27.32 $\pm$ 13.00	ns
IY(g)	241.00 $\pm$ 4.93	244.42 $\pm$ 0.58	247.43 $\pm$ 9.20	ns
PD(days)	34.60 $\pm$ 0.69	32.88 $\pm$ 0.67	32.79 $\pm$ 1.27	ns
PY(g)	359.72 $\pm$ 5.66 ^b	385.00 $\pm$ 38.15 ^a	350.17 $\pm$ 7.89 ^b	*
LL(days)	119.00 $\pm$ 1.57	118.73 $\pm$ 1.54	113.60 $\pm$ 2.99	Ns
No of observation	108	112	30	

Values bearing different superscript at the same row differ significantly ($P < 0.05$);

*= $P < 0.05$; ns = non-significant , ADY = Average Daily Yield, TY = Total Yield,

IY=Initial Yield, PD = Peak Day, PY =Peak Yield, LL = Lactation Length

Potassium types and their Relationship with Milk Yield Characteristics in goats

The effect of potassium types on Milk yield characteristics is shown in table 2. The results indicate that K-types had no significant ($p>0.05$) influence on milk yield characteristics. Although there were no significant ($p>0.05$) difference, animals with HK performed better in all the milk yield trait **evaluated** (Table 2).

Table 2: Mean Values ($\pm$ se) of milk yield characteristics of goats according to potassium Types

Milk yield characteristics	HK	LK
ADY(g)	248.41 $\pm$ 2.01	247.19 $\pm$ 4.69
TY(kg)	29.65 $\pm$ 0.59	29.25 $\pm$ 0.99
IY(g)	244.36 $\pm$ 3.39	239.33 $\pm$ 7.90
PD(days)	33.09 $\pm$ 0.47	32.80 $\pm$ 1.11
PY(g)	360.41 $\pm$ 4.05	349.64 $\pm$ 9.43
LL(days)	118.32 $\pm$ 1.12	117.87 $\pm$ 2.62
No of observation	211	39

ADY = Average Daily Yield, TY = Total Yield, IY = Initial Yield, PD = Peak Day, PY =Peak Yield, LL = Lactation Length, K-type = potassium types, HK = high potassium, LK = low potassium

DISCUSSION

The observed variation in milk yield characteristics (ADY and PY) with Hb-type had earlier been reported in other species. Samarineanu *et al.*(1982) and Gootwine (1988) in cow and ewe, respectively observed that HbAB was superior to both HbAA and HbBB in milk yield. However, Essien (2011) observed that cows with HbAA were superior to HbAB and HbBB in some milk yield characteristics in cows studied in Zaria. On the contrary, Dally *et al.*, (1980) reported that there were no differences in the milk production of ewes with different haemoglobin genotypes. The variation in the performance of the animals due to differences in Hb-types at different location suggests that different Hb-types may have selective advantage in different geographical region (Ndamukong, 1995). The observation in this study suggests that in the study area, Jigawa and Katsina states indigenous goats with HbAA and HbAB have high potential for milk yield than those with HbBB.

The observed non-significant influence of potassium types on milk yield characteristics in this study is contrary to the reports of Gunes *et al.*, (2008) which reported significant influence of potassium types on total milk yield of Awassi sheep, with LK being superior to HK. Limited literature is available on the influence of potassium types and milk yield characteristics in small ruminants. However, a careful study of the means of milk yield characteristics according to potassium types shows that HK, though not significant, were superior to the LK type animals. This could be attributed to association of this gene with aspect of fitness, with animals of HK types being favoured by natural selection in the studied region. **Large sample size is required to confirm the findings of this work as this will remove bias and tendency for results to skew towards one direction.**

Conclusion:

The findings in this work did not give strong reasons to conclude that potassium polymorphism could be used as selection criteria for milk production in goats. However, the non-significant variation of most milk yield characteristics with haemoglobin and potassium genotypes suggests

that haemoglobin and potassium genotypes are more of adaptive traits than productive traits. Therefore, potassium and haemoglobin which are genetic markers can be explored to help in improving milk production traits of the indigenous goat breed by identifying the potassium and haemoglobin types that are more adaptive to a particular environment and has a comparative advantage in milk yield characteristics.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

REFERENCES:

- Adebambo O. A. Samuel, A. O. and Onakade, A. D. (1994). Causes of Variation in Productive performance of West African Dwarf Goat. *Nigerian Journal of Animal Production*, 21:29-36.
- Akpa, G. N., Ibrahim, O. A., Yakubu, H. and Kabir, M. (2011). Sexual dimorphism, haemoglobin polymorphism and body mensuration characteristics of Red Sokoto goats. *Proceedings of the 35th Annual Conference of Genetic Society of Nigeria* 10th -14th October, 2011, Ahmadu Bello University, Zaria. Pp. 2-7
- Al-Samarrae, S.H. and Younis, K.H. (2011). Blood Polymorphism and its Relation with Sheep Production and Reproduction *Diyala Agricultural Sciences Journal*, 3(1) 10-21.
- Antunovic, Z., Sperenda, M., and Steiner, Z., (2004). The influence of age and the reproductive status to the blood indicators of the ewes. *Arch. Tierz*, 47: 265-273.
- Ati, O.F. Iguisi, E.O. and Afolayan, J.O (2007). Are we experiencing drier conditions in the SudanoSahelian Zone of Nigerian? *Journal of Applied Sciences Research*, 3 (12): 1746-751.
- Campbell, N. (1987). Textbook of Clinical Chemistry. Benjamin/Cummings Pub. Co., Menlo Park, Calif, ISBN 0-8053-1840-2, pp:795
- Dally, M. R., Hohenboken, W., Thomas D.L. and Craig, M.C. (1980). Relationship between haemoglobin type and reproduction, lamb wool and milk production and health related traits in crossbred ewes. *Journal Animal Science*, 50: 418-427.
- Das, D. K., Sinha, R., Dattagupta, R. and Senapti, P.K. (2004). Association of Haemoglobin types with some menstruation and reproductive characteristics in Garole sheep. *Indian Journal of Animal Sciences*, 74(4): 382-384.
- FAOSTAT (2008). <http://faostat.fao.org/default.aspx>

213 Ekmewski, S. and Mert,H.(2009).The investigation of heamoglobin,erythrocyte potassium and
214 gluthathion types in Norduzgoats.YuzuncuYil University, (In Turkish). J. Agric. Fac. 20:23-26.

215 Ellory, J. C. and Tucker, E.M (1970). High Potassium type red cell in cattle *J. Agric.*
216 *Science.Camb.* 74, 595-596.

217 Essien, I. C. (2011).Relationship between Haemoglobin polymorphism and Dairy performance
218 of Bunaji and Friesian X Bunaji cows. M.Sc. Thesis. Ahmadu Bello University Zaria.

219 Evans, I. V., King, J.W.B., Cohen, B. L. Harris, H. and Warren, F. L.(1956). Genetic of
220 Hemoglobin and blood potassium differences in Sheep.*Nature*, 4538: 849-850.

221 Galip, N., and Elmaci, c. (2001).Erythrocyte potassium polymorphism and relationship with
222 blood parameter in Turkish Hair Goats. J. Genet, Breeds, 55:183-185.

223 Gunes, N., Aydin, C., udum, C. O. Dikmen, s. and Ozmen, O (2008). Erythrocyte potassium,
224 sadium and glutathione concentrations and their relationship with reproduction, body
225 weight and fleece weight traits in Awassi sheep, Arch. Tierz, 51:479-486.

226 Guney, O., Ozuyanik, O., Torun, O, Gorgulu, M. and Darcan, N, (2003). Relationship between
227 some polymorphic parameters and performances in Damascus goats Pakistan Journal of
228 Biological Sciences 6(8): 738-740.

229 Gurcan, E. K. Erbas, C., Cobanoglue, O., (2010). Biochemical polymorphism of erythrocyte
230 potassium and glutathione protein. The relationship with some blood parameters in Kirvircik
231 sheep breed Afr. J. Agric. Res., 5:1022-1027.

232 Magashi, A. (2011). Jigawa State in brief. WWW.jigawastate.com.

233 Ndamukong, K.J.N.,Sewell ,M.M.H and Asanji, M.F. (1989). Management and productivity of
234 Small Ruminants in the North West Province of Cameroon.*Tropical Animal Health and*
235 *Production*, 21:109-119.

236 Nedjar A. N., Duddois-Deival, Adje, Y. E., Traisnel, J., Krier, F., Mary, P., Kouach., Briand, G.
237 and Guillochon. (2008). Bovine hemoglobin an attractive source of antibacterial.
238 Peptides:29(6):969-977.

239 Nihat, M., Handen, G., and Velat, A. (2003). Correlation between biochemical parameters and
240 production traits in Merino crosses sheep. Haemoglobin and Transferrin tyes.Turk.J.Vet Anim.
241 Sci:27:575-581.

242 Samarineanu, M. Stamatescu, E., Granciu, S. M. and Sotu, N. (1982).The results of
243 electrophoretic studies of some proteins in the blood and milk of Romanian Brown cows in
244 Moldavia.*LucrariStintifice Institute de Cercetar-PentruCrestereaTaurinetor*, 8:45-56.

- 245 Salako, A. E., Ijadunola, T.O. and Agbesola, Y.O. (2007).Haemoglobin polymorphism in
246 Nigerian Indigenous small ruminant populations preliminary investigation. *African Journal*
247 *Biotechnology*, 6(22): 2636-2638.
- 248 Shaharbabak, H. M.,Shaharbabak, M. M., Rahimi, G. H. and Yegane, H. M. (2009).Association
249 of whole blood potassium polymorphism with resistant to saline in two sheep of differnt
250 climates of Iran. *Desert(Biaban)*:14(1):95-99.
- 251 Sobiech, P., Milewski, S., and Zdunczyk, S. (2008). Yield and Composition of Milk and Blood
252 Biochemical Components of Ewes Nursing a Single Lamb or Twins. *Bull Vet Inst Pulawy* 52,
253 591-596
- 254 SAS,2000. Statistical Analysis System, version 8.0. SAS Institute Inc. Cary. North Carolina,
255 USA.
- 256 Tella, M. A., Taiwo,V. O., Agbede, S. A. and Alonge, O. D. (2000). The influence of
257 hemoglobin types of incidence of babesiosis and anaplasmosis in West African Dwarf and
258 Yankasa sheep. *Tropical Veterinary Journal*, 18:121-127.