

Original Research Article

Influence of bio-regulators on the fruit quality and yield attributes of phalsa (*Grewia subinaequalis* DC)

ABSTRACT

Aims: To examine the influence of bio regulators on the fruit quality and yield attributes of phalsa.

Study design: Randomised Block Design.

Place and Duration of Study: Phalsa orchard, Department of Horticulture, Khalsa College, Amritsar during 2015-2016

Methodology: The present study comprised of ten treatments which replicated thrice Uniform sized mature fruits were harvested for carrying out physico-chemical analysis. The observations were recorded viz., fruit length (cm), fruit breadth (cm), fruit weight (g), fruit colour, TSS (%), titratable acidity (%), TSS: acid ratio, reducing sugars (%), total sugars (%), ascorbic acid (mg/100 g pulp), stone length (cm), stone breadth (cm), stone weight (g), harvesting span (days), number of pickings and fruit yield per plant (kg). A sample of ten randomly selected fruits was being taken from each replication for fruit physical and bio chemical analysis and then average was worked out for each parameter.

Results: The results revealed that GA₃ 150 ppm treated plants yielded fruits with maximum fruit length (1.26 cm), fruit breadth (1.43 cm), fruit weight (1.20 g), reducing sugars (7.49 %), ascorbic acid (31.10 mg/100 g pulp) and yield (3.48 kg/plant). Minimum stone length (0.50 cm), stone breadth (0.33 cm) and stone weight (0.045 g) was recorded in fruits obtained from plants treated with NAA 25 ppm. Ethrel 1000 ppm significantly improved TSS (20.23%), fruit colour (purple black), total sugars (12.79 %) whereas reduced the titratable acidity (1.54 %), harvesting span (22.81 days) and number of pickings (3.90).

Conclusion: The results of the present study concluded that plants treated with GA₃ 150 ppm registered significant improvement in fruit physical and biochemical parameters and also fruit yield per plant was increased. The fruits obtained from the plants treated with NAA 25 ppm had small sized and minimum weighed stone. Whereas the yield attributes were enhanced with the application of ethrel 1000 ppm in terms of advancement of harvesting and also reduction in number of pickings. Hence Ethrel 1000 ppm may be applied to improve the fruit quality parameters as well as yield parameters and also to fetch the premium price in the market due to their attractive colour appearance than control.

Keywords: Phalsa; bio-regulators; fruit physical parameters; fruit biochemical parameters; yield

1. INTRODUCTION

The Phalsa (*Grewia subinaequalis* DC) belonging to the family Tiliaceae, is an important indigenous fruit for arid and semi- arid region because of its hardy nature and capacity to tolerate high temperature and even to be grown under prolonged dry spells [1]. India is considered to be the home of phalsa and it is commercially cultivated in Punjab, Uttar Pradesh, Madhya Pradesh, Haryana and Rajasthan [2]. In Punjab it is mostly grown in Ropar, Hoshiarpur, Gurdaspur and Amritsar district. Area under phalsa in Punjab is only 30 hectares with an annual production of 196 tonnes [3].

Comment [AL1]: for the first time the full name and the abbreviation in brackets

Comment [AL2]: what (generally, because bottom there are details) and how (methods)

Comment [AL3]: ?

Comment [AL4]: ?

Plant growth regulators are chemicals that effect flowering, ageing, root growth, prevention and promotion of stem elongation, color enhancement of phalsa fruits. Very small concentration of these substances produces major growth changes. They are widely used for increasing fruit set, controlling fruit drop, enhancing quality and uniform maturity. Among growth regulators GA₃, NAA, Ethrel are widely used for enhancing superior characters of fruit crop. Application of GA₃ results in increased yield and better grade of phalsa [4]. The use of bio regulator like GA₃ has proved effective for increasing the size of berry or fruit and improved quality in crop like grape, citrus, ber etc [2]. Parthenocarpic development of fruits is also a result of gibberellins action [17]. It is thought that NAA may influence the rate of metabolic activity of developing embryos and even also stimulate ethylene biosynthesis which inhibit fruit growth and promote the abscission of weaker fruits. Naphthalene acetic acid shows positive attitude towards the reduction of fruit drop in many fruit crops.

Application of ethrel improves fruit quality in terms of TSS, reducing and total sugar [5]. Ethrel is a compound that decomposes to release the natural plant hormone ethylene. Its main use in fruit production is to enhance fruit ripening by permitting its harvest about one week earlier and also reduces the span of harvesting and number of pickings significantly [5]. The bio regulators have proved to be effective in increasing fruit set and yield in phalsa as well as quality of fruit in terms of TSS, reducing sugars and total sugars. Keeping the role of bio regulators in view, the present study was carried out for improved fruit set, quality and yield of phalsa.

2. MATERIALS AND METHODS

The trial was conducted on well maintained 9 years old phalsa orchard of Department of Horticulture, Khalsa College Amritsar during 2015-16. In the trial 30 uniform sized trees were selected and sprayed with different concentrations of bio regulators as NAA (25, 50 and 75 ppm), GA₃ (50, 100 and 150 ppm) and Ethrel (250, 500 and 1000 ppm). NAA and GA₃ were sprayed at fruit set, whereas Ethrel was sprayed 20 to 25 days prior to harvesting to find out the optimum concentration of these bio regulators. Uniform sized mature fruits were harvested for carrying out physico-chemical analysis. A sample of ten randomly selected fruits was being taken from each treatment and their size was noted using the vernier's caliper. The weight of fruits from randomly drawn sample was computed with the help of electronic weighing balance. The fruits were evaluated for colour rating by a panel of five judges on score card (maximum 10 points) based on colour. Fruits were rated as follows:

Table 1: Score card for colour rating of phalsa fruits

Acceptability	Grading	Colour
Excellent	9.1-10	Purple black
Very good	8.1-9	Cherry red
Good	7.1-8	Light red
Fair	6.1-7	Greenish red
Poor	6 and below	Light green

The stones from the same sample were extracted and their size and weight were recorded with the similar procedures taken for fruits. The fruit TSS was analyzed with the help of hand refractometer. Titratable acidity was calculated by titrating diluted juice against N/10 NaOH by using phenolphthalein as an indicator. The diluted juice was prepared by adding distilled water in the ratio of 1:10. The end point of the titration was permanent pink colour. The acidity was calculated and expressed in terms of citric acid [6] by using the below given formula,

$$\text{Titrateable acidity (\%)} = \frac{0.064 \times \frac{N}{10} \text{NaOH}}{\text{Volume of juice taken}} \times 100$$

Reducing sugars and total sugars were analyzed by Lyan and Neyon method. Ascorbic acid were analysed by Indophenol dye method. The fruits were picked at regular intervals when ripe, the number of pickings was recorded and days taken to ripe from fruit setting were counted as harvesting span. The fruits were picked on alternate days and were weighed by

Comment [AL5]: for all measurement instruments add type and accuracy

Comment [AL6]: for both methods add references

57 using physical balance. Average yield per plant was worked out. The recorded data were computed by randomized block
58 design.

59 **3. RESULTS AND DISCUSSION**
60

61 **3.1 Fruit physical parameters**

62 The data presented in Table 2 revealed that most of the treatments improved the fruit size, however the plants treated
63 with GA₃ 150 ppm (T₉) yielded fruits with maximum length (1.26 cm) and breadth (1.43 cm). The former treatment was
64 also found to be at par with the treatment T₆ (Ethrel 1000 ppm) with fruit length and breadth of 1.18 cm and 1.42 cm
65 respectively.

66 **Table 2. Influence of various bio-regulators on fruit physical parameters of phalsa**

Comment [AL7]: add standard deviations

67

Treatments	Fruit Size		Fruit weight (g)	Fruit colour
	Fruit length	Fruit breadth		
	(cm)	(cm)		
T ₁ - NAA 25 ppm	0.91	1.13	0.98	7.60
T ₂ - NAA 50 ppm	1.00	1.22	1.04	7.80
T ₃ -NAA 75 ppm	1.08	1.28	1.07	7.90
T ₄ - Ethrel 250 ppm	1.19	1.33	1.06	8.80
T ₅ - Ethrel 500 ppm	1.09	1.33	1.08	8.90
T ₆ - Ethrel 1000 ppm	1.18	1.42	1.14	9.20
T ₇ - GA ₃ 50 ppm	1.23	1.35	1.13	8.40
T ₈ - GA ₃ 100 ppm	1.23	1.40	1.13	8.60
T ₉ - GA ₃ 150 ppm	1.26	1.42	1.20	8.70
T ₁₀ - Control	0.95	1.11	0.78	7.10
Mean	1.11	1.30	1.06	8.30
CD (P=.05)	.08	.06	.06	.70

Comment [AL8]: for what. What did you compare.

Comment [AL9]: for what

68

69 Fruit size significantly increased due to application of gibberellin this might be due to the fact that gibberellins indirectly
70 affect the level of auxin that ultimately caused cell elongation by enlargement of vacuoles and loosening of cell wall after
71 increasing its palatability. Also increase in fruit size by the use of ethrel could be attributed due to increase in the
72 availability of photosynthates and nutrients to the remaining fruits thereby increasing the size of individual fruits. Increase
73 in fruit size was also observed by [12] with ethrel and by [7] and [8] with gibberellins. GA₃ 150 ppm (T₉) treated plants
74 gave fruits with maximum weight of 1.20 g, which might be due to the fact that gibberellins increase the cell division and
75 translocation of food material [8]. Treatment T₉ was also found to be at par with the treatment T₆ (Ethrel 1000 ppm) with
76 fruit weight of 1.14 g. Results of these findings were confirmed by [4], [1] and [12]. The plants treated with treatment T₆
77 produced fruits with excellent purple black colour. This might be due to the fact that ethrel enhanced chlorophyll
78 degradation and increased carotene synthesis [9].

Comment [AL10]: where is the statistical proof?

79 3.2 Fruit biochemical parameters

80 The data regarding fruit biochemical parameters as influenced by various concentrations of ethrel, NAA and GA₃ were
81 presented in Table 3. Significant TSS (20.23 %) was observed in fruits obtained from plants treated with ethrel 1000 ppm.
82 This may be due to the quick metabolic transformation of starch and pectin into soluble compounds and rapid
83 translocation of sugars from the leaves to the developing fruits [10]. The results of these findings were also confirmed by
84 [4], [2] and [5].

Comment [AL11]: proof

85 **Table 3.** Influence of various bio-regulators on fruit bio-chemical parameters of phalsa
86

Treatments	TSS (%)	Titrateable acidity (%)	Reducing Sugars (%)	Total sugars (%)	Ascorbic acid (mg/100 g pulp)
T ₁ - NAA 25 ppm	15.23	2.97	5.38	9.78	22.60
T ₂ - NAA 50 ppm	15.81	2.27	5.49	10.52	23.86
T ₃ - NAA 75 ppm	15.75	2.29	5.54	10.47	23.91
T ₄ - Ethrel 250 ppm	16.07	2.22	6.02	10.69	25.03
T ₅ - Ethrel 500 ppm	18.48	1.80	6.77	12.75	25.28
T ₆ - Ethrel 1000 ppm	20.23	1.54	7.49	12.79	25.62
T ₇ - GA ₃ 50 ppm	17.13	2.22	6.69	12.35	30.06
T ₈ - GA ₃ 100 ppm	16.78	2.10	7.32	12.04	30.98
T ₉ - GA ₃ 150 ppm	18.09	1.97	7.28	12.62	31.10
T ₁₀ - Control	15.16	2.42	5.27	10.15	20.02
Mean	16.87	2.18	6.32	11.41	25.84
CD (P=.05)	.38	.52	.41	.82	.99

The fruits with minimum acidic content (1.54 %) were obtained from plants treated with ethrel 1000 ppm. This might be due to the fact that ethrel enhance the conversion of organic acids to sugars which increases the sugar content and decreases the acidity [11]. The present study is in conformity with the findings of [1] and [12].The data related to sugars showed that plants treated with Ethrel 1000 ppm yielded fruits with maximum reducing sugars (7.49 %) and total sugars (12.79 %). This might be due to that ethrel promoted hydrolysis of starch into sugars [5]. The present results are in conformity with the findings of [1] and [12]. Ascorbic acid content in fruits increased significantly by the use of gibberellins. Maximum ascorbic acid content (31.10 mg/100 g pulp) was observed in fruits obtained from plants treated withGA₃ 150 ppm. This might be due to the possible catalytic influence of gibberlic acid on biosynthesis of ascorbic acid from sugar or inhibition of oxidative enzymes or both [13]. The results of these findings are in conformity with [14] and [15].

3.3 Stone parameters

The data presented in Table 4 revealed that all the treatments showed non-significant effect on the stone size in phalsa. Minimum stone length (0.50 cm) and stone breadth (0.33 cm) was registered under the treatment T₁. From the data, it has been observed that NAA 25 ppm treated plants gave fruits with least stone weight (0.045 g). The present study is in conformity with the findings of [8].

Table 4. Influence of various bio-regulators on stone parameters of phalsa

Treatments	Stone length (cm)	Stone breadth (cm)	Stone weight (g)
T ₁ - NAA 25 ppm	0.50	0.33	0.04
T ₂ - NAA 50 ppm	0.56	0.40	0.07
T ₃ -NAA 75 ppm	0.53	0.40	0.06
T ₄ - Ethrel 250 ppm	0.53	0.40	0.11
T ₅ - Ethrel 500 ppm	0.56	0.43	0.08
T ₆ - Ethrel 1000 ppm	0.56	0.43	0.08
T ₇ - GA ₃ 50 ppm	0.53	0.41	0.10
T ₈ - GA ₃ 100 ppm	0.56	0.43	0.11
T ₉ - GA ₃ 150 ppm	0.56	0.46	0.09
T ₁₀ - Control	0.58	0.40	0.11
Mean	0.55	0.41	0.09
CD (P=.05)	NS	NS	0.02

3.4 Yield parameters

The presented data (Table 5) showed that ethrel significantly improved the yield parameters in phalsa. Ethrel 1000 ppm treated plants registered minimum harvesting span (22.81 days) than rest of the treatments. The number of pickings also significantly reduced by the use of ethrel.

Table 5. Influence of various bio-regulators on yield parameters of phalsa

Treatments	Harvesting span (days)	Number of pickings	Yield (kg/plant)
T ₁ - NAA 25 ppm	27.71	6.56	2.51
T ₂ - NAA 50 ppm	27.32	6.31	2.76
T ₃ - NAA 75 ppm	26.72	6.08	2.80

Comment [AL12]: proof

T ₄ - Ethrel 250 ppm	23.02	4.39	2.85
T ₅ - Ethrel 500 ppm	23.51	4.15	3.17
T ₆ - Ethrel 1000 ppm	22.81	3.90	3.36
T ₇ - GA ₃ 50 ppm	26.42	6.01	3.23
T ₈ - GA ₃ 100 ppm	25.13	5.57	3.47
T ₉ - GA ₃ 150 ppm	24.12	5.26	3.48
T ₁₀ - Control	30.52	7.13	1.95
Mean	25.72	5.53	2.95
CD (P=.05)	.07	.58	.15

Minimum number of pickings (3.90) was recorded in plants treated with ethrel 1000 ppm. This might be due to that ethrel hasten the maturation process [16]. These findings are in conformity with the findings of [1] and [5]. The plants treated with GA₃ 150 ppm registered maximum yield (3.48 kg/plant) and was followed by treatments T₈ and T₆ with fruit yield of 3.47 and 3.36 kg/plant respectively. All of these treatments were found to be at par with each other. Yield is a complex character and is characterized by increase in weight of individual fruit and number of fruits. It might be due to the fact that GA₃ increased the translocation and mobilization of photosynthates from source to sink [17]. The results are in accordance with the findings of [4], [14] and [18] in ber, [19] in Plum and [20] in mango.

4. CONCLUSION

The results of the present study concluded that plants treated with GA₃ 150 ppm and ethrel 1000 ppm registered significant improvement in fruit physical and biochemical parameters and also fruit yield per plant was increased. The fruits obtained from the plants treated with NAA 25 ppm had small sized and minimum weighed stone. Whereas the yield attributes were enhanced with the application of ethrel 1000 ppm in terms of advancement of harvesting and also reduction in number of pickings. Hence ethrel 1000 ppm may be applied to improve the fruit quality parameters as well as yield parameters and also to fetch the premium price in the market due to their attractive colour appearance than control.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Singh JP, Kumar S, Katiyar PN, Dwivedi AK. Effect of calcium nitrate, GA₃ and ethrel on fruiting, ripening and chemical traits of phalsa (*Grewia subinaequalis* D. C.). 2011; Annals Hort 4: 72-76.
- Kacha HL, Viradia RR, Leua HN, Jat G, Tank AK. Effects of NAA, GA₃ and ethrel on yield and quality of phalsa under south- surashtra condition. Asian J Hort. 2012; 7: 242-245.
- Pujari S. Phalsa cultivation in India – Production area, Climate, Harvesting and Fruit handling. 2015. <http://www.yourarticlelibrary.com/cultivation/phalsa-cultivation-in-india-production-area-climate-harvesting-and-fruit-handling/24703>.
- Debnath A, Vanajalatha, Umarfarooque M, Reddy M. Effect of NAA, GA₃, Kinetin and Ethrel on yield and quality in phalsa. Asian J Hort. 2011; 6: 474-477.
- Kacha HL, Jat G and Patel SK. Performance of various plant growth regulators on yield and quality of phalsa (*Grewia asiatica* L.). Hortflora Res Spec. 2014; 3:292-294.
- A O A C (1990) Official and tentative, methods of analysis. Association of official analytical chemist, 14th E.D. Washington D.C.

144 7. Shukla HS, Kumar V, Tripathi VK. Effect of gibberellic acid and boron on development and quality of aonla fruit
145 "Banarasi". ISHS Acta Hort. 2011; 890: 375.

146 8. Chandra R, Rawat M, Singh SS, Singh KK. Effect of foliar application of various growth regulators on yield and
147 quality of Aonla cv. Na-7 National Academy Agricultural Sci. 2015; 33:2123-2127.

148 9. Al-Mughrabi MA, Bacha MA, Abdelrahman AO. Influence of pre harvest application of ethrel and 2, 4-D on fruit
149 quality of Balady orange. Agri Sci. 1989; 1: 95-102.

150 10. Tripathi VK, Shukla PK. Effect of plant bio- regulators, zinc sulphate and boric acid on yield and quality of
151 strawberry cv. Chandler. J Asian Hort. 2007; 4: 15-18.

152 11. Singh P, Kumar S, Maji S. Effect of ethrel on post harvest changes in papaya (*Carica papaya* L.) fruits HortFlora
153 Res Spect. 2012; 1: 225-230.

154 12. Meitei SB, Patel RK, Deka BC, Deshmukh NA, Singh A. Effect of chemical thinning on yield and quality of peach
155 cv. Flordasun. African J Agricultural Res. 2013; 8:3558-3565.

156 13. Kher R, Bhat S, Wali VK. Effect of foliar application of GA₃, NAA and CCC on physico-chemical characteristics of
157 guava cv. Sardar. Haryana J Hort Sci. 2005; 34:31-32.

158 14. Tripathi D, Pandey AK, Pal AK, Yadav MP. Studies on effect of plant growth regulators on fruit drop,
159 development, quality and yield of ber cv. Banarasi Karaka. Progressive Horticulture. 2009; 41:184-186.

160 15. Wahdan MT, Habib SE, Bassal MA, Qaoud EM. Effect of some chemicals on growth, fruiting, yield and fruit
161 quality of "Succary Abiad" mango cv. J American Sci 2011; 7:651-658.

162 16. Tyler K, May D, Miller R. Ethrel sprays reduce number of pickings in hand- harvested cantaloupes. California
163 Agriculture. 1970; 24:6-7.

164 17. Krishnamoorthy HN. Gibberellins. Plant growth substances 1981; pp: 77-87.

165 18. Gill KS, JS Bal. Impact of application of growth regulators on Indian jujube. International Jujube Symposium,
166 2011; 993: 119-124.

167 19. Kaur H, Singh A, Gupta M, Randhawa JS. Effect of NAA and gibberellic acid on pre- harvest fruit drop and quality
168 of Satluj Purple plum. Haryana J Hort Sci. 2008; 37:31-32.

169 20. Rohit S. Effect of GA₃ and urea spray on flowering fruit retention and fruit quality of mango cv. Langra. Doctoral
170 dissertation, JNKKV. 2014.

171