

Original Research Article

Zinc and organic fertilization in maize augments enzymatic activities and nutrient bio-availability rendering higher production

Abstract

Low fertility and organic matter contents in soil limit the nutrient bio-availability for crops leading to poor nutrition in the animals and human beings. This field study on maize production employed integration of organic and chemical fertilizers to find out their impact on plant enzymatic activities in relation to crop production, and nutrient dynamics in plants and soil. The treatments compared were: control (no application of N from any source); 100% N from fertilizer (FN); 75% FN + 25% N from organic manure (ON); and 50% FN + 50% ON. Further, all these treatments were superimposed by three levels of zinc (Zn) fertilizer, viz., 0, 4, 8 kg Zn ha⁻¹, applied each one separately. Conjunctive use of FN & ON with the ratio of 75+25 along with Zn application @ 4 kg ha⁻¹ rendered the highest maize grain yield as compared to sole N application from chemical fertilizer. Similarly, enzymatic activities of nitrate reductase (NR) and superoxide dismutase (SOD) were significantly affected with combined use of N and Zn. Nitrogen, phosphorus and zinc contents in diagnostic leaves of maize were also higher in organically substituted treatments. So, integration of FN and ON was proved to render better nutrient supply to the plants throughout the growing season.

Keywords: Nutrient deficiencies; Chemical nitrogen fertilizer; Organic manure; Conjunctive use; Nitrate reductase; Superoxide dismutase.

22 1 Introduction

23 Pakistani soils exhibit multiple nutrient deficiencies including that of zinc (Zn) due to their alkaline-
 24 calcareous nature and low organic matter content, nutrient mining with intensive cropping, and
 25 inadequate and imbalanced fertilizer use. Declined soil organic matter in cultivated lands is a major cause
 26 of soil degradation; converting these soils infertile and thus not sustaining the crop production (Diacono
 27 and Montemurro 2010). Nutrient mining has also gained concerns about the sustainability of agricultural
 28 system. Continuous use of synthetic fertilizers for long time not only turns the land infertile and decreases
 29 crops yield (Sortino *et al.* 2013) but also contaminates the environment (Hu and Qi 2013). Achieving a
 30 balance between nutrient requirements of the plants and nutrient supplies from the soil is essential for
 31 attaining and maintaining high crop yields and good soil fertility. Manures enhance the bio-availability of
 32 macro- and micro-nutrients, viz., N, P, K and Zn, and thus concentration of these nutrients in plant tissues
 33 (Masti *et al.* 2003). Degraded lands can be ameliorated by conjunctive use of organic manures and
 34 inorganic fertilizers (Saleque *et al.* 2004). Appropriate blend of chemical fertilizers and organic manures
 35 varies with land-use pattern, agro-ecology, and socio-economic circumstances. Similarly, organo-mineral
 36 combined fertilizer blend could be a promising alternative for sustaining the crop yields (Deeks *et al.*
 37 2013). Improved management of crop residues / livestock manure and nutrients may also reduce the
 38 environmental impact of agriculture on climate change (Stavi and Lal 2013).

39 Nutrient deficiencies directly affect the enzymatic activities in plants. Zinc deficiency in crops is
 40 a common disorder in most of the cropped lands in Pakistan (Rafique *et al.* 2006). It is mainly due to
 41 calcareous-alkaline soils, low organic matter, nutrient depletion, and injudicious fertilizer application.
 42 Zinc is integral part of superoxide dismutase enzyme, which is important for detoxifying the reactive
 43 oxygen species (ROS) and controlling the free radicals (Cakmak 2000; Wang and Jin 2005) under
 44 drought stress. Similarly nitrate reductase is an important enzyme of N metabolism, its activity in plant is



inducible by nitrate (Shankar *et al.* 2001). Therefore, N deficiency affects the activity of nitrate reductase in plants. Under rain-fed conditions maize production is lower than in irrigated areas, which is mainly due to water stress and poor soil fertility status. Continuous additions of organic amendments to cropland increase soil fertility, organic carbon, microbial biomass and enzymatic activities, and enhance the crop yields (Diacono and Montemurro 2010). With these understandings, a field experiment on integrated nutrient management was conducted on rain-fed maize-crop.

This paper reports the maize crop response (yield, nutrients uptake and enzymes activity) to nitrogen and zinc fertilizers application along with organic manure. It was hypothesized that there will be differences in these crop attributes between organic-N fertilizer combinations and sole mineral N fertilizer treatments. Major objective of this study was to reduce the reliance on chemical fertilizers and enhance the use of organic amendments for sustainable soil health and crop production.



2 Materials and Methods

Field experiment was conducted on maize cv. Swan at the research farm of National Agricultural Research Centre (NARC), Islamabad (latitude 33° 43' N, longitude 73° 04' E, and altitude 490 m). Soil belongs to Nabipur soil series (coarse loamy, mixed, hyperthermic Udic Ustochrept). It is deep, well drained, and moderately calcareous developed on level to nearly level deposition of the flood plain. It lies under sub-humid to humid and medium to high rainfall-zone with annual rainfall ranging from 517 to 1550 mm. Mean maximum temperature during summer ranges from 36 to 42°C. Soil is alkaline (pH, 7.8), calcareous (CaCO₃ equiv., 4.3 %), low in organic matter (0.50 %), and deficient in NO₃-N (3.5 mg kg⁻¹), available P (3.0 mg kg⁻¹), with extractable K (80 mg kg⁻¹) and Zn (0.27 mg kg⁻¹).

The experiment comprised of two factor treatments, viz., nitrogen sources combinations and zinc fertilization. The N application was made in main plots, while Zn was applied in sub-plots. Experimental

design was split-plot under randomized complete block design (RCBD) with three replications. Plot size for main-plot treatments was 5 m × 9 m and each main-plot was divided into three equal sub-plots measuring 3 m × 5 m. Treatments detail is as follows:

Main plots - Nitrogen combinations

Control = No application of N from any source

FN100 = Whole N from chemical fertilizer (FN) 

FN75+ON25 = 75% FN + 25% N from organic manure (ON)

FN50+ON50 = 50% FN + 50% ON

Sub plots - Zinc levels

Zn0 = No application of zinc fertilizer

Zn4 = Zinc fertilizer application @ 4 kg ha⁻¹

Zn8 = Zinc fertilizer application @ 8 kg ha⁻¹

Nitrogen was applied @ 120 kg ha⁻¹ as recommended dose for the maize crop in rainfed areas using different combinations of organic (farm yard manure) and fertilizer N  (urea) sources. Fertilizer N (FN) was applied in two splits, viz., at seed bed preparation and at booting stage. Farm yard manure containing 1.85% N was applied before planting @ 3.3 t ha⁻¹ for replacing 50% recommended mineral N and 1.82 t ha⁻¹ for 25% replacement. Phosphorus fertilizer was equally applied at the rate of 50 kg ha⁻¹ in all the treatments at the time of sowing. 

2.1 Soil and plant analysis

Before sowing the crop, composite soil samples were collected from 0-15cm depth, air dried, ground and passed through 2 mm sieve. Soil samples were analyzed for pH (McLean 1982), AB-DTPA P (Soltanpour and Workman 1979), Electrical conductivity (ECe), (Rhoades 1982), Organic matter (Nelson and Sommer 1982) NH₄OAC extractable K (Knudsen *et al.* 1982) NO₃-N and Zn (Soltanpour and Workman 1979). At blooming stage  recently fully matured leaves from selected plants were collected.

The leaves were washed, cleaned, air dried and then oven dried at 65°C to a constant weight. The samples were ~~grinded~~^{ground}, sieved and stored in small plastic bags for determination of N, P and Zn. Nitrogen was determined using Kjeldahl method as described by Jackson (1973), P was determined by wet digestion as described by Benton *et al.* (1991). Zinc was determined using Atomic Absorption Spectrometry as described by Wright and Stuczynski (1996).

2.2 Measurement of enzymes activity

Samples of 2nd youngest fully expanded leaves at booting stage (diagnostic leaves) were taken, immediately frozen in liquid N, ground to make fine powder, and refrigerated at -80°C. Extraction of antioxidant enzymes in these leaves was performed through the procedure as described by Sankhla *et al.* (1985). Activity of superoxide dismutase (SOD) ~~enzyme~~ was assayed spectrophotometrically at 560 nm by measuring its ability to inhibit photochemical reduction of nitro blue tetrazolium (Dhindsa *et al.* 1981). For nitrate reductase activity, fresh leaf samples were weighed and shifted to plastic vials. In each vial, 2.5 mL phosphate buffer (pH 7.5), 0.2 M KNO₃ and 5% isopropanol solutions were added, and incubated for 2 h in dark at 30°C. After incubation, 1% sulphanilamide and 0.2% NED-HCl (N-1-naphthylethylene-diamine dihydrochloride) were added in to vials, and kept for 20 min for colour development. Absorbance in samples was measured spectrophotometrically at 540 nm as described by Jaworski (1971).

2.3 Statistical analysis

Data on growth, yield and nutrient uptake attributes of the crop were subjected to analysis of variance (Sokal and Rohlf 1997). Data showing significant difference at $P \leq 0.05$ were put to comparison of treatment means by least significant difference (LSD)  test. All the data were processed using MSTAT-C software for statistical analysis.

3. Results and Discussion

3.1 Maize grain yield

Pakistani soils exhibit multiple nutrient deficiencies including that of Zn (Rafique *et al.* 2006) due to their alkaline-calcareous nature and low organic matter content, nutrient mining with intensive cropping, and inadequate and imbalanced fertilizer use. Effect of experimental treatments on maize grain yield is given in Fig 1. Treatment with combination of fertilizer and organic N (FN75+ON25) produced significantly higher grain yield as compared to FN50+ON50 combination and sole chemical N source (FN100). It shows a synergistic effect of organic and inorganic N sources. Secondly, the results of the study also showed positive interaction of N sources and Zn fertilization on maize productivity. The yield increase might be due to enhanced hormonal and enzymatic activities with the application of Zn, and due to balanced and sustained supply of nutrients from organic manure. In case of 100 % chemical N, the nutrients availability might not be sufficient for longer period. (Yadav *et al.* 2006) reported that compared to their sole application, integrated use of N, FYM and Zn is the best choice in term of crop yield, nutrient uptake, and economic return and benefit cost ratio. Shaheen *et al.* (2010) also observed enhanced crop yield and nutrient uptake through synergistic and improved fertilization practices.

3.2 Nitrogen concentration in diagnostic leaf of maize

Nitrogen concentration in maize diagnostic leaf (third leaf below tassel) is given in Fig 2. Compared to that in control, N application significantly increased N concentration. Regarding the comparison of N sources, organic treatments viz. FN75+ON25 and FN50+ON50 showed higher N contents than with sole N application (FN100). It reflects that N was available for long period to plants from organically substituted treatments. It might be due to synergistic effect of organic and inorganic sources on mineralization and sustained supply of essential nutrients. In various studies on IPNM, it has been observed that integration of N sources viz. organic and inorganic origin increased NPK concentration in

alfalfa and maize (Lioveras *et al.* 2004; Sial *et al.* 2007). Under the combined application, decomposition process is enhanced by microbial activity, and energy is readily available from carbon for release of nutrients. Sial *et al.* (2007) also recorded higher N and P concentration in maize leaves and grain, when mineral N was applied in combination with organic sources.

Zinc fertilization also enhanced N concentration significantly in maize diagnostic leaf over control, but Zn application rates of 4 and 8 kg ha⁻¹ were statistically similar. Different physiological processes are inhibited in plants due to Zn deficiency. As a result plant growth and yield are severely affected. Zinc plays fundamental role in critical cellular functions like protein metabolism, gene expression, bio membranes integrity, and IAA metabolism (Marschner 1995). It has also been reported that zinc application increased significantly protein and nitrogen content of maize plant (Tahir *et al.* 2009; Adiloglu and Saglam 2005).

Organically substituted N treatments along with Zn4 or Zn8 showed higher N contents in the diagnostic leaf as compared to that in sole N treatments with zinc, viz., FN100+Zn4 and FN100+Zn8. Overall, there was positive interactive effect on N content in maize. Verma *et al.* (2005) also recorded positive interactive effect of N and Zn on the growth, yield, quality and nutrient uptake of fodder sorghum.

3.3 Phosphorus concentration in diagnostic leaf of maize

Effect of N and Zn treatments on P concentration in diagnostic leaf of maize are given in Fig 3. Nitrogen and Zn treatments significantly differ in P contents. The highest P content was recorded in FN75+ON25 treatment, which was statistically higher than that under FN50+ON50 and FN100. Similar to N, Zn treatments have also increased the P content in diagnostic leaf. Zinc application at 4 and 8 kg ha⁻¹ gave significantly higher P content than in control but statistically alike with each other. Akinrinde *et al.* (2006) also reported that Zn application increased the P contents in maize tissues.

Interaction of N and Zn treatments was highly significant with respect to P concentration in diagnostic leaf. The highest P contents were recorded in organically substituted combinations of N and Zn, viz., FN75+ON25+Zn4, FN75+ON25+Zn8, FN50+ON50+Zn4 and FN50+ON50+Zn8. These treatments were statistically superior to all others including that of 100% fertilizer N application. These findings support the phenomena of higher nutrient availability under combined use of organic and inorganic fertilizers. Balanced application of plant nutrients and IPNM enhance crop yields (Dong *et al.* 2006).

3.4 Zinc Concentration in diagnostic leaf of Maize

Influence of N and Zn application on Zn content in diagnostic leaf is reflected in Fig 4. Zinc contents in diagnostic leaf were significantly increased with N and Zn application. The treatment containing 50% N gave significantly higher Zn content than with FN75+ON25 and FN100, which were statistically alike. Similarly, Zn application also affected Zn content of plant tissues compared to that in control. All the three levels of Zn application differed statistically ~~with each other~~. Maximum contents were observed in Zn8 level, which were significantly higher than in Zn4 level. Interaction of N and Zn was also found significant ~~that might be due~~ to organic sources of N. The highest Zn concentration was observed in FN50+ON50+Zn8 followed by FN50+ON50+Zn4, which are statistically similar but differed with all other combinations of Zn and N fertilization. Other researchers also reported that zinc content and uptake by maize increased with Zn application ~~as it attributed~~ to better root growth (Obrador *et al.* 2002). Concentrations of N and Zn were also increased in maize plant with FYM application (Gondek and Mazur 2005). Akinrinde *et al.* (2006) also found that application of cow manure + ZnSO₄ produced the highest plant shoot biomass and gave the highest Zn uptake by maize.

Activity of superoxide dismutase

Superoxide dismutase (SOD) is an antioxidant enzyme, which counters stress in plants. Zinc is the structural part of SOD, its activity was determined in leaves at booting stage to assess the effect of Zn application on SOD. Data regarding the SOD activity is mentioned in Fig. 5. The effect of N application on SOD activity was significant over control. However, N sources were statistically at par w.r.t. SOD activity. The highest SOD activity was recorded in N₄ followed by N₃ and sole application of N (FN100). The result is in line with the finding of (Alguacil *et al.* 2006), who reported that compost addition significantly enhanced the SOD and POD activities in the shoots of *J. oxycedrus*. Zinc application at 8 kg Zn ha⁻¹ gave significantly higher activity than that with 4 kg Zn ha⁻¹ and control. Interaction of Zn and N treatments was also significantly positive. The highest SOD activity was recorded in FN50+ON50+Zn8 followed by FN75+ON25+Zn8, and FN75+ON25+Zn4 treatments, being statistically similar.

Organic manure treatments with Zn significantly increased the SOD activity. It might be due to structural role of Zn in SOD enzyme. So, it has positive effect on the activity of SOD. Activity of SOD in wheat leaves significantly increased in NO₃ and NH₄ treated plants as compared to N deficient wheat plants (Polesskaya *et al.* 2004). Zn is a cofactor of 300 enzyme and proteins, and it is involved in cell division, nucleic acid metabolism and protein synthesis. Zinc application enhances the activities of antioxidant enzymes (Cakmak 2000). Wang and Jin (2007) also reported that Zn application raised the activity of SOD in maize leaves. Zinc enhances the SOD activity and reduces MDA contents, which is beneficial to counter the free radicals (Gao *et al.* 2001).

Nitrate reductase activity



Nitrate reductase is an important enzyme of N metabolism, its activity in plants is inducible by its substrate (NO₃). So, anything that affects nitrate availability will influence nitrate reductase activity (Pervaiz *et al.* 2002). Nitrogen application significantly enhanced nitrate reductase activity (NRA) over control even N sources significantly differ in NRA with each other (Fig 6). The highest NRA was recorded in FN50+ON50, where N was applied in equal ratio from organic and inorganic sources. Similar

to N application, Zn also significantly affected the NRA. Zinc application at 8 kg ha⁻¹ showed the highest NRA, which was significantly higher than with 4 kg Zn ha⁻¹. Zinc fertilization showed a positive effect on NRA. Maximum NRA was observed in manure supplemented treatments with Zn application.

Soil microbial biomass and enzymatic activities increase with integrated nutrient management (INM), which ultimately activates the micronutrients bio-availability (Tejada *et al.* 2008). Interactive effect of N sources and Zn was also significant in both crops. The highest NRA was recorded in FN50+ON50+Zn8 followed by FN50+ON50+Zn4 and FN75+ON25+Zn8 treatments. Alguacil *et al.* (2006) also found that under drought conditions, only inoculation with exotic arbuscular mycorrhizae fungi increased the shoot and root NRA by 188 and 38%, respectively as compared to the plants without inoculation or compost application. Nardi *et al.* (2004) also reported that NRA correlated positively with organic N, and humus extract of FYM treated soil contained 42% higher NRA as compared to control. Soil microbial biomass also increases enzymatic activities in soil.

Literature Cited

- Adiloglu S, Saglam MT (2005) The effect of increasing nitrogen doses on the zinc content of maize plant in the soils of different properties. Pak J Biol Sci 8:905-909
- Akinrinde EA, Olubakin OA, Omotoso SO, Ahmed AA (2006) Influence of zinc fertilizer, poultry manure and application levels on the performance of sweet corn. Agri J 1:96-103
- Alguacil M, Caravaca F, Diaz-Vivancos P, Hernandez JA, Roldan A (2006) Effect of arbuscular mycorrhizae and induced drought stress on antioxidant enzyme and nitrate reductase activities in *Juniperus oxycedrus* L. grown in a composted sewage sludge-amended semi-arid soil. Plant Soil 279:209-218

- 226 Benton J, Benjamin W, Harry AM (1991) Analysis Hand Books: A practical sampling, preparation and
227 interpretation guide. Micro-macro Publishing, Inc. pp 130
- 228 Cakmak I (2000) Possible role of zinc in protecting plant cells from damage by reactive oxygen species.
229 New Phytologist 146:185-205
- 230 Deeks LK, Chaney K, Murray C, Sakrabani R, Gedara S, Le MS, Tyrrel S, Pawlett M, Read R, Smith GH
231 (2013) A new sludge-derived organo-mineral fertilizer gives similar crop yields as conventional
232 fertilizers. Agron Sustain Dev. doi: 10.1007/s13593-013-0135-z
- 233 Dhindsa RS, Plumb-Dhindsa P, Thorpe TA 1981 Leaf senescence: correlated with increased levels of
234 membrane permeability and lipid peroxidation and decreased levels of superoxide dismutase and
235 catalase. Exp Bot 32:93-101
- 236 Diacono M, Montemurro F (2010) Long-term effects of organic amendments on soil fertility. A review.
237 Agron Sustain Dev 30, 401-422
- 238 Dong J, Hengsdijk H, Ting-Bo DAI, De Boer W, Jing Q, Wei-Xing CAO (2006) Long-term effects of
239 manure and inorganic fertilizers on yield and soil fertility for a winter wheat-maize system in
240 Jiangsu, China. Pedosphere 16:25-32
- 241 Gao Z, Lin B, Zhou W (2001) Effect of zinc nutrition on endogenous plant hormones and free radicals in
242 spring maize. Plant Nutr Fert Sci 1:424-428
- 243 Gondek K, Mazur BF (2005) The effect of mineral treatment and the amendments by organic and
244 organomineral fertilizers on crop yield, plant nutrient status and soil properties. Plant Soil
245 Environ 51:34-45

- 246 Hu C, Qi Y (2013) Effective microorganisms and compost favor nematodes in wheat crops. Agron
247 Sustain Dev. Doi: 10.1007/s13593-012-0130-9
- 248 Jackson ML (1973) Soil Chemical Analysis. Prentice Hall of India Limited, New Delhi
- 249 Jaworski EG (1971) Nitrate reductase assay in intact plant tissues. Biochem Biophys Res Commun 43:
250 1274-1279
- 251 Knudsen D, Peterson GA, Pratt PF (1982) Lithium, sodium and potassium. In: Page AL, Miller RH,
252 Keeney DR (eds) Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties, 2nd
253 Ed. Soil Sci Soc Amer, Madison, WI, pp 115-246
- 254 Lioveras J, Aran M, Villar P, Ballesta A, Arcaya A, Vilanova X, Delgado I, Munoz F (2004) Effects of
255 swine slurry on Alfalfa production and on tissue and soil nutrient concentration. Agron J 96:986-
256 991
- 257 Marschner H (1995) Mineral Nutrition of Higher Plants, (2nd Ed). Academic Press, London. pp 889
- 258 Masti T, Lithourgidis AS, Gagianas AA (2003) Effect over of injected liquid cattle manure on growth and
259 yield of winter wheat and soil characteristics. Agron J 95:592-599
- 260 McLean EO (1982) Soil pH and lime requirement. In: Page AL, Miller RH, Keeney DR (eds) Methods of
261 Soil Analysis. Part 2 - Chemical and Microbiological Properties (2nd Ed). Agronomy 9:199-223
- 262 Nardi S, Morari F, Berti A, Tosoni M, Giardini L (2004) Soil organic matter properties after 40 years of
263 different use of organic and mineral fertilizers. Europ J Agron 21:357-367
- 264 Nelson DW, Sommer LE (1982) Total carbon, organic carbon, and organic matter. In: Page AL (ed)

- 265 Methods of Soil Analysis. 2nd Ed. Amer. Soc. Agron. Madison, WI. pp 539-579
- 266 Obrador A, Alveraz JM, Fernandez MD, Lopez-valdivia LM (2002) Changes with time of zinc forms in a
- 267 acid, a neutral and a calcareous soil amended with three organic zinc complexes. Aust J Soil Res
- 268 40:139-148
- 269 Pervaiz Z, Afzal M, Xi X, Xiaoe Y, Luo (2002) Relationship among nitrate reductase activity, growth,
- 270 mineral contents and salt tolerance of wheat cultivars under saline condition. Pak J Soil Sci
- 271 21:67-72
- 272 Polesskaya OG, Kashirina EI, Alekhina ND (2004) Changes in the activity of antioxidant enzymes in
- 273 wheat leaves and roots as a function of nitrogen source and supply. Russ J Plant Physiol 51:615-
- 274 620
- 275 Rafique E, Rashid A, Ryan J, Bhatti AU (2006) Zinc deficiency in rainfed wheat in Pakistan: Magnitude,
- 276 spatial variability, management, and plant analysis diagnostic norms. Commun Soil Sci Plant
- 277 Anal 37:181-197
- 278 Rhoades JD (1982) Soluble salts. In: Page A. et al. (ed) Methods of Soil Analysis. Part 2. Agronomy
- 279 9:167-178
- 280 Saleque MA, Abedin MJ, Bhuijan NI, Zaman SK, Panaullah GM (2004) Long term effect of inorganic
- 281 and organic fertilizer sources on yield and nutrient accumulating of low land rice. Field Crops
- 282 Res 86:53-65
- 283 Shaheen A, Naeem MA, Jilani G, Shafiq M (2010) Integrated soil management in eroded land augments
- 284 the crop yield and water use efficiency. Acta Agri Scand B - Plant Soil Sci 60(3):274-282

- 285 Shankar N, Khan SR, Srivastava HS (2001) The response of nitrate reductase activity and nitrate
286 assimilation in the maize roots to growth regulators at acidic pH. *Biologia Plant* 44:599-601
- 287 Sial RA, Chaudhary EH, Hussain S, Naveed M (2007) Effect of organic manures and chemical fertilizers
288 on grain yield of maize in rain- fed area. *Soil Environ* 26:130-133
- 289 Soltanpour PN, Workman S (1979) Modification of the NH_4HCO_3 DTPA soil test to omit carbon black.
290 *Commun Soil Sci Plant Anal* 10:1411-1420
- 291 Sokal RR, Rohlf FJ (1997) *Biometry: The Principles and Practice of Statistic in Biological Research*. WH
292 Freeman, New York
- 293 Sortino O, Dipasquale M, Montoneri E, Tomasso L, Avetta P, Prevot AB (2013) 90 % yield increase of
294 red pepper with unexpectedly low doses of compost soluble substances. *Agron Sustain Dev*
295 33:433-441
- 296 Stavi I, Lal R (2013) Agriculture and greenhouse gases, a common tragedy. A review. *Agron Sustain Dev*
297 33:275-289
- 298 Tahir M, Fiaz N, Nadeem MA, Khalid F, Ali M (2009) Effect of different chelated zinc sources on the
299 growth and yield of maize (*Zea-mays* L.). *Soil Environ* 28:179-183
- 300 Tejada M, Gonzalez JL, Garcia-Martinez AM, Parrado J (2008) Application of green manure and green
301 manure composted with beet vinasse on soil restoration: Effect on soil properties. *Bioresour*
302 *Technol* 99:4949-4957
- 303 Verma S, Subehia SK, Sharma SP (2005) Phosphorus fractions in an acid soil continuously fertilized with
304 mineral and organic fertilizers. *Biol Fertil Soils* 41:295-300

305 Wang H, Jin JY (2005) Photosynthetic rate, chlorophyll florescence parameters, and lipid peroxidation of
306 maize leaves as affected by zinc deficiency. *Photosynthetica* 43:591-596

307 Wright RJ, Stuczynski T (1996) Atomic absorption and flame emission spectrometry. In: Sparks DL (ed.)
308 *Methods of Soil Analysis, Part 3. Chemical Methods*. SSSA. pp. 65-90

309 Yadav RP, Singh P, Aggarwal RK, Yadav MK (2006) Improved technology for sustainable production of
310 maize (*Zea mays* L.) in Shivalik foothills of Himachal Pradesh. *Indian J Dryland Agri Res Dev*
311 21:1-6

312

Figure Captions

Fig. 1. Effect of N sources and Zn levels on maize grain yield

Fig. 2. Effect of N sources and Zn levels on nitrogen concentration in maize leaf

Fig. 3. Effect of N sources and Zn levels on phosphorus concentration in maize leaf

Fig. 4. Effect of N sources and Zn levels on zinc concentration in maize leaf

Fig. 5. Effect of N sources and Zn levels on superoxide dismutase activity in maize leaf

Fig. 6. Effect of N sources and Zn levels on nitrate reductase activity in maize leaf





Fig. 1. Effect of N sources and Zn levels on maize grain yield

325




326

327 **Fig. 2.** Effect of N sources and Zn levels on nitrogen concentration in maize leaf

328

329




330

331 **Fig. 3.** Effect of N sources and Zn levels on phosphorus concentration in maize leaf

332

333




334

335 **Fig. 4.** Effect of N sources and Zn levels on zinc concentration in maize leaf

336

337




338

339 **Fig. 5.** Effect of N sources and Zn levels on superoxide dismutase activity in maize leaf

340



341



342

343 **Fig. 6.** Effect of N sources and Zn levels on nitrate reductase activity in maize leaf

344

345

346