

Comparative studies on some of the bioactive components of fluted pumpkin grown in three selected solid media

Abstract:

The amino acids, vitamins and proximate composition of pumpkin grown in three different selected solid media were assessed and compared. The solid media were humus soil, white-sand and wood sawdust. Standard procedure and equipment were followed and used, respectively for the determination of the bioactive components of pumpkin. Proximate composition of pumpkin in the three media were: carbohydrate (12.76%, 20.40% and 6.37%), crude fibre (4.76%, 5.70% and 5.59%), ash content (0.47%, 0.58% and 0.73%), crude lipid (0.30%, 1.00% and 0.10%) and moisture content (81.41%, 72.02% and 86.91%) for humus soil, white-sand and sawdust, respectively while the crude protein was constant at 0.30%. In that same order, the total amino acids were 9.75 g/100g, 7.53 g/100g and 13.46%. The essential amino acids (5.85%, 4.49% and 7.94%) and non-essential amino acids (3.90 g/100g, 3.04 g/100g and 5.52 g/100g) varied for humus soil, white-sand and sawdust, respectively. Among the amino acids, histidine (1.73 g/100g, 1.34 g/100g, and 2.45 g/100g), cysteine (1.26 g/100g, 0.98 g/100g and 1.78 g/100g) and threonine (1.23 g/100g, 0.96 g/100g and 1.75 g/100g) were the most abundant while proline (0.01 g/100g) was the least for humus soil, white-sand and sawdust as a medium of growth for pumpkin. The percentage water soluble vitamins (B₁, B₂, B₃, B₆, B₁₂ and C): 26.16%, 71.42% and 135.53% and fat-soluble vitamins (A, E and K): 18.87%, 19.95% and 41.73% for humus soil, white-sand and sawdust, respectively. The water-soluble vitamins accounts for the high total vitamins obtained. The study has shown that pumpkin can be grown in white-sand and sawdust without losing the bioactive compounds rather improving its availability.

Keywords: Bioactive compounds, fluted pumpkin, solid media

Introduction:

A well balanced diet of natural foods generally provide the vitamins requirements of man. In contrast to the inorganic mineral elements, they are essential dietary constituents being required for growth, health and reproduction. Vegetables are highly perishable and require careful processing in order to preserve the nutrients, especially the water soluble vitamins. Ascorbic acid or vitamin C is acquired human nutrient. Many nations of the world depend on vegetables as their main sources of several nutrients and medicine especially the less developed and developing nations. Nigeria and most African countries consumed green leafy vegetable as diet or component of their meal in most cases. According to Hossain *et al.* [1], leafy vegetables are widely consumed in many countries because of their nutritional quality and anthocyanin, ascorbic acid, β -carotene, flavonoids, folic acid, polyphenol and alkaloid contents. These components were earlier reported to have a wide range of biological functions such as antiallergic, anticancer, antidiabetic, antimicrobial, antioxidant, anticardiovascular diseases [2]. Among the 20 amino acids required by humans for protein, only twelve can be produced within the body, whereas the other eight, isoleucine, leucine,

lysine, methionine, phenylalanine, threonine, tryptophan and valine must be obtained from the diet. In addition, adults are capable of synthesizing arginine and histidine but these amino acids are believed to be essential to growing children, meaning that children cannot produce them on their own and must be obtained from other external supplies [3]. The amino acids (Asp, Glu, Ser, Gly, Ala, and Leu) are naturally involved in osmolyte synthesis, cell metabolism, ammonia detoxification, antioxidant activity and alkaloid synthesis [4]. It is interesting to know that many depend on vegetables intake in their diet instead of inorganic supplement for health concerns. In fact, food not only provides vital nutrients which are essential for life, but also other bioactive compounds which are necessary to promote health and prevent disease [5].

Fluted pumpkin (*Telfairia occidentalis* Hook F.) is a perennial angiosperm plant with great economic importance in Nigeria. This plant is indigenous to southern Nigeria, and used primarily in soups and herbal medicine [6]. The tender leaves and vines are consumed by humans as vegetables as well as livestock as forage, while the young seeds are eaten as food and snacks [6,7,8]. The cooked leaves are staple vegetable in soups and stews of various cultures throughout equatorial Africa [9,10]. In Nigeria, several works on *T. occidentalis* leaf as a leafy vegetable in the dietary food or as an infusion in medicinal preparation is being encouraged as a result of its medicinal properties which includes antidiabetic, antianemic, hepatoprotective effect, and as a purgative leafy vegetable [11,12,13,14].

Considering the fact that intake of green leafy vegetables in most regions of the world is on the increase either for food or medicine, the study seek to identify and quantify some of the bioactive compounds in *Telfairia occidentalis* leaf grown in three selected solid media.

Materials and methods:

The seeds of *Telfairia occidentalis* and sawdust (SD) used were sourced from Choba Market and Rumuosi Sawmill respectively while the white sand (WS) and humus soil (HS) were obtained from Choba River and a garden in University of Port Harcourt in that order. The seeds were planted in three different media namely; white sand, sawdust and humus soil. After germination, the seedlings were allowed to stand for one month when it is matured and can be harvested to prepare food. The leafy vegetables were harvested and the following analyses were carried out: proximate composition, amino acids and vitamins.

Proximate Analysis:

Proximate analysis (moisture, ash, protein, carbohydrate and lipid content) was determined using standard method of AOAC [15]. Fibre was determined by difference.

Determination of crude protein (Kjeldahl method): Plant samples weighing 0.1 g were put into different conical flasks, 3 g of digestion catalyst and 20 ml of concentrated sulphuric acid was added into the flask. The flasks were then heated gently to boil in a fume chamber, until charred particles disappeared and clear greenish grey solution was obtained. The resulting solution in the conical flask was heated for an additional 20 minutes and allowed to cool. The digest was diluted with water to 100 ml capacity and 20 ml was then measured into a distillation flask and this was connected to a condenser adapted to a receiver beaker

containing 10 ml of 2% boric acid with 2 drops of double indicator. NaOH (40%) was added to the digest, the distillation flask was then heated to distil the nitrogen present as ammonia. The boric acid in the receiver was titrated with standard 0.1N hydrochloric acid. The volume of HCl used was recorded as titre value. The formula for the calculation of total Nitrogen is:

$$\% \text{Nitrogen} = (\text{Titre value} \times 1.4 \times 100) / (1000 \times 20 \times 0.1)$$

Titre value = Volume of Hydrochloric acid used.

1.4 = Nitrogen equivalent in relation to the normality of HCl used in titration.

100 = Percentage factor

1000 = Conversion factor from gram to milligram.

20 = Integral volume of digest analyzed or distilled

0.1 = Weight of sample in gram digested

% Crude protein = %Nitrogen X 6.25

Determination of carbohydrate (Clegg Anthrone method): Plant samples weighing 1 g were put into a 250 ml volumetric flask. Distilled water (10 ml) and 13 ml of 62% perchloric acid was added and the mixture was shaken in order for it to homogenize completely. The flask was made up to 250 ml with distilled water; the solution formed was filtered through a glass filter paper. Filtrate (10 ml) was collected and transferred into a 100 ml test tube; this was also diluted to volume with distilled water. The hydrolyzed solution was pipetted into a clean test tube and 5 ml of Anthrone reagent was added, they were then mixed together. The whole mixture was read at 630 nm wavelength using the 1 ml distilled water and the 5 ml anthrone prepared as blank. Glucose solution of 0.1 ml was also prepared and this was treated with the anthrone reagent. Absorbance of the standard glucose was calculated using the formula below:

$$\% \text{CHO} = (25 \times \text{Absorbance of sample}) / (\text{Absorbance of standard glucose} \times 1 \text{g of sample used})$$

Determination of moisture using the air oven method: Samples weighing 1 g were placed in a clean dry porcelain evaporation dish. This was placed in an oven to maintain a temperature of 105°C for six hours. The evaporating dish was cooled to room temperature in a desiccator and re-weighed.

$$\% \text{Moisture} = [(\text{weight of fresh sample} - \text{weight of dried sample}) / \text{Weight of sample used}] \times 100$$

Determination of lipid by soxhlet extraction method: Samples weighing 2 g was inserted into a filter paper and placed into the soxhlet extractor. The extractor was fitted into a pre-weighed round bottomed dry distillation flask and the solvent acetone was added into it through the condenser. The extractor and the flask were held in place with a retort stand clamp. Cold water was passed into the condenser via the rubber tubing and the flask was heated such that the solvent refluxed continuously within the enclosure. The lipid in the solvent chamber was extracted through this process of continuous refluxing. After the lipid had been extracted completely from the sample, the condenser and the extractor were disconnected, the acetone solvent was distilled off and the lipid concentrate was cooled in the oven and re-weighed.

126 % Lipid = [(Weight of flask and extract – Weight of empty flask) / Weight of sample
127 extracted] X 100

128 **Determination of ash by furnace method:** The dried sample weighing 1 g was placed into a
129 porcelain crucible which had been preheated and weighed. The crucible was inserted into a
130 muffle furnace and regulated to a temperature 630°C. After three hours it was removed from
131 the furnace and allowed to cool to room temperature, and then it was re-weighed.

132 **Determination of fibre:** The fibre content was determined by difference. The other five
133 proximate components were summed and the value gotten was subtracted from 100% giving
134 the fibre content (100 - per cent estimated proximate components represented the per cent
135 fibre in the sample).

136

137 **Determination of Amino acids using waters 616/626 LC (HPLC) Instrument:**

138 The sample preparation and determination were carried out in the following four stages:

139 **Hydrolysis:** The samples (0.5 g) were weighed into a sterile furnaces hydrolysis tube 5 nmols
140 Norleucine was added to the samples and then dried under a vacuum. The tube was again
141 placed in a vial containing 10.05 N HCl with a small quantity of phenol, thereby hydrolyzing
142 the protein by the HCl vapours under vacuum. This stage of hydrolysis of the sample lasted
143 for between 20 – 23 hours at 108°C. After the hydrolysis, the samples were dissolved in ultra-
144 pure water grade, containing ethylene diamine tetraacetic acid (EDTA). The EDTA chelates
145 the metal was present in the samples. The hydrolysed samples now are stored in HPLC amino
146 acid analyzer bottles for further analytical operations.

147 **Derivatisation:** The hydrolysed samples were derivatised automatically on the water 616/626
148 HPLC by reacting the five amino acid, under basic situations with phenylisothiocyanate
149 (PITC) to get phenylthiocarbamyl (PTC) amino acid derivatives. The duration for this is 45
150 minutes per sample, as calibrated on the instrument. A set of standard solutions of the amino
151 acids were prepared from Pierce Reference standards H (1000 µmol) into auto-sampler crops
152 and they were also derivatised. These standards (0.0, 0.5, 1.0, 1.5, 2.0 µmol) were used to
153 generate a calibration file that was used to determine the amino acids contents of the samples.
154 After the derivatisation, a methanol solution (1.5N) containing the PTC-amino acids was
155 transferred to a narrow bore waters 616/626 HPLC system for separation.

156 **HPLC separation and Quantization:** The separation and quantization of the PTC-amino
157 acids were done on a reverse phase (18 silica column) and the PTC chromophore were
158 automatically and digitally detected at the wavelength of 254 nm. The elution of the whole
159 amino acids in the samples took 30 minutes. The buffer system used for separation was 140
160 mm sodium acetate pH 5.50 as buffer A and 80% acetonitrile as buffer B. The program was
161 ran using a gradient of buffer A and buffer B concentration and ending with a 55% buffer B
162 concentration at the end of the gradient.

163 **Data processing / interpretation and calculation:** The intensity of the chromatographic
164 peaks areas were automatically and digitally identified and quantified using a Dionex
165 chromeleon data analysis system which is attached to the waters 616/626 HPLC System. The
166 calibration curve or file prepared from the average values of the retention times (in minutes)
167 and areas (in Au) at the amino acids in 5 standards runs was used.

Since a known amount of each amino acid in the standard loaded into the HPLC, a response factor (Au/pmol) was calculated by the software that was interphased with the HPLC. This response factor was used to calculate the amount of each of the amino acid (in pmols) in the sample and displayed on the system digitally. The amount of each amino acid in the sample is finally calculated by the software by dividing the intensity of the peak area of each (corrected for the differing molar absorptivity's of the various amino acids) by the internal standard (Pierce) in the chromatogram and multiplying this by the total amount of internal standard added to the original sample.

After the picomole by the intensity of the height of each amino acid has been ascertained by the software, the data, the digital chromatographic software extrapolate back to 5 nmoles of the internal standard (Norleucine), and displays for the total amount that was pipetted into the hydrolysis tube at the beginning of the analysis as below:

Calculation:

mg/ml (in Extract) = Dilution factor x Peakheight intensity

mg/ml (in sample) = $\frac{\mu\text{g/ml in extract} \times \text{sample volume}}{\text{Wt. of sample}}$

Wt. of sample

Vitamins:

Vitamin C was determined using titrimetric method [16].

Vitamin A extraction and determination using waters 616/626 HPLC

Plant sample (0.5 g) was weighed into a cornical flask, 20 ml of 0.2N HCl dispensed and allowed to stand for 1.5 hours. The solution was cooled and the pH adjusted to pH 6, using NaOH. Also, 1N HCl added to lower the pH to 4.5. The solution was made up to 50 ml, shake and centrifuged for 10 minutes at 3000 rpm. The supernatant was separated, 1 ml of acetic acid (CH₃COOH) added and mixed properly. Also, 0.5 ml of 3% H₂O₂ added and mixed well. Finally, 20 mg of sodium hydrogen sulphate added and shake properly. The extract was ran on HPLC (Waters 616/626). Water 616/626 accessories used had Merck Lichrosphere WOH-18/2 (5 μm) at 40°C column (stationary phase) and mobile phase (Solvent 'A' was 30 mM sodium acetate, pH 6.5 containing 5% dimethyl flormamide and solvent 'B' was acetonitrile) with fluorescence detector, range of working standard (0, 2, 4, 6, 8 ppm) and determination was carried out wavelength of 328 nm.

2.7.2 Vitamin B₂, B₆, B₁₂ combined extraction and determination using Waters 616/626 HPLC

Plant sample (2.5 g) was weighed into a set of digestion tubes, and an extraction solution (Ultra-pure water : HCl : 0.1N H₂SO₄, in the ratio 5:2:3) dispensed. The tube was warmed at the temperature of 40% for 2 hours, allowed to cool to room temperature and transferred to a set of plastic centrifuged tubes. The later was shake for 10 minutes and centrifuged at 3000 rpm. The supernatant was set in autoanalyser tubes and ran on HPLC. Water 616/626 accessories used had Merck Lichrosphere WOH-18/2 (5 μm) at 40°C column (stationary phase) and mobile phase (Solvent 'A' was 30 mM sodium acetate, pH 6.5 containing 5% dimethyl flormamide and solvent 'B' was acetonitrile) with fluorescence detector, range of

working standard (0, 0.2, 0.4, 0.6, 0.8 ppm) and determination was carried out wavelength range of 240 - 465 nm.

2.7.3 Vitamin E extraction and determination using Waters 616/626 HPLC

Plant sample (0.5 g) each was weighed into a set of digestion tubes, 20 ml of diluted hydrochloric acid (HCl) added and shake vigorously for 2 hours. The extract was further treated with phosphatase to liberate free vitamin E into the solution. The extract was purified by passing through base exchange silicate alkaline column to remove interfering compound. Thereafter, the extract was stored in a set of vials for analysis using HPLC. Water 616/626 accessories used had Merck Lichrosphere W0CH-18/2 (5 µm) at 40°C column (stationary phase) and mobile phase (Solvent 'A' was 30 mM sodium acetate, pH 6.5 containing 5% dimethyl flormamide and solvent 'B' was acetonitrile) with fluorescence detector, range of working standard (0, 0.2, 0.4, 0.6, 0.8 ppm) and determination was carried out wavelength of 356 nm.

Results and Discussion:

Proximate composition of the leaf of *T. occidentalis* grown in humus soil, white-sand and sawdust are presented in Table I. The results varied in the three growth media. *Telfairia occidentalis* leaf showed appreciable levels of crude protein (0.30%, 0.30% and 0.30%), carbohydrate (12.76%, 20.40% and 6.37%), crude lipid (0.30%, 1.00% and 0.10%), crude fibre (4.76%, 5.70% and 5.59%), ash (0.47%, 0.58% and 0.73%), and moisture content (81.41%, 72.02% and 86.91%) respectively. The moisture contents of *T. occidentalis* were high in the three media, which directly affects the shelf life. This has serious economic implications. According to Abdullahi [17], high moisture content is linked with increased microbial activities during storage. Interestingly in white-sand growth medium, *T. occidentalis* with the least moisture content had the highest carbohydrate, crude fibre and crude lipid. The particle sizes of the growth media probably affected the availability of water to the plant. Crude lipid is surprisingly low, ranging from 0.10 – 1.00%. This values is low compared to values (1.83 - 27%) reported by other researchers in some leafy vegetables consumed in Nigeria and Niger Republic [18,19]. These low crude lipids is of great importance, considering that excess intake of lipid leads to health challenges such as obesity and heart related problems.

Table I: Proximate composition of *T. occidentalis* grown in different media

Proximate composition (%)	Growth medium		
	Humus soil	White Sand	Sawdust
Crude protein	0.30	0.30	0.30
Carbohydrate	12.76	20.40	6.37
Crude lipid	0.30	1.00	0.10
Crude fibre	4.76	5.70	5.59
Ash	0.47	0.58	0.73
Moisture content	81.41	72.02	86.91

Amino acids content of *T. occidentalis* nurtured in three different solid medium are presented in Table II. The presence of essential (9) and non-essential (11) amino acids were observed. These amino acids were below the limit documented by FAO [20], indicating that *T. occidentalis* can possibly as one of the natural sources of amino acids for human consumption. Amongst the solid media, *T. occidentalis* grown in sawdust had the highest essential and non-essential amino acids with histidine (2.45 g/100g) and cysteine (1.78 g/100g) for each respectively. Also, histidine (1.73 g/100g; 1.34 g/100g) and cysteine (1.26 g/100g; 0.98 g/100g) were the most concentrated essential and non-essential amino acids in humus soil and white sand respectively. The essential amino acids content of *T. occidentalis* in three solid media with their values ranged from 0.04 – 2.45 g/100g, the least concentrated amino was tryptophan with values of 0.05, 0.04 and 0.08 g/100g in that order which is comparably lower than the value (0.43 g/100g) reported for *Cucurbita maxima* leaves [21].

Table II: Amino acids present in *T. occidentalis* grown in different media

Types of amino acids	Amino Acids (g/100g)	Growth medium			*FAO (g/100gcp)
		Humus soil	White Sand	Sawdust	
Essential	Threonine	1.23	0.96	1.75	4.00
	Leucine	0.22	0.17	0.32	7.00
	Isoleucine	0.62	0.48	0.87	4.00
	Lysine	0.92	0.67	1.00	5.50
	Methionine	0.52	0.40	0.67	3.50
	Phenylalamine	0.49	0.38	0.70	6.00
	Tryptophan	0.05	0.04	0.08	1.00
	Valine	0.07	0.05	0.10	5.00
	Histidine	1.73	1.34	2.45	
Non-essential	Alanine	0.51	0.40	0.73	
	Aspartic acid	0.19	0.15	0.27	
	Asparagine	0.27	0.21	0.38	
	Glutamic acid	0.14	0.11	0.20	
	Glutamine	0.70	0.55	1.00	
	Glycine	0.18	0.14	0.26	
	Proline	0.01	0.01	0.01	
	Serine	0.05	0.04	0.06	
	Arginine	0.43	0.33	0.61	
	Cysteine	1.26	0.98	1.78	2.00
	Tyrosine	0.16	0.12	0.22	2.80
Total Amino acid (g/100g)		9.75	7.53	13.46	

*Source: FAO [20]

The proportion of vitamins in *T. occidentalis* leaf under different growth media are shown in Table III. There is variation in vitamins content of same plant grown in different medium. Surprisingly, *T. occidentalis* leaf grown in sawdust had the highest water-soluble and fat-soluble vitamins compared to others. The study suggest that soil physicochemical

composition affects the mineral nutrition of plants grown in it. B-Vitamins obtained is within the recommended daily allowance for adults (19-64 years) in the United Kingdom. However, *T. occidentalis* contains high B-vitamins in relation to the work of Ball [22], who reported that *Geranium robertianum* is a good source of vitamin complex ($B_1 - 288.17 \pm 0.12$ mg/100g; $B_2 - 818.21 \pm 0.07$ mg/100g; $B_3 - 319.13 \pm 0.12$ mg/100g). The vitamin C value present in *T. occidentalis* leaf grown in sawdust and white-sand (17.27%, 9.75% respectively) were higher compared to that reported (5.6 g/100g) by Wasagu *et al.* [19] for *Pistia stratiotes* leaf. Lawal *et al.* [23] reported that fresh pumpkin contains 64.33 ± 4.48 mg/100g vitamin C. The mitochondrial functions are affected by low and dietary intake of B-vitamins and vitamin C [24,25,26,27,28]. Okwu and Josiah [29] reported that ascorbic acid is crucial for body performance. Vitamin C is also antioxidant which performs as an electron donor for eight human enzymes [30]. The vitamin E content of *T. occidentalis* varied in different solid media. Kornteiner *et al.* [31] earlier reported that vitamin E always increases with temperature during seed maturation and also during drought. This observation is in line with result for vitamin E, which showed *T. occidentalis* grown in sawdust had the highest vitamin E content, a reflection of the temperature difference in the media. Vitamin E in pumpkin seed oil makes it an important oil especially in cosmetic applications, human diet, nutrition and health [32]. The vitamin E content of *T. occidentalis* leaf in different solid media were higher compared to pumpkin seed oil as reported by Karanja *et al.* [32]. Presence of vitamin K in *T. occidentalis* as a fat soluble vitamin is important to the overall human health. Douthit *et al.* [33] reported the importance of vitamin K intake on cardiovascular health of an adult. Previous works on the intake of vitamin K have been inconsistent [34,35,36,37,38,39,40,41,42,43]. Also, Jie *et al.* [34] suggested that vitamin K intake might be linked to the development of atherosclerosis, because they found that phylloquinone intakes were significantly lower in postmenopausal women with aortic calcifications than in those without aortic calcifications.

Table III: Percentage vitamins present in *T. occidentalis* grown in different medium

Types of vitamins	Vitamins (%)	Growth medium			*FSA (%)
		Humus soil	White Sand	Sawdust	
Water-soluble	Vitamin B ₁	6.13	9.22	12.54	13
	Vitamin B ₂	0.80	10.34	21.04	28
	Vitamin B ₃	1.25	3.02	7.66	
	Vitamin B ₆	2.12	5.01	11.66	10
	Vitamin B ₁₂	12.70	34.08	65.36	
	Vitamin C	3.16	9.75	17.27	
Fat-soluble	Vitamin A	17.27	16.63	29.65	
	Vitamin E	0.97	2.15	8.07	
	Vitamin K	0.63	1.17	4.01	
Total Vitamins (%)		45.03	91.37	177.26	

*FSA - Food Standards Agency [44]

Conclusion:

The use of humus soil for the cultivation of plants has been a long standing practice for most farmers. It has been reported severally that seeds need water for germination to commence. On the basis of this prerequisite for seeds, pumpkin grown in other solid media shown the capacity to sustain growth and development of several bioactive compounds such as vitamins, amino acids and proximate composition, which serve as the building blocks of living system. Pumpkin grown in sawdust gave the histidine, threonine and cysteine content of the amino acids when compared to other media. Similar results were obtained for water-soluble vitamins and fat-soluble vitamins with variation in proximate composition among pumpkin grown in humus soil and white-sand.

References:

- Abdullahi S.A. (2002). Evaluation of the Nutrient composition of some freshwater fish families in Northern Nigerian. *Journal of Agriculture and Environment*, (2): 141-150.
- Adesina, A. J. and Adeyeye, E. I. (2013). Amino acid profile of three non-conventional leafy vegetables: *Cucurbita maxima*, *Amaranthus viridis* and *Basella alba*, consumed in Ekiti State, Nigeria. *IJAPS*, 3(1): 1-10.
- Akoroda, M. O. (1990). Ethnobotany of *Telfairia occidentalis* (Cucurbitaceae) among Igbos of Nigeria. *Economy Botany* 44 (1): 29-39.
- Akwaowo, E. U., Ndon, B. A. and Etuk, E. U. (2000). Minerals and anti-nutrients in fluted pumpkin (*Telfairia occidentalis* Hook f.). *Food Chemistry*, 70: 235-240.
- Badifu, G. I. O. and Ogunsua, A. O. (1991). Chemical composition of kernels from some species of Cucurbitaceae grown in Nigeria. *Plants Foods Hum. Nutri.* 41: 35-44
- Ball, F. G. (2006). Riboflavin in vitamins in foods analysis: Bioavailability and stability, Taylor and Francis group, USA, 168-173.
- Beulens JW, Bots ML, Atsma F, Bartelink ML, Prokop M, Geleijnse JM, Witteman JC, Grobbee DE, van der Schouw YT. High dietary menaquinone intake is associated with reduced coronary calcification. *Atherosclerosis* 2009;203:489–93.
- Braam L, McKeown N, Jacques P, Lichtenstein A, Vermeer C, Wilson P, Booth S. Dietary phylloquinone intake as a potential marker for a hearthealthy dietary pattern in the Framingham Offspring cohort. *J Am Diet Assoc* 2004;104:1410–4.
- Cox, M. M. and Nelson, D. L. (2011). *Lehninger Principles of Biochemistry*, 5th edn. W. H. Freeman and Company, New York.
- Depeint, F., Bruce, W. R., Shangarl, N., *et al* (2006). Mitochondrial function and toxicity: role of the B-vitamin family on mitochondrial energy metabolism. *Chem. Biol. Interact.*, 163: 94-112.

- 329 Egedigwe, C. A. (2010). Effect of dietary incorporation of *Vernonia amygdalina* and
330 *Vernonia colorata* on blood lipid profile and relative organ weights in albino rats
331 (Thesis). Department of Biochemistry. MOUAU, Nigeria.
- 332 Erkkila " AT, Booth SL, Hu FB, Jacques PF, Lichtenstein AH. Phylloquinone intake and risk
333 of cardiovascular diseases in men. *Nutr Metab Cardiovasc Dis* 2007;17:58–62.
- 334 Erkkila " AT, Booth SL, Hu FB, Jacques PF, Manson JE, Rexrode KM, Stampfer MJ,
335 Lichtenstein AH. Phylloquinone intake as a marker for coronary heart disease risk but
336 not stroke in women. *Eur J Clin Nutr* 2005;59:196–204.
- 337 Food Standards Agency (2003). The national diet and nutritional survey: Adults aged 19 to
338 64 years, Vol. 3. London: The Stationery Office.
- 339 Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, Geleijnse JM, Witteman JC, Grobbee
340 DE, Peeters PH, van der Schouw YT. A high menaquinone intake reduces the
341 incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis* 2009;19:504–10.
- 342 Geleijnse JM, Vermeer C, Grobbee DE, Schurgers LJ, Knapen MH, van der Meer IM,
343 Hofman A, Witteman JC. Dietary intake of menaquinone is associated with a reduced
344 risk of coronary heart disease: the Rotterdam Study. *J Nutr* 2004;134:3100–5.
- 345 Hossain, S. J., Sultana, M. S., Iftexharuzzaman, M., Hossain, S. A. and Taleb, M. A. (2015).
346 Antioxidant potential of common leafy vegetables in Bangladesh. *Bangladesh Journal*
347 *of Botany*, 44(1): 51-57.
- 348 Institute of Medicine (1997). Dietary Reference Intakes for calcium, phosphorus, magnesium,
349 vitamin D and fluoride. Washington DC: National Academic Press.
- 350 Institute of Medicine (1998). Dietary Reference Intakes for thiamine, riboflavin, niacin,
351 vitamin B₆, folate, vitamin B₁₂, pantothenic acid, biotin and choline. Washington DC:
352 National Academic Press.
- 353 Institute of Medicine (2000). Dietary Reference Intakes for vitamin C, vitamin E, selenium
354 and carotenoids. Washington DC: National Academic Press.
- 355 Institute of Medicine (2001). Dietary Reference Intakes for vitamin A, vitamin K, arsenic,
356 boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon,
357 vanadium and zinc. Washington DC: National Academic Press.
- 358 Jie KS, Bots ML, Vermeer C, Witteman JC, Grobbee DE. Vitamin K intake and osteocalcin
359 levels in women with and without aortic atherosclerosis: a population-based study.
360 *Atherosclerosis* 1995;116:117–23.
- 361 Karanja, J. K., Mugendi, B. J., Khamis, F. M. and Muchugi, A. N. (2013). Nutritional
362 composition of the pumpkin (*Cucurbita* spp.) seed cultivated from selected regions in
363 Kenya. *Journal of Horticulture Letters*, 3(1): 17-22.
- 364 Kokwaro, J. (2009). *Medicinal plants of East Africa*. 3rd ed. Nairobi, Kenya. University of
365 Nairobi Press.

- 366 Kumar, H., Choudhary, N., Varsha, N., Kumar, S. and Seth, R. (2014). Phenolic compounds
367 and their health benefits: A review. *Journal of Food Research and Technology*, 2(2):
368 46-59.
- 369 Lawal, O. O., Essien, N. C., Essien, N. M. and Ochalla, F. (2015). Vitamin C content of some
370 processed green leafy vegetables. *European Journal of Experimental Biology*, 5(2):
371 110-112.
- 372 Maas AH, van der Schouw YT, Beijerinck D, Deurenberg JJ, Mali WP, Grobbee DE, van der
373 Graaf Y. Vitamin K intake and calcifications in breast arteries. *Maturitas*
374 2007;56:273– 9.
- 375 Maeda, K., Kuzuya, M., Cheng, X. W., Asai, T., Kanda, S., Tamaya-Mori, N., Sasaki, T.,
376 Shibata, T. and Ighuchi, A. (2003). Green tea catechins inhibit the cultured smooth
377 muscle cell invasion through the basement barrier. *Atherosclerosis*, 166: 23-30.
- 378 Mary K Douthit, Mary Ellen Fain, Joshua T Nguyen, Celestine F Williams, Allison H Jasti,
379 Bernard Gutin, and Norman K Pollock (2017). Phylloquinone Intake Is Associated
380 with Cardiac Structure and Function in Adolescents. *The Journal of Nutrition*, 1960-1967.
- 381 Moran-Palacio, E. F., Tortoledo-Ortiz, O., Yanez-Farias, G. A., Zamora-Alvarez, L. A.,
382 Stephens-Camacho, N. A., Sonanez-Organis, J. G., Ochoa-Lopez, L. M., Rosas-
383 Rodriguez, J. A. (2014). Determination of amino acids in medicinal plants from
384 Southern Sonora, Mexico. *Tropical Journal of Pharmaceutical Research*, 13(4): 601-
385 606.
- 386 Mussatto, G., Dragone, I. and Roberto, C. (2007). Ferulic and p-coumaric acids extraction by
387 alkaline hydrolysis of brewer's spent grain. *Industrial Crops and Products*, 25: 231–
388 237.
- 389 Nakajima, D., Kim, C. S., Oh, T. W., Yang, C. Y., Naka, T., Igawa, S. and Ohta, F. (2001).
390 Suppressive effects of genistein dosage and resistance exercise on bone loss in
391 ovariectomized rats. *Journal Physiological Anthropology and Applied Human*
392 *Science*, 20: 285-291.
- 393 Nwozo, S. O., Adaramoye, O. A., and Ajaiyoba, E. O. (2004). Antidiabetic and
394 hypolipidemic studies of *Telfairia occidentalis* on alloxan-induced diabetic rats. *Nig.*
395 *J. Nat. Prod. and Med.*, 8: 45-47.
- 396 Oboh, G. (2006). Hepatoprotective property of ethanolic and aqueous extract of fluted
397 pumpkin (*Telfairia occidentalis*) leaves against garlic-induced oxidative stress. *J. Med.*
398 *Food.*, 8: 560-563.
- 399 Okwu, D. E. (2004). Phytochemicals and vitamin content of indigenous spices of
400 Southeastern Nigeria. *Journal for Sustaining Agricultural Environment*, 6(1): 30- 37.
- 401 Okwu, D. E. and Josiah, C. (2006). Evaluation of the chemical composition of two Nigerian
402 medicinal plants. *African Journal of Biotechnology*, 5: 357-361.

- 403 Olaniyan, M. F., and Adeleke, A. A. (2005). Study of the effect of pumpkin (*Telfairia*
404 *occidentalis*) milk and raw egg mixture in the treatment of anaemic pregnant women
405 in rural area. *Afri. J. Trad. Complem. and Alt. Med.*, 2: 269-273.
- 406 Olorunfemi, A. E., Arnold, C. I., Emmanuel, O., Nkaima, N., and Akeem, A. (2005).
407 Hypoglycaemic activity of *Telfairia occidentalis* in rats. *J. Pharm. and Biores.*, 2: 36-
408 42.
- 409 Pan Y, Jackson RT. Dietary phylloquinone intakes and metabolic syndrome in US young
410 adults. *J Am Coll Nutr* 2009;28:369–79.
- 411 Sami, R., Li, Y., Qi, B., Wang, S., Zhang, Q., Han, F., Ma, Y., Jing, J. and Jiang, L. (2014).
412 HPLC analysis of water-soluble vitamins (B2, B3, B6, B12 and C) and fat-soluble
413 vitamins (E, K, D, A and β -Carotene) of Okra (*Abelmoschus esculentus*). *Journal of*
414 *Chemistry*, 1-6
- 415 Scalbert, A., Manach, C., Morand, C., Remesy, C. and Jimenez, L. (2005). Dietary
416 polyphenols and the prevention of diseases. *Crit. Rev. Food Sci.* 45: 287-306.
- 417 Senna, L.P., Vanderjert, D.J., River, C.T., Sim, A.T.C. and Muhammad, R.H. (1998).
418 Analysis of nutritional components of eight famine foods of Republic of Niger. *Plant*
419 *Foods for Human Nutrition*, 52: 17-30.
- 420 Smith, L. J., Holbrook, J. T., Wise, R. and Blumenthal, M. (2004). Dietary intake of soy
421 genistein is associated with lung function in patients with asthma. *Journal of Asthma*,
422 41: 833-843.
- 423 Sousa, C., Taveira, M., Valentão, P., Fernandes, F., Pereira, J. A., Estevinho, L., Bento, A.,
424 Ferreres, F., Seabra, R. M. and Andrade, P. B. (2008). Inflorescences of Brassicacea
425 species as source of bioactive compounds: A comparative study. *Food Chemistry*,
426 110: 953-961.
- 427 Stefova, M., Stafilov, T. and Kulevanova, S. (2003). HPLC analysis of flavonoids.
428 *Encyclopedia of chromatography*. Marcel Dekker Inc. New York, pp 1-7.
- 429 Uyoh, E. A., Ita, E. E. and Nwofia, G. E. (2013). Evaluation of the chemical composition of
430 *Tetrapleura tetraptera* (Schum and Thonn.) Taub. Accessions from Cross River State,
431 Nigeria. *Int. J. Med. Arom. Plants*, 3(3): 386-394.
- 432 Villines TC, Hatzigeorgiou C, Feuerstein IM, OMalley PG, Taylor AJ. Vitamin K1 intake
433 and coronary calcification. *Coron Artery Dis* 2005;16:199–203.
- 434 Wasagu, R.S.U., Lawal, M., Shehu, S., Alfa, H.H. and Muhammad, C. (2013). Nutritive
435 values, mineral and antioxidant properties of *Pistia stratiotes* (Water lettuce).
436 *Nigerian Journal of Basic and Applied Science*, 21(4): 253-257.
- 437
- 438 15 AOAC (1984). *Official Methods of Analysis*. 14th Edition, Association of Analytical
439 Chemists, Washington DC, 249-252.

- 17 Abdullahi S.A. (2002). Evaluation of the Nutrient composition of some freshwater
fish families in Northern Nigerian. *Journal of Agriculture and Environment*, (2): 141-150.
- 21 Adesina, A. J. and Adeyeye, E. I. (2013). Amino acid profile of three non-
conventional leafy vegetables: *Cucurbita maxima*, *Amaranthus viridis* and *Basella alba*,
consumed in Ekiti State, Nigeria. *IJAPS*, 3(1): 1-10.
- 6 Akoroda, M. O. (1990). Ethnobotany of *Telfairia occidentalis* (Cucurbitaceae) among
Igbos of Nigeria. *Economy Botany* 44 (1): 29-39.
- 8 Akwaowo, E. U., Ndon, B. A. and Etuk, E. U. (2000). Minerals and anti-nutrients in
fluted pumpkin (*Telfairia occidentalis* Hook f.). *Food Chemistry*, 70: 235-240.
- 7 Badifu, G. I. O. and Ogunsua, A. O. (1991). Chemical composition of kernels from
some species of Cucurbitaceae grown in Nigeria. *Plants Foods Hum. Nutri.* 41: 35-
44
- 22 Ball, F. G. (2006). Riboflavin in vitamins in foods analysis: Bioavailability and
stability, Taylor and Francis group, USA, 168-173.
- 41 Beulens JW, Bots ML, Atsma F, Bartelink ML, Prokop M, Geleijnse JM, Witteman
JC, Grobbee DE, van der Schouw YT. High dietary menaquinone intake is associated
with reduced coronary calcification. *Atherosclerosis* 2009;203:489–93.
- 35 Braam L, McKeown N, Jacques P, Lichtenstein A, Vermeer C, Wilson P, Booth S.
Dietary phyloquinone intake as a potential marker for a hearthealthy dietary pattern
in the Framingham Offspring cohort. *J Am Diet Assoc* 2004;104:1410–4.
- 3 Cox, M. M. and Nelson, D. L. (2011). *Lehninger Principles of Biochemistry*, 5th edn.
W. H. Freeman and Company, New York.
- 24 Depeint, F., Bruce, W. R., Shangarl, N., *et al* (2006). Mitochondrial function and
toxicity: role of the B-vitamin family on mitochondrial energy metabolism. *Chem.*
Biol. Interact., 163: 94-112.
- 33 Mary K Douthit, Mary Ellen Fain, Joshua T Nguyen, Celestine F
Williams, Allison H Jasti, Bernard Gutin, Norman K Pollock. *Journal of*
Nutrition 147 (10), 1960-1967. 2017
- 10 Egedigwe, C. A. (2010). Effect of dietary incorporation of *Vernonia amygdalina* and
Vernonia colorata on blood lipid profile and relative organ weights in albino rats
(Thesis). Department of Biochemistry. MOUAU, Nigeria.
- 39 Erkkila AT, Booth SL, Hu FB, Jacques PF, Lichtenstein AH. Phyloquinone intake
and risk of cardiovascular diseases in men. *Nutr Metab Cardiovasc Dis* 2007;17:58–
62.

- 474 38 Erkkila AT, Booth SL, Hu FB, Jacques PF, Manson JE, Rexrode KM, Stampfer MJ,
475 Lichtenstein AH. Phylloquinone intake as a marker for coronary heart disease risk but
476 not stroke in women. *Eur J Clin Nutr* 2005;59:196–204.
- 477 20 FAO (1970). Amino Acid content of foods and Biological Data on Proteins, Food and
478 Agricultural Organization of the United Nations, Rome, Italy p. 285.
- 479 44 Food Standards Agency (2003). The national diet and nutritional survey: Adults aged
480 19 to 64 years, Vol. 3. London: The Stationery Office.
- 481 43 Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, Geleijnse JM, Witteman JC,
482 Grobbee DE, Peeters PH, van der Schouw YT. A high menaquinone intake reduces
483 the incidence of coronary heart disease. *Nutr Metab Cardiovasc Dis* 2009;19:504–10.
- 484 36 Geleijnse JM, Vermeer C, Grobbee DE, Schurgers LJ, Knapen MH, van der Meer IM,
485 Hofman A, Witteman JC. Dietary intake of menaquinone is associated with a reduced
486 risk of coronary heart disease: the Rotterdam Study. *J Nutr* 2004;134:3100–5.
- 487 1 Hossain, S. J., Sultana, M. S., Iftekharuzzaman, M., Hossain, S. A. and Taleb, M. A.
488 (2015). Antioxidant potential of common leafy vegetables in Bangladesh. *Bangladesh*
489 *Journal of Botany*, 44(1): 51-57.
- 490 25 Institute of Medicine (1997). Dietary Reference Intakes for calcium, phosphorus,
491 magnesium, vitamin D and fluoride. Washington DC: National Academic Press.
- 492 26 Institute of Medicine (1998). Dietary Reference Intakes for thiamine, riboflavin,
493 niacin, vitamin B₆, folate, vitamin B₁₂, pantothenic acid, biotin and choline. Washington DC:
494 National Academic Press.
- 495 27 Institute of Medicine (2000). Dietary Reference Intakes for vitamin C, vitamin E,
496 selenium and carotenoids. Washington DC: National Academic Press.
- 497 28 Institute of Medicine (2001). Dietary Reference Intakes for vitamin A, vitamin K,
498 arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel,
499 silicon, vanadium and zinc. Washington DC: National Academic Press.
- 500 34 Jie KS, Bots ML, Vermeer C, Witteman JC, Grobbee DE. Vitamin K intake and
501 osteocalcin levels in women with and without aortic atherosclerosis: a population-
502 based study. *Atherosclerosis* 1995;116:117–23.
- 503 32 Karanja, J. K., Mugendi, B. J., Khamis, F. M. and Muchugi, A. N. (2013). Nutritional
504 composition of the pumpkin (*Cucurbita* spp.) seed cultivated from selected regions in
505 Kenya. *Journal of Horticulture Letters*, 3(1): 17-22.
- 506 31 M. Kornsteiner, W. Karl-Heinz, I. Elmadfa **Tocopherols and total phenolics in 10 different**
507 **nut types** *Food Chemistry*, 98 (2006), pp. 381-387
508
- 509 9 Kokwaro, J. (2009). *Medicinal plants of East Africa*. 3rd ed. Nairobi, Kenya.
510 University of Nairobi Press.

- 511 23 Lawal, O. O., Essien, N. C., Essien, N. M. and Ochalla, F. (2015). Vitamin C content
512 of some processed green leafy vegetables. *European Journal of Experimental*
513 *Biology*, 5(2): 110-112.
- 514 40 Maas AH, van der Schouw YT, Beijerinck D, Deurenberg JJ, Mali WP, Grobbee DE,
515 van der Graaf Y. Vitamin K intake and calcifications in breast arteries. *Maturitas*
516 2007;56:273– 9.
- 517 4 Moran-Palacio, E. F., Tortoledo-Ortiz, O., Yanez-Farias, G. A., Zamora-Alvarez, L.
518 A., Stephens-Camacho, N. A., Sonanez-Organis, J. G., Ochoa-Lopez, L. M., Rosas-
519 Rodriguez, J. A. (2014). Determination of amino acids in medicinal plants from
520 Southern Sonora, Mexico. *Tropical Journal of Pharmaceutical Research*, 13(4): 601-
521 606.
- 522 11 Nwozo, S. O., Adaramoye, O. A., and Ajaiyoba, E. O. (2004). Antidiabetic and
523 hypolipidemic studies of *Telfairia occidentalis* on alloxan-induced diabetic rats. *Nig.*
524 *J. Nat. Prod. and Med.*, 8: 45-47.
- 525 14 Oboh, G. (2006). Hepatoprotective property of ethanolic and aqueous extract of fluted
526 pumpkin (*Telfairia occidentalis*) leaves against garlic-induced oxidative stress. *J.*
527 *Med. Food.*, 8: 560-563.
- 528 16 Okwu, D. E. (2004). Phytochemicals and vitamin content of indigenous spices of
529 Southeastern Nigeria. *Journal for Sustaining Agricultural Environment*, 6(1): 30-
530 37.
- 531 29 Okwu, D. E. and Josiah, C. (2006). Evaluation of the chemical composition of two
532 Nigerian medicinal plants. *African Journal of Biotechnology*, 5: 357-361.
- 533 12 Olaniyan, M. F., and Adeleke, A. A. (2005). Study of the effect of pumpkin (*Telfairia*
534 *occidentalis*) milk and raw egg mixture in the treatment of anaemic pregnant women
535 in rural area. *Afri. J. Trad. Complem. and Alt. Med.*, 2: 269-273.
- 536 13 Olorunfemi, A. E., Arnold, C. I., Emmanuel, O., Nkaima, N., and Akeem, A. (2005).
537 Hypoglycaemic activity of *Telfairia occidentalis* in rats. *J. Pharm. and Biores.*, 2: 36-
538 42.
- 539 42 Pan Y, Jackson RT. Dietary phylloquinone intakes and metabolic syndrome in US
540 young adults. *J Am Coll Nutr* 2009;28:369–79.
- 541 2 Scalbert, A., Manach, C., Morand, C., Remesy, C. and Jimenez, L. (2005). Dietary
542 polyphenols and the prevention of diseases. *Crit. Rev. Food Sci.* 45: 287-306.
- 543 18 Senna, L.P., Vanderjert, D.J., River, C.T., Sim, A.T.C. and Muhammad, R.H. (1998).
544 Analysis of nutritional components of eight famine foods of Republic of Niger. *Plant*
545 *Foods for Human Nutrition*, 52: 17-30.
- 546 5 Sousa, C., Taveira, M., Valentão, P., Fernandes, F., Pereira, J. A., Estevinho, L.,
547 Bento, A., Ferreres, F., Seabra, R. M. and Andrade, P. B. (2008). Inflorescences of

548 Brassicacea species as source of bioactive compounds: A comparative study. *Food*
549 *Chemistry*, 110: 953-961.

550 30 Uyoh, E. A., Ita, E. E. and Nwofia, G. E. (2013). Evaluation of the chemical
551 composition of *Tetrapleura tetraptera* (Schum and Thonn.) Taub. Accessions from
552 Cross River State, Nigeria. *Int. J. Med. Arom. Plants*, 3(3): 386-394.

553 37 Villines TC, Hatzigeorgiou C, Feuerstein IM, OMalley PG, Taylor AJ. Vitamin K1
554 intake and coronary calcification. *Coron Artery Dis* 2005;16:199–203.

555 19 Wasagu, R.S.U., Lawal, M., Shehu, S., Alfa, H.H. and Muhammad, C. (2013).
556 Nutritive values, mineral and antioxidant properties of *Pistia stratiotes* (Water lettuce).
557 *Nigerian Journal of Basic and Applied Science*, 21(4): 253-257.

558