

Original Research Article

Influence of bio-regulators on the fruit quality and yield attributes of phalsa (*Grewia subinaequalis* DC)

ABSTRACT

Aims: To examine the influence of bio regulators on the fruit quality and yield attributes of phalsa.

Study design: Randomised Block Design.

Place and Duration of Study: Phalsa orchard, Department of Horticulture, Khalsa College, Amritsar during 2015-2016

Methodology: The present study comprised of ten treatments which replicated thrice Uniform sized mature fruits were harvested for carrying out physico-chemical analysis. The observations were recorded viz., fruit length (cm), fruit breadth (cm), fruit weight (g), fruit colour, TSS (%), titratable acidity (%), TSS: acid ratio, reducing sugars (%), total sugars (%), ascorbic acid (mg/100 g pulp), stone length (cm), stone breadth (cm), stone weight (g), harvesting span (days), number of pickings and fruit yield per plant (kg). A sample of ten randomly selected fruits was being taken from each replication for fruit physical and bio chemical analysis and then average was worked out for each parameter.

Results: The results revealed that GA₃ 150 ppm treated plants yielded fruits with maximum fruit length (1.26 cm), fruit breadth (1.43 cm), fruit weight (1.20 g), reducing sugars (7.49 %), ascorbic acid (31.10 mg/100 g pulp) and yield (3.48 kg/plant). Minimum stone length (0.50 cm), stone breadth (0.33 cm) and stone weight (0.045 g) was recorded in fruits obtained from plants treated with NAA 25 ppm. Ethrel 1000 ppm significantly improved TSS (20.23%), fruit colour (purple black), total sugars (12.79 %) whereas reduced the titratable acidity (1.54 %), harvesting span (22.81 days) and number of pickings (3.90).

Conclusion: The results of the present study concluded that plants treated with GA₃ 150 ppm registered significant improvement in fruit physical and biochemical parameters and also fruit yield per plant was increased. The fruits obtained from the plants treated with NAA 25 ppm had small sized and minimum weighed stone. Whereas the yield attributes were enhanced with the application of ethrel 1000 ppm in terms of advancement of harvesting and also reduction in number of pickings.

Keywords: Phalsa; bio-regulators; fruit physical parameters; fruit biochemical parameters; yield

1. INTRODUCTION

The Phalsa (*Grewia subinaequalis* DC) belonging to the family Tiliaceae, is an important indigenous fruit for arid and semi- arid region because of its hardy nature and capacity to tolerate high temperature and even to be grown under prolonged dry spells [1]. India is considered to be the home of phalsa and it is commercially cultivated in Punjab, Uttar Pradesh, Madhya Pradesh, Haryana and Rajasthan [2]. In Punjab it is mostly grown in Ropar, Hoshiarpur, Gurdaspur and Amritsar district. Area under phalsa in Punjab is only 30 hectares with an annual production of 196 tonnes [3].

Plant growth regulators are chemicals that effect flowering, ageing, root growth, prevention and promotion of stem elongation, color enhancement of phalsa fruits. Very small concentration of these substances produces major growth changes. They are widely used for increasing fruit set, controlling fruit drop, enhancing quality and uniform maturity.

Among growth regulators GA₃, NAA, Ethrel are widely used for enhancing superior characters of fruit crop. Application of GA₃ results in increased yield and better grade of phalsa [4]. Application of ethrel improves fruit quality in terms of TSS, reducing and total sugar [5].

Ethrel is a compound that decomposes to release the natural plant hormone ethylene. Its main use in fruit production is to enhance fruit ripening by permitting its harvest about one week earlier and also reduces the span of harvesting and number of pickings significantly [5]. It is thought that NAA may influence the rate of metabolic activity of developing embryos and even also stimulate ethylene biosynthesis which inhibit fruit growth and promote the abscission of weaker fruits. The use of bio regulator like GA₃ has proved effective for increasing the size of berry or fruit and improved quality in crop like grape, citrus, ber etc [2].

The bio regulators have proved to be effective in increasing fruit set and yield in phalsa as well as quality of fruit in terms of TSS, reducing sugars and total sugars. Keeping the role of bio regulators in view, the present study was carried out for improved fruit set, quality and yield of phalsa.

2. MATERIALS AND METHODS

The trial was conducted on well maintained 9 years old phalsa orchard of Department of Horticulture, Khalsa College Amritsar during 2015-16. In the trial 30 uniform sized trees were selected and sprayed with different concentrations of bio regulators as NAA (25, 50 and 75 ppm), GA₃ (50, 100 and 150 ppm) and Ethrel (250, 500 and 1000 ppm). NAA and GA₃ were sprayed at fruit set, whereas Ethrel was sprayed 20 to 25 days prior to harvesting to find out the optimum concentration of these bio regulators. Uniform sized mature fruits were harvested for carrying out physico-chemical analysis. A sample of ten randomly selected fruits was being taken from each treatment and their size was noted using the vernier's caliper. The weight of fruits from randomly drawn sample was computed with the help of electronic weighing balance. The fruits were evaluated for colour rating by a panel of five judges on score card (maximum 10 points) based on colour. Fruits were rated as follows:

Acceptability	Grading	Colour
Excellent	9.1-10	Purple black
Very good	8.1-9	Cherry red
Good	7.1-8	Light red
Fair	6.1-7	Greenish red
Poor	6 and below	Light green

The stones from the same sample were extracted and their size and weight were recorded with the similar procedures taken for fruits. The fruit TSS was analyzed with the help of hand refractometer. Titratable acidity was calculated by titrating diluted juice against N/10 NaOH by using phenolphthalein as an indicator. The diluted juice was prepared by adding distilled water in the ratio of 1:10. The end point of the titration was permanent pink colour. The acidity was calculated and expressed in terms of citric acid [6] by using the below given formula,

$$\text{Titrate acidity(\%)} = \frac{0.064 \times \frac{N}{10} \text{NaOH}}{\text{Volume of juice taken}} \times 100$$

Reducing sugars and total sugars were analyzed by Lyan and Neyon method. Ascorbic acid were analysed by Indophenol dye method. The fruits were picked at regular intervals when ripe, the number of pickings was recorded and days taken to ripe from fruit setting were counted as harvesting span. The fruits were picked on alternate days and were weighed by using physical balance. Average yield per plant was worked out. The recorded data were computed by randomized block design.

3. RESULTS AND DISCUSSION

3.1 Fruit physical parameters

The data presented in Table 1 revealed that most of the treatments improved the fruit size, however GA₃ 150 ppm registered maximum fruit length (1.26 cm) and fruit breadth (1.43 cm). Fruit size significantly increased due to application of gibberellin this might be due to the fact that gibberellins indirectly affect the level of auxin that ultimately caused cell elongation by enlargement of vacuoles and loosening of cell wall after increasing its palatability.

Table 1. Influence of various bio-regulators on fruit physical parameters of phalsa

Treatments	Fruit Size		Fruit weight (g)	Fruit colour
	Fruit length	Fruit breadth		
	(cm)	(cm)		
T ₁ - NAA 25 ppm	0.91	1.13	0.98	7.60
T ₂ - NAA 50 ppm	1.00	1.22	1.04	7.80
T ₃ -NAA 75 ppm	1.08	1.28	1.07	7.90
T ₄ - Ethrel 250 ppm	1.19	1.33	1.06	8.80
T ₅ - Ethrel 500 ppm	1.09	1.33	1.08	8.90
T ₆ - Ethrel 1000 ppm	1.18	1.42	1.14	9.20
T ₇ - GA ₃ 50 ppm	1.23	1.35	1.13	8.40
T ₈ - GA ₃ 100 ppm	1.23	1.40	1.13	8.60
T ₉ - GA ₃ 150 ppm	1.26	1.42	1.20	8.70
T ₁₀ - Control	0.95	1.11	0.78	7.10
Mean	1.11	1.30	1.06	8.30
CD (<i>P</i> =.05)	0.08	0.06	0.06	0.70

The present analysis is in accordance with [7] and [8]. GA₃ 150 ppm (T₉) treated plants gave fruits with maximum weight of 1.20 g, which might be due to the fact that gibberellins increase the cell division and translocation of food material [8].

Treatment T₉ was also found to be at par with the treatment T₆ (Ethrel 1000 ppm) with fruit weight of 1.14 g. Results of these findings were confirmed by [4] and [1]. The plants treated with treatment T₆ produced fruits with excellent purple black colour. This might be due to the fact that ethrel enhanced chlorophyll degradation and increased carotene synthesis [9].

3.2 Fruit biochemical parameters

The data regarding fruit biochemical parameters as influenced by various concentrations of ethrel, NAA and GA₃ are presented in Table 2. Significant TSS (20.23 %) was observed in fruits obtained from plants treated with ethrel 1000 ppm. This may be due to the quick metabolic transformation of starch and pectin into soluble compounds and rapid translocation of sugars from the leaves to the developing fruits [10]. The results of these findings were also confirmed by [4], [2] and [5]. The fruits with minimum acidic content (1.54 %) were obtained from plants treated with ethrel 1000 ppm. This might be due to the fact that ethrel enhance the conversion of organic acids to sugars which increases the sugar content and decreases the acidity [11]. The present study is in conformity with the findings of [1] and [12]. The data related to sugars showed that plants treated with Ethrel 1000 ppm yielded fruits with maximum reducing sugars (7.49 %) and total sugars (12.79 %). This might be due to that ethrel promoted hydrolysis of starch into sugars [5].

Table 2. Influence of various bio-regulators on fruit bio-chemical parameters of phalsa

Treatments	TSS (%)	Titrateable acidity (%)	Reducing Sugars (%)	Total sugars (%)	Ascorbic acid (mg/100 g pulp)
T ₁ - NAA 25 ppm	15.23	2.97	5.38	9.78	22.60
T ₂ - NAA 50 ppm	15.81	2.27	5.49	10.52	23.86
T ₃ - NAA 75 ppm	15.75	2.29	5.54	10.47	23.91
T ₄ - Ethrel 250 ppm	16.07	2.22	6.02	10.69	25.03
T ₅ - Ethrel 500 ppm	18.48	1.80	6.77	12.75	25.28
T ₆ - Ethrel 1000 ppm	20.23	1.54	7.49	12.79	25.62
T ₇ - GA ₃ 50 ppm	17.13	2.22	6.69	12.35	30.06
T ₈ - GA ₃ 100 ppm	16.78	2.10	7.32	12.04	30.98
T ₉ - GA ₃ 150 ppm	18.09	1.97	7.28	12.62	31.10
T ₁₀ - Control	15.16	2.42	5.27	10.15	20.02
Mean	16.87	2.18	6.32	11.41	25.84
CD (P=.05)	.38	.52	.41	.82	.99

The present results are in conformity with the findings of [1] and [12]. Ascorbic acid content in fruits increased significantly by the use of gibberellins. Maximum ascorbic acid content (31.10 mg/100 g pulp) was observed in fruits obtained from

plants treated with GA₃ 150 ppm. This might be due to the possible catalytic influence of gibberlic acid on biosynthesis of ascorbic acid from sugar or inhibition of oxidative enzymes or both [13]. The results of these findings are in conformity with [14] and [15].

3.3 Stone parameters

The data presented in Table 3 revealed that all the treatments showed non-significant effect on the stone size in phalsa. Minimum stone length (0.50 cm) and stone breadth (0.33 cm) was registered under the treatment T₁. From the data, it has been observed that NAA 25 ppm treated plants gave fruits with least stone weight (0.045 g). The present study is in conformity with the findings of [8].

Table 3. Influence of various bio-regulators on stone parameters of phalsa

Treatments	Stone length (cm)	Stone breadth (cm)	Stone weight (g)
T ₁ - NAA 25 ppm	0.50	0.33	0.04
T ₂ - NAA 50 ppm	0.56	0.40	0.07
T ₃ -NAA 75 ppm	0.53	0.40	0.06
T ₄ - Ethrel 250 ppm	0.53	0.40	0.11
T ₅ - Ethrel 500 ppm	0.56	0.43	0.08
T ₆ - Ethrel 1000 ppm	0.56	0.43	0.08
T ₇ - GA ₃ 50 ppm	0.53	0.41	0.10
T ₈ - GA ₃ 100 ppm	0.56	0.43	0.11
T ₉ - GA ₃ 150 ppm	0.56	0.46	0.09
T ₁₀ - Control	0.58	0.40	0.11
Mean	0.55	0.41	0.09
CD (P=.05)	NS	NS	0.02

3.4 Yield parameters

The presented data (Table 4) showed that ethrel significantly improved the yield parameters in phalsa. Ethrel 1000 ppm treated plants registered minimum harvesting span (22.81 days) than rest of the treatments. The number of pickings also significantly reduced by the use of ethrel. Minimum number of pickings (3.90) was recorded in plants treated with ethrel 1000 ppm. This might be due to that ethrel hasten the maturation process [16]. These findings are in conformity with the findings of [1] and [5]. The plants treated with GA₃ 150 ppm registered maximum yield (3.48 kg/plant) and was followed by treatments T₈ and T₆ with fruit yield of 3.47 and 3.36 kg/plant respectively. All of these treatments were found to be at par with each other. Yield is a complex character and is characterized by increase in weight of individual fruit and number of fruits. It might be due to the fact that GA₃ increased the translocation and mobilization of photosynthates from source to sink [17]. The results are in accordance with the findings of [4], [14] and [18] in ber, [19] in Plum and [20] in mango.

111 **Table 4. Influence of various bio-regulators on yield parameters of phalsa**

Treatments	Harvesting span (days)	Number of pickings	Yield (kg/plant)
T ₁ - NAA 25 ppm	27.71	6.56	2.51
T ₂ - NAA 50 ppm	27.32	6.31	2.76
T ₃ - NAA 75 ppm	26.72	6.08	2.80
T ₄ - Ethrel 250 ppm	23.02	4.39	2.85
T ₅ - Ethrel 500 ppm	23.51	4.15	3.17
T ₆ - Ethrel 1000 ppm	22.81	3.9	3.36
T ₇ - GA ₃ 50 ppm	26.42	6.01	3.23
T ₈ - GA ₃ 100 ppm	25.13	5.57	3.47
T ₉ - GA ₃ 150 ppm	24.12	5.26	3.48
T ₁₀ - Control	30.52	7.13	1.95
Mean	25.72	5.53	2.95
CD (P=.05)	.07	.58	.15

112
113 **4. CONCLUSION**

114 The results of the present study concluded that plants treated with GA₃ 150 ppm registered significant improvement in fruit
115 physical and biochemical parameters and also fruit yield per plant was increased. The fruits obtained from the plants
116 treated with NAA 25 ppm had small sized and minimum weighed stone. Whereas the yield attributes were enhanced with
117 the application of ethrel 1000 ppm in terms of advancement of harvesting and also reduction in number of pickings.

118
119
120 **COMPETING INTERESTS**

121
122 Authors have declared that no competing interests exist.

123
124
125 **REFERENCES**

- 126
127 1. Singh JP, Kumar S, Katiyar PN, Dwivedi AK. Effect of calcium nitrate, GA₃ and ethrel on fruiting, ripening and
128 chemical traits of phalsa (*Grewia subinaequalis* D. C.). 2011; Annals Hort 4: 72-76.
- 129 2. Kacha HL, Viradia RR, Leua HN, Jat G, Tank AK. Effects of NAA, GA₃ and ethrel on yield and quality of phalsa
130 under south- surashtra condition. Asian J Hort. 2012; 7: 242-245.
- 131 3. Pujari S. Phalsa cultivation in India – Production area, Climate, Harvesting and Fruit handling. 2015.
132 [http://www.yourarticlelibrary.com/cultivation/phalsa-cultivation-in-india-production-area-climate-harvesting-and-](http://www.yourarticlelibrary.com/cultivation/phalsa-cultivation-in-india-production-area-climate-harvesting-and-fruit-handling/24703)
133 [fruit-handling/24703](http://www.yourarticlelibrary.com/cultivation/phalsa-cultivation-in-india-production-area-climate-harvesting-and-fruit-handling/24703).
- 134 4. Debnath A, Vanajalatha, Umarfarooque M, Reddy M. Effect of NAA, GA₃, Kinetin and Ethrel on yield and quality
135 in phalsa. Asian J Hort. 2011; 6: 474-477.
- 136 5. Kacha HL, Jat G and Patel SK. Performance of various plant growth regulators on yield and quality of phalsa (*Grewia asiatica* L.). Hortflora Res Spec. 2014; 3:292-294.
- 137
138 6. A O A C (1990) Official and tentative, methods of analysis. Association of official analytical chemist, 14th E.D.
139 Washington D.C.

- 140 7. Shukla HS, Kumar V, Tripathi VK. Effect of gibberellic acid and boron on development and quality of aonla fruit
141 "Banarasi". ISHS Acta Hort. 2011; 890: 375.
- 142 8. Chandra R, Rawat M, Singh SS, Singh KK. Effect of foliar application of various growth regulators on yield and
143 quality of Aonla cv. Na-7 National Academy Agricultural Sci. 2015; 33:2123-2127.
- 144 9. Al-Mughrabi MA, Bacha MA, Abdelrahman AO. Influence of pre harvest application of ethrel and 2, 4-D on fruit
145 quality of Balady orange. Agri Sci. 1989; 1: 95-102.
- 146 10. Tripathi VK, Shukla PK. Effect of plant bio- regulators, zinc sulphate and boric acid on yield and quality of
147 strawberry cv. Chandler. J Asian Hort. 2007; 4: 15-18.
- 148 11. Singh P, Kumar S, Maji S. Effect of ethrel on post harvest changes in papaya (*Carica papaya* L.) fruits HortFlora
149 Res Spect. 2012; 1: 225-230.
- 150 12. Meitei SB, Patel RK, Deka BC, Deshmukh NA, Singh A. Effect of chemical thinning on yield and quality of peach
151 cv. Flordasun. African J Agricultural Res. 2013; 8:3558-3565.
- 152 13. Kher R, Bhat S, Wali VK. Effect of foliar application of GA₃, NAA and CCC on physico-chemical characteristics of
153 guava cv. Sardar. Haryana J Hort Sci. 2005; 34:31-32.
- 154 14. Tripathi D, Pandey AK, Pal AK, Yadav MP. Studies on effect of plant growth regulators on fruit drop,
155 development, quality and yield of ber cv. Banarasi Karaka. Progressive Horticulture. 2009; 41:184-186.
- 156 15. Wahdan MT, Habib SE, Bassal MA, Qaoud EM. Effect of some chemicals on growth, fruiting, yield and fruit
157 quality of " Succary Abiad" mango cv. J American Sci 2011; 7:651-658.
- 158 16. Tyler K, May D, Miller R. Ethrel sprays reduce number of pickings in hand- harvested cantaloupes. California
159 Agriculture. 1970; 24:6-7.
- 160 17. Krishnamoorthy HN. Gibberellins. Plant growth substances 1981; pp: 77-87.
- 161 18. Gill KS, JS Bal. Impact of application of growth regulators on Indian jujube. International Jujube Symposium,
162 2011; 993: 119-124.
- 163 19. Kaur H, Singh A, Gupta M, Randhawa JS. Effect of NAA and gibberellic acid on pre- harvest fruit drop and quality
164 of Satluj Purple plum. Haryana J Hort Sci. 2008; 37:31-32.
- 165 20. Rohit S. Effect of GA₃ and urea spray on flowering fruit retention and fruit quality of mango cv. Langra. Doctoral
166 dissertation, JNKKV. 2014.