

Water deficit as a limiting factor to the initial growth of coffee Conilon variety Diamante

ABSTRACT

The water deficit is considered one of the main limiting factors of agricultural production, studies that aim to understand it become essential for improving productivity and rational use of water resources. Thus, the objective of this work was to evaluate the influence of the reduction of the fraction of transpirable soil water (FTSW) under the growth variables of the nine clones that compose the variety "Diamante Incaper ES8112" of the coffee Conilon and to estimate the critical FTSW (when the growth potential is reduced by the limitation of the transpiration process). The nine experiments were conducted in a greenhouse, located in the Center of Agrarian Sciences and Engineering of the Federal University of Espírito Santo, in the city of Alegre-ES. Each experiment consisted of one of the nine clones that compose the cultivar Diamante Incaper ES 8112, with two treatments: T0 (without water deficit) and Td (water deficit induced until the plants reach 10% of the relative transpiration of the T0 treatment) with 8 replications. The variables evaluated were: relative transpiration (RT), plant height (PH) and leaf area (LA). At the end of the experiments it was verified that the clones are affected at different times, showing that they have different water needs, being clone 104 characterized as the most resistant to the water deficit in the soil, for keeping the TR to a lower value of FTSW (0.47) and clone 108 was characterized as the most sensitive to soil moisture variation, with critical FTSW value of 0.82.

Keywords: Water resources, *Coffea canephora*, FTSW, irrigation management.

1. INTRODUCTION

Coffee production is a highlight in the Brazilian agricultural scenario, and according to the National Supply Company [1] the coffee production for this year is estimated between 54.4 and 58.5 million bags benefited. The total area, in formation and production, amounts to 2,202.6 thousand hectares, of which 286.5 are in formation and 1,916.1 thousand hectares in production. The coffee conilon accounts for approximately 24% of the country's total coffee production, estimated at between 12.7 and 13.96 million bags [1], with the state of Espírito Santo being the largest national producer, with around 7.7 and 8.7 million bags estimated, corresponding to a growth between 29.5 and 46.3% over the previous year.

In Brazil and in other regions of the world, drought is considered the main climatic aggravating factor affecting coffee production [2, 3]. The water deficit besides affecting the water relations in the plants, altering the metabolism, is a phenomenon that occurs to a great extent of the cultivable areas [4], being responsible for most of the variability of the final production [5].

CONAB [6] affirms that the occurrence of a severe drought in the State of Espírito Santo accompanied by high temperatures and sunshine between December 2014 and February 2015, directly impacted productivity (a decrease of 22% in relation to the previous year). The prolonged period of drought in the State generated a serious water crisis that served as a warning for studies related to the water aspect.

Faced with the problem of water deficit, irrigation becomes a practice constantly employed in regions with water deficit, because it plays a major role for production, since investments in the sector result in a substantial increase in productivity and production value, which causes pressure reduction by the incorporation of new areas for cultivation [7]. However, for the adequate use of this technique,

knowledge about the water requirement of crops is essential in order to guarantee satisfactory results and the rational use of available water resources. One of the factors that can be used to express the content of water in the soil is the fraction of transpirable soil water (FTSW), widely used for the evaluation of plant responses to water deficit. It is assumed in this concept that the water content in the soil used by the plant for transpiration oscillates between the soil water content in the field capacity when the transpiration is maximum; and the water content in the soil, when the transpiration of the plant is equal to 10% of the maximum [8]. This seems to be the concept that most closely approximates as an indicator of the actual amount of water in the soil that can be extracted by the plants for transpiration [9]. For the determination of FTSW, two stages [8, 10, 11] are considered: the stage I is when water is freely available in the soil, the plant has no water deficiency and its stomatal conductance and transpiration are maximal, that is, the content of transpirable water in the soil does not limit stomatal conductance and transpiration; and the stage II when the available water in the soil decreases and the plant reduces its stomatal conductance and transpiration to maintain its water balance and cellular turgescence. Lago et al. [12] affirms that several studies have been conducted to verify critical FTSW values ranging from 0.30 to 0.71 in annual agricultural species and from 0.70 to 0.90 in forest species. But, the knowledge of the influence of these values on the growth of conilon coffee is still limited. Therefore, the objective of this study was to quantify the influence of the soil water deficit, represented by FTSW, on the transpiration, plant height and leaf area of conilon coffee (*Coffea canephora*) and to determine critical FTSW, that is, the moment in that the fraction of transpirable soil water begins to limit the growth variables.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The experiments were conducted in a greenhouse located at the Center of Agrarian Sciences and Engineering of the Federal University of Espírito Santo (CCAUE-UFES), located in the city of Alegre-ES (20°45' S, 41°32' W and altitude of 269.0 m). The climate of the region is of type "Aw" with dry season in winter, according to the classification of Köppen, with average annual temperature of 23°C and annual precipitation around 1,200 mm.

Nine experiments were carried out with the conilon coffee (*Coffea canephora*), variety "Diamante Incaper ES8112", composed of nine clones. Each experiment was composed of one clone of the cultivar and constituted by two treatments: T0 (irrigated throughout the experiment and Td - induced water deficit until the plants reached 10% of the relative transpiration of the T0 treatment) with 8 replicates.

The seedlings of the conilon coffee were obtained in a suitable nursery, certified, these are available in tubes with commercial substrate, free of pathogens and presenting a pattern of three pairs of leaves. During the experiment, the monitoring and management of pests and diseases were carried out.

Each experimental plot was consisted of a 12-liter pot filled with soil characteristic of the region, containing one plant per plastic pot. The eight T0 replicates were used as controls, where the plants grew without water deficit, and the other 8 Td replicates, the water deficit was imposed until reaching the pre-established limit.

The soil used as substrate was classified as a Red-Yellow Latosol, with a medium texture, according to the textural triangle of EMBRAPA, being collected in the depth of 0.00 to 0.30 m, disturbed, passed in a sieve of 4 millimeter and homogenized. The soil samples were sent to the laboratory for chemical and physical analysis, according to the methodology proposed by Almeida et al. [13]. The chemical and physical attributes of the Latosol used as substrate are presented in Table 1.

Table 1. Chemical and physical attributes of the Latosol used as substrate.

pH	P	K	Ca	Mg	BS	CEC	V	Sand	Silt	Clay
	mg dm ⁻³		cmol _c dm ⁻³ %					%		
5.25	2.28	82	0.79	0.43	1.46	4.84	30.09	73	3.0	24

Extraction and determination: pH in water (1:2.5); P, K, Na: Mehlich 1; Ca, Mg, Al: KCl (1M); H+Al: calcium acetate (0.5M), CTC a pH 7.0. For physical analysis slow stirring at 50 rotations per minute for 16 hours, with

wagner type stirrer; chemical dispersant: NaOH 0.1 mol L⁻¹ and determination of the silt and clay fractions by the pipette method [13].

With the results obtained, the soil water retention curve was determined. Then, the soil acidity was corrected according to the methodology described by Prezotti et al. [14] and fertilization as recommended by Novais, Neves and Barros [15].

The pots were coated with white paper to reduce the absorption of solar radiation to minimize the heating of the soil. Subsequently, the pots were placed on a metal bench (3.0 x 0.80 meter) with 1 meter height and weekly the standardization procedure was carried out between the pots of the same treatment.

The was performance of the plants submitted to the water deficit verified by the method of the fraction of transpirable soil water (FTSW) in two stages of water deficit [8]: in stage I, water is freely available in the soil, with no water deficit for the plants, and stomatal conductance and transpiration are potential; in stage II, the available water in the soil decreases, and the stomatal conductance and the transpiration decrease, to maintain the water balance and the cellular turgescence.

For the beginning of the experiment it was necessary to determine the initial weight of each plot (Wi), which corresponds to its weight with the soil moisture near the field capacity. In order to do so, all pots with already established seedlings were saturated with water and left in free drainage for 48 hours in order to determine the weight of the experimental units with the soil moisture corresponding to the field capacity. The soil surface of each pot was covered with white polystyrene to ensure that all lost water was from the transpiration process.

The applications of the water regimes began 30 days after planting of the seedlings, applying the water deficit in the pots referring to the Td, which were not irrigated until the end of the experiment, that is, until the relative transpiration (RT) of the plants of the Td to reach 10% of the transpiration of the T0 plants, which occurred before the plants reached the permanent wilting point.

All experimental plots were weighed daily at 5:00 p.m. on an electronic scale with a capacity of 25 kilograms, and a variation of 0.5 g. Subsequently, each T0 treatment pot represented the response to the amount of water the plant transpired from the previous day. The daily loss of water by the T0 plants was determined according to equation 1 and for Td according to equation 2, as demonstrated by Kelling et al. [16]:

$$RT_0 = W_d - W_i \quad (1)$$

where:

RT₀ – Relative transpiration;

W_d – weight of each pot on the day of weighing; and

W_i – weight of the same pot on the day of the start of the experiment.

$$RT_d = W_d - W_{da} \quad (2)$$

where:

RT_d – Relative transpiration;

W_d – weight of each pot on the day of weighing; and

W_{da} – weight of the same pot on the day of the start of the experiment.

Then, the average daily loss of water in the 8 plants constituting the T0 treatment in each experiment was calculated. With the obtained value the water lost by sweating on the day was returned, each plot returning to its value of Wi. The Td plants did not have water replacement in order to reach the pre-established limit.

With these variables, it was possible to monitor the decrease of RT by the equation 3 [8]. The 10% limit of relative transpiration was adopted because it is assumed that below this transpiration rate the stomas are closed and the loss of water is due only to epidermal conductance.

$$RT = \frac{DTT_d}{ADT_0} * 100 \quad (3)$$

where:

RT – Relative transpiration;

DTT_d – Daily transpiration of treatments that suffer deficit;

ADT₀ – Average daily transpiration of treatment T0.

The end of the water deficit period was determined when the Td treatment plots reached 10% of the relative transpiration of the T0 treatment, at which time the weight of each plot was determined and it was identified as final weight (Wf).

In order to evaluate coffee responses to water deficit, used the concept of the fraction of transpirable soil water (FTSW) [8, 10, 11, 16]. The fraction of transpirable soil water was calculated by equation 4 [8].

$$FTSW = \frac{(W_{daily} - W_f)}{(W_i - W_f)} \quad (4)$$

where:

FTSW – Fraction of transpirable soil water;

W_{daily} – Weight of the experimental plot in each day;

W_i – Initial weight of each experimental plot; and

W_f – Final weight of each experimental plot.

The growth variables evaluated were: plant height (PH, cm - being the length of the main stem at ground level up to the apical bud) and leaf area (LA, cm²), calculated by the sum of the LA of all individual leaves of the plant. The growth analyzes were performed every 3 days, biometrically with the aid of a graduated ruler.

Based on total daily LA data, the daily leaf area (LA) and daily height (PH) increment for each clone was calculated, considering the first normalization with the aim of normalizing the data in the range of 0 and 1, performed by Equation 5 [16].

$$V = V_{d_{total}} - V_{0_{total}} \quad (5)$$

where:

V – Variable in study (LA; PH);

V_{d_{total}} – Increase of the total variable in each Td plant; and

V_{0_{total}} – Increase of the total variable in each T0 plant.

The second normalization was performed with the aim of reducing the variations among plants, caused by the environmental conditions of the greenhouse, according to equation 6 [17]:

$$Nv = \frac{value \ RT \ 10\% - value \ (n)}{value \ RT \ 10\% - v_{initial}} \quad (6)$$

where:

Nv – Normalized variable;

Value RT 10% – final value of the variable (when RT was 10%);

Value (n) – value of the variable on the specific day; and

V_{initial} – value of the variable on the first day of the experiment.

The growth data of the dependent variables obtained from the second normalization were adjusted to a logistic function of the independent FTSW variable in a non-linear sigmoidal model (Equation 7) [11].

$$y = \frac{1}{(1 + \exp(a \cdot (X - b)))} \quad (7)$$

where:

y – Dependent variable (Height and leaf area of plants);

X – Fraction of transpiration soil water; e

a and b - coefficients estimated using non-linear regression procedures.

Adjusted curves were used to determine the critical FTSW value at which relative transpiration reduction was initiated.

3. RESULTS AND DISCUSSION

The value of the fraction of transpirable soil water, in which the relative transpiration reduction (RT) begins, occurs when the curve estimated by the logistic equation moves away from 1 and starts the linear decrease (critical point) [8].

Thus, it is observed in the cultivar Diamante that the critical FATS values for the RT variable were concentrated between a range of values of 0.47 and 0.82, confirming the hypothesis that the clones have different water needs. Souza [18] affirms that it is possible to occur variation among clones in different cultures, since tolerance to water deficit is a genotype-dependent character.

Lago et al. [12] verified variation among clones of the same species in the potato crop, which presented a critical FTSW variation between 0.28 and 0.49, which indicates the existence of genetic variability for tolerance to water deficit. Kelling et al. [16] found variation between different cultivars in chrysanthemum, with oscillation from 0.48 to 0.65. Rodrigues et al. [19] found FTSW variation between species identical, but with different ages and that converged from 0.40 to 0.61.

The critical FTSW determination is a character of interest, since it represents the ability of the genotype to respond to soil water deficit to maintain foliar turgescence [12]. The indication of the exact moment when the growth variables were affected is a fundamental factor to inhibit the effect of the water deficit, avoiding changes in the plant behavior whose irreversibility will depend on the genotype, duration, severity and stage of development of the plant [9].




Figure 1. Normalized relative transpiration (NRT) in function of the decrease of the fraction of transpirable soil water (FTSW), in the nine clones of conilon coffee Diamante Incaper ES8112 ("A"- Clone 101; "B"- Clone 102; "C"- Clone 103; "D"- Clone 104; "E"- Clone 105; "F"- Clone 106; "G"- Clone 107; "H"- Clone 108; "I"- Clone 109).

106; “G”- Clone 107; “H”- Clone 108; “I”- Clone 109). Esm is the efficiency of the statistical model and Sde is the standard error.

The clones that obtained the lowest values were clones 104 and 103 presenting critical FTSW of 0.47 and 0.56, respectively. These results are considered inferior to the results obtained by Rodrigues et al. [21], in the tropical Robusta species with critical FATS of 0.71. But, close to the results obtained for arabica coffee, variety Catucaí Vermelho 785-15 (0.52) [20].

It is possible to indicate that clones 104 and 103 have more efficient stomatal control mechanisms than the other clones that compose the variety, because they can reduce the water consumption, probably due to the less stomatal opening, delaying the effect of reduction in leaf growth [12]. Probably, a form of adaptation to water deficit conditions [11, 17].

The clones 107, 108 had the highest critical FATS values 0.79 and 0.82, respectively. The FTSW value at which the start of the stomas closure occurred is considered high when compared to other annual crops, as soybean (0.40) [8]; for maize (0.36 to 0.60) [21]; for grapevine (0.35) [11]; but close to those obtained for Robusta Tropical of 0.61 [19] and of 0.80 [22].

The productive point of view, the reduction of RT caused by the premature closure of stomas causes impacts, because during a cycle of short-term deficit there will be a loss of productivity [21]. However, from the biological point of view, the species that closes the stomas in higher FTSW will save water, since stomatal closure in a higher soil moisture condition results in water conservation in the soil and constitutes an adaptation of the species, a characteristic that contributes possibly to better withstand prolonged water deficit [10].

The relationships between RT and FTSW and stages I and II of transpiration observed in this species, in which the plants dehydrated under the same conditions of temperature and water demand, reinforces the theory that stomatal closure, to prevent or reduce water loss is determined by soil water content [8].

The behavior of the plant height variable is shown in Figure 2.




Figure 2. Normalized relative height (NRH) in function of the decrease of the fraction of transpirable soil water (FTSW), in the nine clones of conilon coffee Diamante Incaper ES 8112 ("A"- Clone 101; "B"- Clone 102; "C"- Clone 103; "D"- Clone 104; "E"- Clone 105; "F"- Clone 106; "G"- Clone 107; "H"- Clone 108; "I"- Clone 109). Esm is the efficiency of the statistical model and Sde is the standard error.

As a response to the water deficit applied to the conilon coffee Diamante, we obtained for the relative height variable a FTSW range that oscillated from 0.48 to 0.84. The clones 104 and 108 may be characterized as the two most resistant (0.48 and 0.51), followed by clone 105 (0.52). Clones that performed poorly, with growth paralyzing at higher levels were clones 107 and 102, obtaining a critical FTSW of 0.84 and 0.85, respectively.

In some clones it is observed that the critical FTSW values for height variable were larger than the variable RT, that is, the height variable was more sensitive to the water variation, being the first variable affected. This is a statement by Larcher [23], where he reports that the first and most sensitive response to the water deficit is the reduction of turgescence and, associated to this event, is the decrease of the plant growth and extension process.

The results differ from that verified by Araújo [22] who found a critical FTSW value of 0.95 for the coffee plant, but resembles the results of Rodrigues et. al [19] who applied water deficit at 30 days after planting in Robusta Tropical coffee plants where they observed a critical FTSW of 0.67.

Dardengo et al. [24] verified the negative influence of the water deficit on the height of the conilon coffee, the plants that were submitted to the water deficit presented smaller height. Busato et al. [25] when evaluating the initial development of the conilon coffee plant under different irrigation layers found higher values of coffee height for greater water availability in the soil.

The Figure 3 shows the behavior of the relative leaf area variable in function of FATS decrease.




Figure 3. Normalized relative leaf area (NRLA) in function of the decrease of the fraction of transpirable soil water (FTSW), in the nine clones of conilon coffee Diamante Incaper ES 8112 ("A"- Clone 101; "B"- Clone 102; "C"- Clone 103; "D"- Clone 104; "E"- Clone 105; "F"- Clone 106; "G"- Clone 107; "H"- Clone 108; "I"- Clone 109). Esm is the efficiency of the statistical model and Sde is the standard error.

The clones 101, 103 and 104 (critical FTSW 0.37, 0.41 and 0.42) behaved as resistant to the induced water deficit, because they did not paralyze leaf area growth, however the clones 102, 107 and 109 were those with the highest critical FTSW values, 0.68, 0.74 and 0.68, respectively. Thus, they ceased their leaf area growth prematurely in a condition of water stress.

These results indicate that under conditions of low soil water availability, stomatal closure occurs earlier in clones 102, 107 and 109. And this response is an adaptive criterion of clones to withstand prolonged water deficit, conserving for longer the water in the soil [10]. Kelling et al. [16] presents this inhibition of cell expansion as a defense strategy of the plant, because the plants respond to the water deficit according to the intensity of soil water stress, the genotype, the stage of development and the type of organ and cell considered, and may present responses at the morphological, physiological, cellular and metabolic level [26].

However, this plant's strategy to inhibit leaf growth due to the reduction of soil water availability affects metabolism, transport, solutes translocation, cell turgor, cell expansion, and stomas opening and closing [27], which induces the reduction of the growth and yield losses of the culture, since the stomatal closure decreases the CO₂ entry, and consequently reduces the photosynthetic process [28].

In this study, critical FTSW values in which leaf growth begins to decrease ranged from 0.37 to 0.74 and are within the range of values found by Araujo et al. [29] (critical FTSW ranging from 1.0 to 0.40) and Rodrigues et al. [19] (critical FTSW ranging from 0.71 to 0.47) in which they worked with three times of water deficit in coffee.

The data of the present work are also in the range of values of foliar growth described by other works with other cultures, such as 0.55 in peas [19], 0.35 in grapevine [11], 0.44 in rice [30], 0.70 and 0.90 in eucalyptus seedlings [31] and 0.48 to 0.71 in chrysanthemum cultivars [16].

It is observed that the RLA variable presented lower values of critical FTSW in all nine clones worked in relation to RT. This fact indicates that the paralization of the foliar growth occurred after the stomatal closure mechanism that caused the RT limitation, since there was no decrease of the cellular turgescence, thus leading to cell expansion and consequent to the foliar growth [33, 27]. According to Lago et al. [12], this result indicates that the plants may have an efficient stomatal control mechanism, since they can reduce water consumption, probably due to the reduced stomatal opening, delaying the reduction effect on leaf growth.

Although most of the results converge to a single range of critical values of the fraction of transpirable soil water, there are some variations between the different studies. These variations in the critical values of the fraction of transpirable soil water could be attributed to differences in the size of the pots used in each study, soil type or variations in air evaporative demand conditions [34].

Studies of tolerance to crop water deficit should continue to be carried out, to characterize more tolerant genotypes [18] and adaptations to the climatic conditions of each region, to minimize the risks in agricultural production.

4. CONCLUSION

The water deficit limited the growth of the clones in different moments, being verified that these have different water needs.

The RT variable was the first to be affected by FTSW reduction in soil, followed by plant height and leaf area.

Clones 104 and 103 were the most tolerant to FTSW reduction and clones 107 and 108 were the most susceptible soil moisture variation.

REFERENCES

1. Companhia Nacional de Abastecimento – CONAB. Acompanhamento da safra brasileira de café: V. 5 - Safra 2018, N.1 - Primeiro levantamento. 2018; 1-73. Accessed: 20 March 2018. Available:

- http://www.conab.gov.br/OlalaCMS/uploads/arquivos/18_01_24_17_34_27_cafe_janeiro2018.pdf. Portuguese.
2. DaMatta FM. Exploring drought tolerance in coffee: a physiological approach with some insights for plant breeding. *Brazilian Journal of Plant Physiology*. 2004; 16(1): 1-6.
3. Bierhals JD, Ferroni PH. C. Café: Safra 2008/2009 encolhe e preços sobem. In: *AGRIANUAL* (2008): anuário da agricultura brasileira. FNP Consultoria & Comercio. 2007; 213-215. Portuguese.
4. Nogueira RJMC, Moraes JAPV, Burity HA. Alterações na resistência à difusão de vapor das folhas e relações hídricas em aceroleira submetidas a déficit de água. *Revista Brasileira de Fisiologia Vegetal*. 2001; 13(1): 75-87. Portuguese.
5. Costa JO. Padrões de resposta termal ao déficit hídrico na cultura do café irrigada por gotejamento. Dissertação (Mestre em Ciências) –Escola Superior de agricultura Luiz de Queiroz, Piracicaba. 2016. Portuguese.
6. Companhia Nacional de Abastecimento – CONAB. Acompanhamento da safra brasileira de café: V. 2 - Safra 2015, N. 4 - Quarto levantamento. 2015; 1-60. Accessed: 20 March 2018. Available: http://www.conab.gov.br/OlalaCMS/uploads/arquivos/15_12_17_09_02_47_boletim_cafe_dezembro_2015_2.pdf. Portuguese.
7. Agência Nacional de Águas – ANA. Conjuntura dos recursos hídricos no Brasil: informe 2014. Accessed: 08 March 2018. Available: <http://www.brasil.gov.br/meio-ambiente/2015/03/ana-divulga-relatorio-de-conjuntura-dos-recursos-hidricos>. Portuguese.
8. Sinclair TR, Ludlow MM. Influence of soil water supply on the plant water balance of four tropical grain legumes. *Australian Journal Plant Physiology*. 1986; 13(3): 319-340.
9. Santos RF, Carlesso R. Déficit hídrico e os processos morfológico e fisiológico das plantas. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 1998; 2(3): 287-294. Portuguese.
10. Muchow RC, Sinclair TR. Water deficits effects on maize yields modeled under current and “greenhouse” climates. *Agronomic Journal*. 1991; 83(6): 1052-1059.
11. Bindi M, Bellesi S, Orlandini S, Fibbi F, Moriondo M, Sinclair T. Influence of water deficit stress on leaf area development and transpiration of Sangiovese Grapevines grown in pots. *American Journal of Enology and Viticulture*. 2005; 56(3): 68-72.
12. Lago I, Streck NA, Zanon AJ, Hanauer JG, Bisognin DA, Silva MR. Transpiração e Crescimento Foliar de Clones de Batata em Resposta à Fração de Água Transpirável no Solo. *Revista Brasileira de Ciência do Solo*. 2012; 36(3): 745-754. Portuguese.
13. Almeida BG, Donagemma GK, Ruiz HA, Braidia JA, Viana JHM, REICHERT JMM, Oliveira LB, Ceddia MB, Wadt OS, Fernandes RBA, Passos RR, Dechen SCF, Klein VA, Teixeira WG. Padronização de Métodos para Análise Granulométrica no Brasil. Rio de Janeiro: Embrapa, 2012 (Comunicado técnico 66). Portuguese.
14. Prezotti LC, Gomes JÁ, Dadalto GG, Oliveira JA. Manual de Recomendação de Calagem e Adubação para o Estado do Espírito Santo. 5th ed. Vitória: SEEA; 2007. Portuguese.
15. Novais RF, Neves JCL, Barros NF. Ensaio em ambiente controlado. In: Oliveira AJ, Garrido WE, Araújo JD, Lourenço S. (Coord.). Métodos de pesquisa em fertilidade do solo. Brasília: Embrapa-SEA; 1991. Portuguese.
16. Kelling CRS, Reichardt K, Streck NA, Lago I, Zanon AJ, Rodrigues MA. Transpiração e crescimento foliar de crisântemo em função da fração de água transpirável no substrato. *Pesquisa Agropecuária Brasileira*. 2015; 50(9): 735-744. Portuguese.

17. Sinclair TR, Holbrook NM, Zwieniecki MA. Daily transpiration rates of woody species on drying soil. *Tree Physiol.* 2005; 25(11): 1469-1472.
18. Souza AT. Transpiração e crescimento foliar de genótipos de batata em função da fração de água transpirável do solo. Dissertação (Mestre em Engenharia Agrícola) – Universidade Federal de Santa Maria, Santa Maria. 2012.
19. Rodrigues RR, Pizetta SC, Reis EF dos, Ribeiro WR. Fração de água transpirável no solo no desenvolvimento inicial do cafeeiro conilon. *Coffee Science.* 2015; 10(3): 337-345. Portuguese.
20. Pizetta SC, Rodrigues RR, Ribeiro WR, Reis EF dos. Análise do crescimento do cafeeiro arábica, em relação à Fração de água transpirável do solo. *Coffee Science.* 2016; 11(1): 46-54. Portuguese.
21. Ray JD, Sinclair TR. Stomatal closure of maize hybrids in response to drying soil. *Crop Science.* 1997; 37(3): 803-807.
22. Araújo GL. Déficit hídrico no solo e sua influência sobre a transpiração, o crescimento e desenvolvimento do cafeeiro. Monografia (Graduação em Agronomia) - Universidade Federal do Espírito Santo, Alegre, 2011. Portuguese.
23. Larcher, W. *Ecofisiologia vegetal.* São Carlos: RIMA Artes e Textos. 2006. Portuguese.
24. Dardengo MCJD, Reis EF dos, Passos RR. Influence of the water readiness in the initial growth of the coffee plant conilon. *Bioscience Journal,* 2009; 25(6): 1-14.
25. Busato C, Reis EF dos, Martins CC, Pezzopane JEM. Lâminas de irrigação aplicadas ao café conilon na fase inicial de desenvolvimento. *Revista Ceres.* 2007; 54(314): 351-357. Portuguese.
26. Bray EA. Plant responses to water deficit. *Trends in Plant Science.* 1997; 2(2): 48-54.
27. Taiz L, Zeiger E. *Fisiologia vegetal.* 4th ed. Porto Alegre: Artmed; 2009. Portuguese.
28. Streck NA. Do we know how plants sense a drying soil? *Ciência Rural.* 2004; 34(2): 581-584. 33. Lecoeur J, Sinclair RT. Field pea transpiration and leaf growth in response to soil water deficits. *Crop Science.* 1996; 36(2): 331-335.
29. Araujo GL, Reis EF dos, Moraes WB, Garcia GO, Nazário AA. Influência do déficit hídrico no desenvolvimento inicial de duas cultivares de café conilon. *Irriga.* 2011; 16(2): 115-124. Portuguese.
30. Lecoeur J, Sinclair RT. Field pea transpiration and leaf growth in response to soil water deficits. *Crop Science.* 1996; 36(2): 331-335.
31. Davatgar N, Neishabouri MR, Sepaskhah AR, Soltani A. Physiological and morphological responses of rice (*Oryza sativa* L.) to varying water stress management strategies. *International Journal of Plant Production.* 2009; 3(4): p. 19-32.
32. Martins FB, Streck NA, Silva JC, Moraes WW, Susin F, Navroski MC, Vivian MA. Deficiência hídrica no solo e seu efeito sobre transpiração, crescimento e desenvolvimento de mudas de duas espécies de eucalipto. *Revista Brasileira de Ciência do Solo.* 2008; 32(3): 1297-1306. Portuguese.
33. Shao H, Chu L, Jaleel CA, Zhao C. Water-deficit stress-induced anatomical changes in higher plants. *Compens Rendus Biologies.* 2008; 331(3): 215-225.
34. Lago I, Streck NA, Bisognin DA, Souza AT de, Silva MR da. Transpiração e crescimento foliar de plantas de mandioca em resposta ao déficit hídrico no solo. *Pesquisa Agropecuária Brasileira.* 2011; 46(11): 1415-1423. Portuguese.