

Moisture behavior to the spatial variability of physical attributes in different soil management

ABSTRACT

The technique of temporal stability is used by several researchers to adequately represent the water content in the soil with reduced sampling effort from the identification of points on the field that reflect the average behavior of soil moisture. Therefore, the aim of this work was to analyze the moisture behavior to the spatial variation of density and particle size in a Hapludox under three different soil tillage systems (No-Till, Crop-Livestock Integration and Conventional Planting) in the municipality of Jataí, Southwest of Goiás, Brazil. It were collected nine sampling points information for each handling system in three depth ranges (0 - 0.12 m, 0.12 to 0.24 m, 0.24 to 0.36 m). For this, it was used a portable type sensor TDR HydroSence model, rods with length 0.12 m. The determination and location of points for data collection was done by georeferencing, using a GPS system ("Global Positioning System"), GARMIN, MAP785 model. Bulk density and soil particle size (clay) were determined according to the recommendation by EMBRAPA. The experimental area was composed of three sub-areas of approximately 1 ha each, cultivated with soybean (*Glycine max*) in the season 2013/2014 (first crop) with different soil management systems. Statistical analyzes were performed using analysis of variance and Pearson correction. The spatial variation was represented using the SURFER software. The results of variance analysis indicate statistical differences between values of bulk density and soil particle size between depths, between the different management and spatial distribution within each management. It was perceived spatial relationship between moisture and attributes studied for some depths and types of management, but there was little influence of physical attributes in the behavior of moisture to the soil and conditions studied.

Keywords: soil physical attributes; soil water content; soil tillage systems; surfer.

1. INTRODUCTION

Water is the main factor determining the productivity of crops. During the whole development, the plant absorbs water and loses using the ground as a reservoir of water and nutrients. The amount of water consumed by a crop during its development varies with the spatial and temporal distribution of climate, the variety of culture and management of the agricultural system, which modifies the physical properties of the soil directly related to storage of water [1].

It is noteworthy that due to the large spatial variability of physical and water attributes become often onerous costs sampling and analysis of soil to represent the average of any attributes.

Soils are naturally heterogeneous and their properties vary continuously in space and time. Among the factors responsible for the variability, we can mention the processes of training, involving the physical and chemical characteristics, interactions with the biological system and anthropogenic interference.

Soil has higher variability in their attributes, both vertically and horizontally, resulting from the interaction of the processes that govern the factors of their formation [2]. Knowledge of the distribution of physical water soil attributes becomes a basic requirement when seeking to establish appropriate management practices for soil and crops, for failure to comply with these concepts will result in errors in sampling and soil management. This stems from the large spatial variation of soil attributes and meaning and direction of water flows [3].

Over the years, some authors have studied the temporal stability of soil moisture and concluded that specific sites can represent the mean values of moisture in a given area of study over a period of time

and that the identification of these points is fundamental in planning of monitoring programs for hydrological and hydroagricultural variables, as it allows the reduction of costs related to the measurement of field data [4]; [5]; [6]; [7]; [8]; [9]; [10]; [11]; [12].

Therefore, the aim of this work was to analyze the response of moisture to the spatial variation of density and particle size in a Hapludox under three different managements (Tillage, Crop-Livestock Integration and Conventional Planting) in the municipality of Jataí, southwest of Goiás, Brazil.

2. MATERIAL AND METHODS

2.1 Experiment Location

The experiment was conducted in an experimental area at the Federal University of Goiás (UFG), Regional Jataí, in the southwest of the State of Goiás - Brazil (Figure 1), 17° 52' 53" S latitude and 51° 42' 52" W longitude, with 700 m of altitude, located in a climatic region type Cw, tropical savanna, mesothermal, with well-defined dry and rainy season, according to Kopen rating. The average annual rainfall varies around 1600 mm [13]. The studied soil was classified as Hapludox, according to the classification of EMBRAPA (2009) [14].

2.2 Experiment Area

The experimental area was composed of three sub-areas of approximately 1 ha each, cultivated with soybean (*Glycine max*) in the season 2013/2014 (first crop) with different soil management systems (Figure 1, b and c).

Subareas have been cultivated over the years with soybean in the first crop and maize (*Zea mays*) and sorghum (*Sorghum bicolor*) in the second (off-season) in no-tillage (NT) since the year 2008 (subarea 1) in crop-Livestock Integration system (ILP), consorting soybean crops and pasture since 2009 (subarea 2) and conventional tillage (CONV) associated with the use of harrowing at the time of planting. Each subfield is divided into a regular grid containing nine cells measuring 30 x 30 m each (Figure 1, b, and c) considering the center of each cell as the reference point for sampling.




Figure 1. Experiment location (State System of Geoinformation of Goiás - SGEI, 2014) (a); satellite image of the study area (Google Earth, 20/04/2014) (b); Sketching and distribution of the sampling points in the study areas (c). NT: No-tillage; ICL: integrated crop livestock; CT: conventional tillage.

2.3 Sampling and Data Collection

The determination and location of points for data collection was done by georeferencing, using a GPS system ("Global Positioning System"), GARMIN, MAP785 model.

On predetermined points, it was collected the straw (crop residues deposited on the soil surface) using an arc with internal area of 1.0 m², released randomly near the sampling point. Immediately after sampling the straw at each point, it was made monitoring of soil moisture. Moisture data were collected on nine points on each soil management system (NT, ICL and CT) at depths from 0 to 0.12, from 0.12 to 0.24 and 0.24 to 0.36 m, between the months of November 2013 and February 2014, with instantaneous readings every 14 days, by means of an equipment type TDR (time domain reflectometry). At the same points, deformed samples were collected for determination of soil organic matter (SOM). This portable Time Domain Reflectometry (TDR) type sensor, "HydroSence" (CD620, CS620), 0.12 m long rod manufactured by Campbell Scientific, provided instantaneous readings of soil

water content based on volume, providing monitoring of layers 0.12 m depth. Four repetitions were made, collected randomly within a radius of 1 m from the reference point.

For calibration of the TDR sensor, methodology similar to Abbas et al. (2011) [15] was adopted. 270 deformed soil samples were collected at five of the nine points of each management system, with 4 replicates per point (20 replicates per depth analyzed), collected randomly within a radius of 1 m from the point, at three depths.

With the predetermined points, it was held to monitor the soil moisture, as well as the removal of undisturbed soil samples collected in containers volumes known for determining the density and deformed sample for determining of soil particle size (clay), as recommended by EMBRAPA, (2009) [14].

3. RESULTS AND DISCUSSION

3.1 Area Characteristics

Although Schneider et al. (2008) [16] mention that the continuing existence of the pattern of the water content in the soil can be influenced by the topography, the relief of the areas used in the study (the gently rolling plan) significantly reduces the interference of relief in the spatiotemporal pattern of moisture soil in these areas, significantly reducing the influence of the topography in the temporal analysis of soil moisture (Figure 2).




Figure 2. Topography in the study areas. NT: No-tillage; ICL: integrated crop livestock; CT: conventional tillage.

As the topography, vegetation also had little influence in the comparative analysis of soil moisture between the three types of soil management (NT, ICL and CT), since the plantation was carried out with the same crop (soybean) and on the same date for both.

3.2 Soil Characteristics

It was determined the soil pH (H₂O), which showed values close to 5.47 for all conditions.

After completion of analysis of variance and test "Scott-Knott" the 5% significance, realized a statistical difference between the mean values of soil attributes evaluated in this study. Different letters placed after the average values of the attributes (Table 1) represent these differences.

The increased stability in CT probably occurred due to the frequent disturbance of the three depths under review. Higher values of soil density in layers close to the soil surface, in NT and ICL systems indicate a greater degree of compression in these layers (Table 1), which can be attributed to traffic historical of agricultural machinery in soil slightly upturned to over the years, confirming effects observed by Costa et al. (2003) [17] and Assis et al. (2005) [18]. In ICL, such compaction becomes more pronounced due to animal trampling [19].

Table 1. Organic matter content of the studied Latosol (ANOVA and test "Scott-Knott," $\alpha = 0.05$, $n = 9$)

Items	Parameter	NT-A	NT-B	NT-C	ICL-A	ICL-B	ICL-C	CT-A	CT-B	CT-C
d (g cm ⁻³)	Mean	1,26a	1,19b	1,17b	1,32a	1,22a	1,14b	1,24 ^a	1,26a	1,27a
	Max	1,44	1,37	1,32	1,38	1,37	1,25	1,37	1,39	1,38
	Min	1,09	0,99	1,02	1,14	1,09	0,99	1,07	1,14	1,09
	Σ	0,10	0,12	0,11	0,07	0,10	0,08	0,10	0,08	0,09
	CV	7,98	10,23	9,69	5,46	7,98	6,92	7,69	6,23	6,73
	Default E.	0,03	0,04	0,04	0,02	0,03	0,03	0,03	0,03	0,03
Clay (%)	Mean	45,1c	44,3c	48,2b	43,1c	46,8b	48,3b	57,6 ^a	58,0a	58,9a
	Max	49,32	50,16	52,74	48,59	50,48	53,12	64,85	63,19	63,84
	Min	42,19	36,31	44,21	35,12	39,96	44,19	52,06	47,83	54,16
	σ	2,83	4,64	2,74	4,42	3,36	3,10	4,87	5,37	3,78
	CV	6,27	10,46	5,68	10,25	7,17	6,43	8,45	9,25	6,42
	Default E.	0,94	1,55	0,91	1,05	1,12	1,03	1,62	1,79	1,26

d = bulk density; σ = standard deviation; CV = coefficient of variation; Default E. = standard error of the mean; n = sample size; Means followed by the same letter on the line, do not differ statistically; NT: tillage; ICL: integrated crop livestock; CT: conventional tillage; A = 0 to 0.12; B = 0.12 to 0.24 m; C = 0.24 to 0.36 m

Analyzing Table 1 and Figure 3, it is noted a tendency toward greater stability of density values along the soil profile in CT system, since the three layers of this system are equal to each other, while in other systems the first layers showed higher density values also represented by a greater homogeneity in color pattern between the three layers of CT system, compared to other systems.




Figure 3. Spatial distribution of density (g cm^{-3}) soil in three management systems and three depths.

Also in Table 1 and in Figure 4, it is possible to see similarities between the NT and ICL systems as the clay content in the three analyzed layers and growing trend with soil depth, unlike CT system, that is statistically different when presenting higher values of clay content at all depths evaluated compared to previous systems, with mean values close to 58%.




Figure 4. Spatial distribution of soil clay content in three management systems and three depths.

Similar to Gao & Shao (2012) [11], the correlation coefficient of "Pearson" was used to examine the dependence of the water content in the soil (WCS) in relation to the soil properties to NT, ICL and CT at three different depths (Table 2).

Table 2. Correlation coefficient of "Pearson" between soil moisture and variables investigated for the three types of management in different soil depths

Management		Depth	Bulk Density	Clay
System				
NT	A	0,29	- 0,14	
	B	- 0,24	- 0,16	
	C	- 0,23	0,22	
ICL	A	- 0,14	- 0,20	
	B	0,03	0,64**	
	C	- 0,52**	- 0,16	
CT	A	0,13	- 0,35	
	B	- 0,35	- 0,56**	
	C	- 0,79**	0,09	

** Significant to 1% (ANOVA); A = 0 to 0.12 m; B = 0.12 to 0.24 m; C = 0.24 to 0.36 m; NT: no-tillage; ICL: integrated crop livestock; CT: conventional tillage.

The physical attributes analyzed showed spatial relationship with moisture in two soil management systems (ICL and CT), with two events in each system. The increase in density caused a reduction of soil moisture in depth from 0.24 to 0.36 m for both, ICL and to CT. However, to the clay, there was both a positive (ICL-B) and negative (CT-B) relation. Note also that greater depths favor greater influence of these attributes on the variation of soil moisture in ICL and CT systems (Table 2).

4. CONCLUSION

Although there was some significance between the analyzed variables, it can be considered that there was little influence of physical attributes in the behavior of moisture to the studied soil and conditions.

REFERENCES

- MORETI, D; LIBARDI, PL; ROCHA, GC; LOVATTI, MJ; AGUIAR, LIG. Avaliação espaço-temporal das armazenagens gravimétrica e volumétrica da água num Latossolo com citros. *Revista Brasileira de Ciência do solo*, v.31, p. 1281-1290, 2007. Portuguese.
- CARVALHO, MAC DE; SORATTO, RP; ATHAYDE, MLF; ARF, O; SÁ, ME de. Produtividade do milho em sucessão a adubos verdes no sistema de plantio direto e convencional. *Pesquisa Agropecuária Brasileira*, Brasília, v.39, n.1, p.47-53, 2004. Portuguese.
- IQBAL, J. et al. Relationships between soil landscape and dryland cotton lint yield. *Soil Science Society American Journal*, Madison, v.69, n.6, p.872-882, 2005. English.
- GRAYSON, RB & WESTERN, AW. Towards areal estimation of soil water content from point measurements: Time and space persistence of mean response. *Journal of Hydrology*, 207:68-82, 1998. English.
- GÓMEZ-PLAZA, A; ALVAREZ-ROGEL, J. & ALBALADEJO, J. Spatial patterns and temporal stability of soil moisture across a range of scales in a semi-arid environment. *Hydrological Processes*, 14: 1261-1277, 2000. English.
- STARR, GC. Assessing temporal stability and spatial variability of soil water patterns with implications for precision water management. *Agric. Water Manage.* 72, 223–243. 2005. English.
- BROCCA, L; MORBIDELLI, R; MELONE, F; MORAMARCO, T. Soil moisture spatial variability in experimental areas of central Italy. *Journal of Hydrology* 333, 356–373, 2007. English.
- BROCCA, L; MELONE, F; MORAMARCO, T; MORBIDELLI, R. Soil moisture temporal stability over experimental areas in Central Italy. *Geoderma*, v. 148, n. 3-4, p. 364-374, 2009. English.
- ÁVILA, LF; MELLO, CR de; SILVA, AM da. Estabilidade temporal do conteúdo de água em três condições de uso do solo, em uma bacia hidrográfica da região da Serra da Mantiqueira, MG. *Revista Brasileira de Ciência do Solo*, 34: 2001-2009. 2010. Portuguese.
- SOUZA, ER de; MONTENEGRO, AADA; MONTENEGRO, SMG; MATOS, JDA de. Temporal stability of soil moisture in irrigated carrot crops in Northeast Brazil. *Agricultural Water Management* 99, 26–32. 2011. English.
- GAO, L; SHAO, M. Temporal stability of shallow soil water content for three adjacent transects on a hillslope. *Agricultural Water Management*, 110, 41-54. 2012. English.
- JIA, YH & SHAO, MA. Temporal stability of soil water storage under four types of revegetation on the northern Loess Plateau of China, *Agricultural Water Management*, 117, 33–42, 2013. English.
- KOETZ, M; MASCA, MGCC; CARNEIRO, LC; RAGAGNIN, VA; SENA JUNIOR, DG de; GOMES FILHO, RR. Caracterização agrônômica e °brix em frutos de tomate Industrial sob irrigação por gotejamento no sudoeste de Goiás. *Revista Brasileira de Agricultura Irrigada* v.4, n.1, p.14–22, 2010. Portuguese.
- EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária. Sistema brasileiro de classificação de solos. Rio de Janeiro: Centro Nacional de Pesquisa de Solos/Embrapa Solos, 412 p., 2009. Portuguese.
- ABBAS F; FARES, F; FARES, S. Field Calibrations of Soil Moisture Sensors in a Forested Watershed. *Sensors*, 11, 6354-6369, 2011. English.
- SCHNEIDER, K; HUISMAN, JA; BREUER, L; ZHAO, Y; FREDE, HG. Temporal stability of soil moisture in various semi-arid steppe ecosystems and its application in remote sensing. *Journal of Hydrology* 359:16-29, 2008. English.
- COSTA, FS; ALBUQUERQUE, JA; BAYER, C; FONTOURA, SMV; WOBETO, C. Propriedades físicas de um latossolo bruno afetadas pelos sistemas plantio direto e preparo convencional. *Revista Brasileira de Ciência do Solo*, 27: 527-535, 2003. Portuguese.

- 231 18. ASSIS, RL de; LANÇAS, KP. Avaliação dos atributos físicos de um Nitossolo vermelho
232 distroférico sob sistema plantio direto, preparo convencional e mata nativa. Revista Brasileira de
233 Ciência do Solo, 29: 515-522, 2005. Portuguese.
- 234 19. ALBUQUERQUE, JA; SANGOI, L; ENDER, M. Efeitos da integração lavoura-pecuária nas
235 propriedades físicas do solo e características da cultura do milho. Revista Brasileira de Ciência
236 do Solo, 25:717-723, 2001. Portuguese.