

PRODUCTION COMPONENTS OF THE COWPEA UNDER DIFFERENT DOSES OF ORGANIC FERTILIZER

ABSTRACT

The bean plant has a very high demand for nutrients, and as it has a short cycle, requires that the nutrients be readily available when needed, so as not to limit productivity. The use of organic fertiliser in beans is efficient, since due to the short cycle, the crop displays a satisfactory response to this type of fertilisation. The aim of this study was to evaluate the effects of organic fertiliser on the biometric parameters of the cowpea that reflect the productivity of bean plants. The experiment was carried out in the vegetable garden of the Federal University of Ceará, Pici Campus, and consisted of five treatments: control, mineral fertiliser, and organic fertiliser at doses of 100, 200 and 300% of the nitrogen recommendation for mineral fertilisation. At the time of harvest, 10 plants were collected from the working area of each plot to evaluate the number of pods per plant, grains per pod, 100-grain weight and mean pod length. The variables evaluated in the field experiment were submitted to statistical analysis using the SISVAR 5.3 software (System for the variance analysis of balanced data). The mean pod length, number of pods per plant, and 100-grain weight responded to the organic fertiliser relative to the mineral. Organic fertilisation is essential to achieve production components that indicate good crop yield. The absence of any nutrient can cause metabolic and nutritional disorders that prevent this aim from being reached. Organic fertiliser is associated with an improvement in the commercial quality of the beans.

Keywords: Organic fertilization; nutrient cycling; growth; bean; cowpea.

1. INTRODUCTION

High population density generates large amounts of organic waste throughout the food production and consumption chain. Among the possible forms of disposal of this waste, soil application is one of the most promising, since in addition to improving the chemical, physical and biological properties of the soil, it facilitates nutrient cycling in agricultural production, avoiding the consumption of non-renewable resources to meet the nutrient demand of agricultural crops [1].

Native to Africa, the cowpea (*Vigna unguiculata* L. Walp.) is cultivated in various regions of the world, particularly in India, West Africa, some regions of America, and in the north and northeast of Brazil [2]. In this region, it has an important socioeconomic, cultural and especially nutritional aspect, as it is a rich source of proteins, carbohydrates, vitamins and minerals [3].

According to [4], the bean plant has a very high demand for nutrients, and as it has a short cycle, requires that the nutrients be readily available when needed, so as not to limit productivity.

Research involving fertilisation with organic materials refers to the use of manure to promote improvements in the soil and to supply nutrients [5], and deals mainly with bovine, goat, pig and poultry manure [6];[7]. Even though the composition of these fertilisers is different, there is agreement that they are good for supplying nutrients, especially nitrogen, phosphorus and potassium.

The use of organic fertilisers in the bean is efficient, as it is a short-cycle crop with a superficial root system and has a satisfactory response to this type of fertilisation [8]. The productivity of organic bean production systems is comparable to conventional systems, showing that crop production with the use of organic fertilisers is viable [9]. The use of organic fertilisers has been effective in increasing productivity in the cowpea [10]; [11]; [12] and runner bean [13].

Numerous studies demonstrate the effects of organic fertiliser on different bean species. [14] achieved the highest mean yield for dry cowpea beans, of 2 Mg ha⁻¹ fertilised with a dose of 25 Mg ha⁻¹ cattle manure, in which there was an increase of 47.9 kg ha⁻¹ in green grain for each additional ton of organic fertiliser. In the runner bean, [15] found a linear increase in pod growth and a quadratic effect on maximum pod yield of 26, 30 and 23 t ha⁻¹ when adding a dose of 13, 24 and 16 t ha⁻¹ of chicken, cattle and goat manure respectively. Similarly, in the fava bean, [16] found that an increase in the doses of pig manure had a linear effect on productivity, both with and without mineral fertiliser.

The main objective of adding organic compost to the soil is not only the immediate availability of

nutrients to the plants, but also an improvement in soil structure, enabling mobilisation of the nutrient to the plant, and allowing for a more sustainable balance in the soil [17]. Another advantage of using compost is the gradual release of nutrients, where they are released in amounts that make it impossible to pollute the soil or poison the soil microbiota [18].

Organic fertilisers are able to provide essential nutrients to plants [19], which are indispensable for the productivity of legumes. [20] states that fertilising the soil with organic compost is fundamental to the formation and maintenance of soil fertility, since it affects various soil properties, such as the slow and gradual release of N, P and S. Therefore, the aim of this study was to evaluate the effect of organic fertiliser on the biometric parameters of the cowpea that reflect the productivity of bean plants.

49

50 2. MATERIAL AND METHODS

51

The field experiment was carried out in the vegetable garden of the Federal University of Ceará, Pici Campus (Fortaleza, Ceará) (Figure 1). The crop under evaluation was the cowpea (*Vigna unguiculata* (L.) Walp) 'Pitiúba', under a system of localised drip irrigation. The experimental design was of randomised blocks with five replications. The treatments comprised a standard treatment with mineral fertiliser (Treatment 1), treatment with no fertiliser (Treatment 2) and treatments with doses of organic compost equivalent to 100 (Treatment 3), 200 (Treatment 4) and 300% (Treatment 5) of the standard treatment.

The plots consisted of five rows spaced 100 cm apart with 30 cm between planting holes along the row, giving an area 5 m long and 1.8 m wide. Fertilisation in the standard treatment included N (urea), P (single superphosphate) and K (potassium chloride) in mineral form, and was defined based on an analysis of the soil (table 1) and on the crop requirements, as per the Recommendation and Liming Manual for the state of Ceará [21]. The mean temperature during the field experiment was $28 \pm 2^\circ\text{C}$ (Figure 2).

65

66 **Table 1 – Soil Analysis of experiment**

Characteristic	Value
Coarse Sand (g kg^{-1})	523
Thin Sand	370
Silt (g kg^{-1})	52
Clay (g kg^{-1})	55
Bulk Density (g cm^{-3})	1,46
Particle Density (g cm^{-3})	2,68
pH (água)	6,92
Calcium ($\text{mmol}_c \text{ dm}^{-3}$)	16,0
Magnesium ($\text{mmol}_c \text{ dm}^{-3}$)	13,0
Sodium ($\text{mmol}_c \text{ dm}^{-3}$)	3,0
Potassium ($\text{mmol}_c \text{ dm}^{-3}$)	1,0
Hydrogen + Aluminium ($\text{mmol}_c \text{ dm}^{-3}$)	18,2
Aluminium ($\text{mmol}_c \text{ dm}^{-3}$)	1,5
Carbon (g kg^{-1})	7,02
Nitrogen (g kg^{-1})	0,68
Soil Organic Matter (g kg^{-1})	12,10
P assimilável (mg dm^{-3})	23
C:N Ratio	10

67




Figure 1. Map of the area of the field experiment.




Figure 2. Temperature and rainfall data during the field experiment.

The dose of 100% in the treatment with organic fertiliser was defined based on the nitrogen equivalent of the standard treatment. Nutrient availability in the organic compost was determined from the equation proposed by [22] modified by [23], that takes into account the rate of nutrient mineralisation.

Planting holes, 0.25 m in depth and 0.25 m in diameter, were opened manually; the removed soil was mixed with the doses of organic and mineral fertilisers and returned to the holes. Organic fertilisation was carried out by incorporating the fertiliser into the soil in a single dose 10 days prior to sowing, since organic fertilisers tend to release nutrients gradually. The doses of N and K in the standard treatment with mineral fertiliser were applied to the planting hole in stages, with half applied together with the P when sowing and the remainder later at development stage V4, sub-stage V4-5, i.e. when the fifth trifoliolate leaf on the main stem was open in 50% of the plants [24]. Localised drip irrigation was continued throughout the experiment.

Sowing was by hand, using three seeds per hole. Thinning was later carried to leave two plants per hole, with the aim of obtaining a population of approximately 67,000 plants ha⁻¹. To control the weeds, two weedings between the rows were carried out with the help of a hoe, with one manual weeding in the holes. The cowpea aphid (*Aphis craccivora*) was controlled by spraying with a 4% solution of Ypê® Glycerin detergent with coconut [25]. Manual harvesting was carried out at the end of stage R9, i.e. at full maturation, giving a total biological cycle of 72 days. At the time of harvesting, 10 plants were collected from a predetermined location in the working area of each plot to evaluate the number of pods per plant (total number of pods divided by the number of plants), number of grains per pod (ratio between the total number of grains and the total number of pods), 100-grain weight (determined by randomly sampling and then weighing, on a precision balance to three decimal places, eight samples of 100 grains per plot, corrected for 13% wet basis) and mean pod length (determined by measuring with a millimetre rule 10 pods from each plot per replication).

The variables evaluated in the field experiment were submitted to statistical analysis using the SISVAR 5.3 software (System for the variance analysis of balanced data) described by [26]. Analysis of variance was performed between the five treatments, and regression analysis for the doses of organic compost and the contrast between the standard dose of mineral fertiliser and the doses of organic compost. Correlation analysis was applied to analyse the relationship between the production data and the nutrient content of the leaves and soil.

3. RESULTS AND DISCUSSION

The joint analysis of variance of the data for the yield components during the field experiment detected a significant effect on grain yield and its components, and all were influenced by the treatment (Table 2). No significant effects from the blocks were detected on the number of grains per pod, average pod weight or 100-seed weight.

Table 2 - Summary of the analysis of variance and coefficients of variation for mean pod length (MPL), number of grains per pod (NGP), number of pods per plant (NPP), mean pod weight (MPW), 100-grain weight (W100) and productivity in the cowpea.

SV	Mean square				
	MPL	NGP	NPP	MPW	W100
Treatment	7.93**	4.84**	52.63**	0.89**	8.09**
Block	3.76*	1.83 ^{ns}	11.16**	0.37 ^{ns}	0.43 ^{ns}
Error	1.44	2.03	3.16	0.25	0.42
CV (%)	5.21	7.70	11.21	13.62	4.16

Treatment	Mean				
	cm	-----um-----	-----g-----		
Mineral	23.30	18.56	17.7	3.78	16.10
0%	22.51 ⁽⁻⁾	18.18 ^{ns}	12.95 ⁽⁻⁾	3.58 ^{ns}	15.37 ⁽⁻⁾
100%	23.10 ^{ns}	18.32 ^{ns}	12.95 ⁽⁻⁾	3.59 ^{ns}	15.49 ⁽⁻⁾
200%	22.99 ^{ns}	18.50 ^{ns}	15.21 ^{ns}	3.90 ^{ns}	15.02 ⁽⁻⁾
300%	23.59 ^{ns}	19.00 ^{ns}	20.48 ^{ns}	3.72 ^{ns}	16.01 ^{ns}

SV = source of variation; CV (%) = Coefficient of variation; *significant at 5% probability by F-test; **significant at 1% probability by F-test. Mean values followed by (+) were higher than the absolute control at a level of 5% by Dunnett's test ($P \leq 0.05$). Mean values followed by (-) were lower than the absolute control at a level of 5% by Dunnett's test ($P \leq 0.05$). ^{ns} mean values do not differ from the absolute control at a level of 5% by Dunnett's test.

For mean pod length, the treatment corresponding to 300% organic fertiliser and the treatment of mineral fertiliser displayed the greatest pod lengths, 23.59 and 23.30 cm respectively, with no significant difference between them (Table 1). The control treatment (no fertiliser) had the lowest mean pod length, of 22.51 cm. [27], found mean values of 21.60 cm for the same variety as in the present study, demonstrating that the values found are close.

Mean pod length showed a linear response to the doses of organic fertiliser (Figure 3). Similar results were found by [15], according to whom increasing doses of chicken, cattle and goat manure resulted in a linear increase in pod length. [28] concluded that there was an increase in mean pod length in the fava bean when cattle manure was applied with and without NPK. [12] saw significant effects from the use of cattle manure on this production component in cowpea plants.

132




Figure 3. Increase in mean pod length for doses of organic fertiliser. Significant at 5% probability.

133

134

135

136

137 The MPL displayed a strong correlation with the number of grains per pod ($r=0.709$); the highest
138 number of grains per pod was seen with an increase in the mean pod length.

139 Although the analysis of variance indicated a significant response ($p < 0.05$) to the treatments for the
140 number of grains per pod (Table 1), there was no regression adjustment for the different treatments of
141 organic fertiliser, and no difference between the standard treatment of mineral fertiliser and the other
142 treatments. The data found were superior to those found by [27] and by [29], who found values of 16
143 and 17.59 grains per pod respectively. The number of grains per pod correlated moderately with
144 productivity ($r=0.639$) and had a strong correlation with mean pod length (0.709).

145 The SOM provides the macro- and micronutrients that are essential for plant development. Organic
146 materials in the soil gradually release nutrients, showing that the application of organic compost in
147 bean production is promising. Among the nutrients, the most important are sulphur and the
148 micronutrients zinc and boron, which are responsible for grain formation.

149 The mean number of pods per plant showed a quadratic increase for increasing doses of organic
150 fertiliser (Figure 4). The control (2) and the 100% fertiliser (3) displayed the lowest performance for
151 this variable, differing significantly from the treatment with mineral fertiliser (Table 1).

152




Figure 4. Increase in the number of pods per plant for different doses of organic fertiliser. Significant at 1% probability.

[30] found the highest number of pods per plant, 20.75 units, at a dose of 50 mL biofertiliser. Similar values were found in this study at the maximum organic fertiliser dose of 15696 t ha⁻¹ (Treatment 5). [14], when evaluating doses of cattle manure in the cowpea, found that there was an increase in pod yield with increases in the doses of manure. [31], evaluating cowpea genotypes without the addition of fertiliser, since the soil already contained the necessary nutrients, saw a yield of 15.4 pods per plant in the pitiúba cultivar, close to that found in this study with the dose of 200% organic fertiliser (Treatment 4). Alcântara *et al.* (2014) found 12.85 pods per plant in the pitiúba variety inoculated with strains of rhizobia.

For mean pod weight (MPW) as well as for NGP, although the analysis of variance indicated a significant response ($p < 0.01$) to the treatments (table 2) there was no regression adjustment for the different treatments of organic fertiliser, neither was there a difference between the standard treatment of mineral fertiliser and the other treatments. The variable displayed a moderate correlation with the 100-grain weight ($r=0.6049$), productivity ($r=0.5735$), number of pods per plant ($r=0.6382$) and mean pod length ($r=0.454$). According to [32], the presence of available essential nutrients at sufficient levels influences plant vigour and the morphological and metabolic processes that, as a consequence, increase the number of pods per plant and total productivity of the cowpea.

The maximum value found, 3.90 g, corresponding to a dose of 10464 t ha⁻¹ organic compost, was similar to the result found by [29]. [33], studying the cowpea, saw a greater increase in mean pod weight at a dose of 7.5 t ha⁻¹ cattle manure applied together with biofertiliser.

Treatments 1 (mineral) and 5 (300% organic fertiliser) had the highest mean values for 100-grain weight, not differing from each other and with higher values than in the control treatment (Table 1), demonstrating the positive effect of applying organic compost when growing cowpea. The values found in these treatments are similar to those obtained by [34], who found mean values of 16.17 g, and [29], who found a value of 16.59 g for the pitiúba variety. However, the values were lower than the value of 19.4 g found by [27] for the same variety as in the present study.

Several authors found no increase in the 100-grain weight for increases in fertiliser dose. [35], testing different doses of N and P in the common bean, found no significant results for this variable. [30] concluded that there was no significant effect on the cowpea from the addition of different doses of biofertiliser when compared to the control.

The 100-grain weight is a variable of extreme commercial importance, both domestically and internationally, where the varieties with a W100 greater than 20g are preferred [27]. The variety under study has this drawback, as it has smaller seeds and consequently less weight.

4. CONCLUSION

Organic fertiliser is essential to achieving production components that indicate good crop profitability. The absence of any of the nutrients can cause metabolic and nutritional disorders that prevent this goal from being achieved. Organic fertiliser is associated with an improvement in the commercial quality of the beans.

REFERENCES

- Blum J, Toma RS, Silva CMF. *Matéria Orgânica e Compostagem*. 1. ed. Fortaleza: Expressão Gráfica e Editora, 2016; v.1. 84p.
- Peksen A. Fresh Pod Yield and come Pod Characteristic of Cowpea (*Vigna unguiculata* L. Walp.) Genotypes from Turkey. **Asian Journal of Plant Sciences**, v.3, p. 269-273, 2004.
- Freire Filho FR, Ribeiro VQ, Rocha MM, Silva KJD, Nogueira MSR, Rodrigues EV. Feijão-caupi no Brasil: produção, melhoramento genético, avanços e desafios. Teresina: Embrapa Meio-Norte, 2011, 84 p.
- SILVA, C.C. da; SILVEIRA, P.M. Influência de sistemas agrícolas na resposta do feijoeiro (*Phaseolus vulgaris* L.) irrigado à adubação nitrogenada em cobertura. **Pesquisa Agropecuária Tropical**. 2000.
- VILLAS BOAS, R.L.; PASSOS, J.C.; FERNANDES, D.M.; BULL, L.T.; CEZAR, V.R.S.; GOTO, R. Efeitos de doses e tipos de compostos orgânicos na produção de alface em dois solos sob ambiente protegido. **Horticultura Brasileira** v. 22, p. 28-34, 2004.
- MALAVOLTA, E. **ABC da adubação**. São Paulo: Ceres, 1989. 250 p.
- CAVALCANTE, I.H.L.; ROCHA, L.F.; SILVA JÚNIOR, G.B.; AMARAL, F.H.C.; FALCÃO NETO, R. NOBREGA, J.C.A. Fertilizantes orgânicos para o cultivo da melancia em Bom Jesus- PI. **Revista Brasileira de Ciências Agrárias**, Recife – PE, v. 5, p. 518-524, 2010.
- PARRA, M.S. **Calagem e adubação**. In: PARRA, M.S. Feijão: tecnologia e produção. Londrina: IAPAR, 2000. p. 79-100.
- CARVALHO, W. P.; WANDERLEY, A. L. Avaliação de cultivares de feijão (*Phaseolus vulgaris*) para o plantio em sistema orgânico no Distrito Federal. **Ciência e Agrotecnologia**, Lavras, v. 31, n. 3, p. 605-611, 2007.
- CAVALCANTE, S.N.; DUTRA, K.O.G.; MEDEIROS, R.; LIMA, S.V.; SANTOS, J.R.; ANDRADE, R.; MESQUITA, E.F. Comportamento da produção do feijoeiro macassar (*Vigna unguiculata* L. Walp) em função de diferentes dosagens e concentrações de biofertilizante, **Revista de Biologia e Ciências da Terra**, v. 1, p. 10-14, 2009.
- MELO, R. F.; BRITO, L.T.L.; PEREIRA, L.A.; ANJOS, J.B. dos. Avaliação do uso de adubo orgânico nas culturas de milho e feijão caupi em barragem subterrânea. **Revista Brasileira de Agroecologia**, Porto Alegre, v. 4, n. 2, p. 1264-1267, 2009.
- PEREIRA, R.F.; CAVALCANTE, S.N.; LIMA, A.S.; MAIA FILHO, F.C.F.; SANTOS, J.G.R. Crescimento e rendimento de feijão vigna submetido à adubação orgânica. **Revista Verde**, Mossoró, v. 8, n. 3, p. 91-96, 2013.
- ALVES, E.U.; OLIVEIRA, A.P.; BRUNO, R.L.A.; ARAÚJO, E.; SILVA, J.A.L.; GONÇALVES, E.P.; COSTA, C.C. Produção de sementes de feijão- vagem em função de fontes e doses de matéria orgânica. **Horticultura Brasileira**, Brasília, DF, v. 18, n. 3, p. 215- 221, 2000.
- OLIVEIRA, A.P.; ARAUJO, J. S.; ALVES, E. U.; NORONHA, M. A. S.; CASSIMIRO, C. M.; MENDON, A. F. G. Rendimento de feijão-caupi cultivado com esterco bovino mineral. **Horticultura Brasileira**, v. 19, n. 1, p. 81-84, março, 2001.
- SANTOS, G.M.; OLIVEIRA, A.P.; SILVA J.A.L.; ALVES, E.U.; COSTA, C.C. Características e rendimento de vagem do feijão-vagem em função de fontes e doses de material orgânica. **Horticultura Brasileira**, Brasília, v. 19, n. 1, p. 30-35, março, 2001.
- ARAÚJO, J.S.; OLIVEIRA, A.P.; SILVA, J.A.L.; RAMALHO, C.I.; NETO, F.L.C. Rendimento do feijão-vagem cultivado com esterco suíno e adubação mineral. **Revista Ceres**, v. 48, p. 501-510, 2001.
- DE BERTOLDI, M; VALLINI, G.; PERA, A. The biology of composting: a review. *Waste Management and Resource*, vol. 1, n. 2, p. 157-176, 1983.
- CANELLAS, L.P.; OLIVARES, F.L.; AGUIAR, N.O.; JONES, D.L.; NEBIOSO, A.; MAZZEI, P.; PICCOLO, A. Humic and fulvic acids as biostimulants in horticulture. **Science horticulture**, v. 196, p. 15-27. 2015.

19. GILLER, M.V.; CARBALLO, M.T.; CALVO, L.F. Modelling N mineralization from bovine manure and sewage sludge composts. **Bioresource Technology**, v. 102, p. 863-871, 2011.
20. VAN RAIJ, B. **Fertilidade do solo e adubação**. Piracicaba, SP: Ceres: Potafós, 1991. 343 p.
21. AQUINO, A.B.; AQUINO, B.F.; HERNANDEZ, F.F.F.; HOLANDA, F.J.M.; FREIRE, J.M.; CRISÓSTOMO, L.A.; COSTA, R.I.; UNCHÔA, S.C.P.; FERNANDES, V.L.B. **Recomendações de adubação e calagem para o estado do Ceará**, UFC, 1993. 247 p.
22. FURTINI NETO, A. E.; VALE, F.R.; RESENDE, A.V.; GUILHERME, L.R.G.; GUEDES, G.A.A. **Fertilidade do solo**. Lavras: UFLA, 2001. 261 p.
23. SILVA, C. A. Uso de resíduos orgânicos na agricultura. In: SANTOS, G. A. (Ed.) *et al.* **Fundamentos da matéria orgânica do solo: ecossistemas tropicais e subtropicais**. 2ª edição. Porto Alegre: Metrópole, 2008, p. 597-624.
24. FERNÁNDEZ, F.; GEPTS, P.; LÓPEZ, M. Etapas de desarrollo en la planta de frijol. In: LÓPEZ, M.; FERNÁNDEZ, F.; SCHOONHOVEN, A. van. (eds.). **Frijol: investigación y producción**. Cali: CIAT, 1985. p.61-78.
25. LAVOR, M.T.F.C. Atividade biológica de produtos domissanitários para o controle alternativo do pulgão-preto no feijão-de-corda. 2006. 56 f. **Dissertação** (Mestrado em Agronomia/fitotecnia) – Centro de Ciências Agrárias, Universidade Federal do Ceará, Fortaleza, 2006.
26. FERREIRA, D.F. Sisvar: a computer statistical analysis system. **Ciência e Agrotecnologia**, v.35, n.6, p. 1038-1042, 2011.
27. PAIVA, J.B.; FREIRE FILHO, F. R.; TEÓFILO, E. M.; RIBEIRO, V. Q. **Feijão-caupi: melhoramento genético no Centro de Ciências Agrárias**. Fortaleza: Edições UFC, 2014. 261 p. il.
28. ALVES, A.U.; OLIVEIRA, A.P.; ALVES, A.U.; DORNELAS C.S.M.; ALVES, E.U.; CARDOSO, E.A.; OLIVEIRA, A.N.P.; CRUIZ, I.S. Lima beans production and economic revenue as function of organic and mineral fertilization. **Horticultura Brasileira**, v. 26, p. 251-254, 2008.
29. ALCÂNTARA, R.M.C.M.; XAVIER, G.R.; RUMJANEK, N.G.; ROCHA, M.M.; CARVALHO, J.S. Eficiência simbiótica de progenitores de cultivares brasileiras de feijão-caupi. **Revista Ciência Agrônoma**, v.45, n.1, p.1-9, 2014.
30. ALVES, G.S.; SANTOS, D.; SILVA, J.A.; MEDEIROS, J.A.; NASCIMENTO, L.F.C.; DANTAS, T.A.G. Estado nutricional do pimentão cultivado em solo tratado com diferentes tipos de biofertilizantes. **Acta Scientiarum. Agronomy**. v. 31, n. 4, p. 661-665, 2009.
31. FERREIRA, J.M.; SILVA, P.S.L. Produtividade de “feijão verde” e outras características de cultivares de caupi. **Pesquisa Agropecuária Brasileira**, Brasília, v.22, p. 55-58, janeiro, 1987.
32. CHATTERJEE, R.L.; BANDYOPADHYAY, S. Effect of boron, molybdenum and biofertilizers on growth and yield of cowpea (*Vigna unguiculata* L. Walp.) in acid soil of eastern Himalayan region. **Journal of the Saudi Society of Agricultural Science**, v. 16, p 332-336, 2017.
33. VIANA, J.S.; BRUNO, R.L.A.; ALMEIDA, F.A.C.; ARAÚJO, M.E.R.; CAVALCANTI, M.F.B.S.; SANTOS, J.F. Produção de feijão verde, cv. IPA 206, sob diferentes doses de esterco bovino e com biofertilizante. **Horticultura Brasileira**, Brasília, v. 23, n. 2, p.479, 2005.
34. OLIVEIRA, F.J.; ANUNCIAÇÃO FILHO, C.J.; BASTOS, G.Q.; REIS, O.V. Divergência genética entre cultivares de caupi. **Pesquisa Agropecuária brasileira**, Brasília, v. 38, n. 5, p. 605-611. 2003.
35. VIANA, T. O.; VIEIRA, N.M.B.; MOREIRA, G.B.L.; BATISTA, R.O.; CARVALHO, S.J.P. de; RODRIGUES, H.F.F. Adubação do feijoeiro cultivado no norte de Minas Gerais com nitrogênio e fósforo. **Revista Ceres**, Viçosa-MG, v. 58, n. 1, p. 115-120, jan/fev, 2011.