

Yield, quality and soil fertility as influenced by *rabic*castor (*Ricinus communis* L.) based intercropping system

Abstract

An experiment was conducted at C. P. College of Agriculture, Sardarkrushinagar on loamy sand soil using randomized block design with three replications during 2013-14. Results revealed that ~~S~~sole castor, castor + lucerne and castor + fenugreek (1:2) found equally effective with respect to seed yield of castor. Among the intercrops ~~viz-~~ lucerne, chicory, fenugreek and carrot, the maximum green forage yield of lucerne / chicory, seed yield of fenugreek and root yield of carrot were obtained when they were sown as sole crop. Similar trend was also observed in stalk / dry fodder / straw / green fodder yield of castor and intercrops. ~~R~~Rabicastor intercropped with lucerne, carrot, fenugreek and chicory at 1:2 row ratio produced significantly higher castor equivalent yield, 3297, 3057, 2731 and 3237 kg ha⁻¹, respectively over sole castor and other intercrops as sole. While intercropping of castor + fenugreek and castor + lucerne at 1:2 row ratio recorded significantly higher LER (Land equivalent ratio) value than sole crops and other intercropping systems. Oil content in castor / crude protein and crude ~~fiber~~ fibre content in lucerne and chicory / total soluble salts in carrot were found higher in their sole crop treatments as compared to intercrop. Oil yield was found significantly higher in sole castor than other treatments. Nutrient status of soil ~~viz-~~ organic carbon, available N, P, K and S after harvest of crops did not differ significantly due to different treatments.

Key words: Castor, Castor equivalent yield, intercropping, LER, row ratio and sole crop

INTRODUCTION

Castor (*Ricinus communis* L.) is a non-edible oilseed having high industrial importance due to presence of unique fatty acid and ricinoleic acid. Castor is extensively cultivated in India, China, Brazil, Ethiopia, Thailand ~~etc.~~ In India total acreage, production and productivity of castor crop during 2011-12 were 11.20 lakh hectares, 15.80 lakh tonnes and 1240 kg/ha⁻¹, respectively (SEA., 2013). The contribution of India in the world castor acreage and production are 56 and 84 percent, respectively. Area, production and

Formatted: English (United Kingdom)

Formatted: English (United Kingdom),
Superscript

Formatted: English (United Kingdom)

productivity of castor in Gujarat were 4.22 lakh ha, 8.33 lakh tonnes and 1972 kg ha⁻¹, respectively (SEA, 2013).

It is the major non-edible oil seed crop of India, which contributes 3-4 per cent to the total oilseeds production of the country. A significant feature of mixed/intercropping is that it is biologically more dynamic than a pure crop and is therefore less likely to succumb to vagaries of weather, for example reduction in yield of one component may be compensated by the other component. Thus, a mixer is intrinsically more secure and dependable in providing some major of returns than sole cropping. Intercropping is growing subsidiary crops between two rows of widely spaced main crops. The main objective of intercropping is to utilize the space left between two rows of main crop and natural resources as well as to produce more grain per unit area by better use of resources. This practice leads to some benefits like yield advantages as compared to sole cropping, greater stability of yield over different seasons, insurance against aberrant weather conditions, build-up or maintenance of soil fertility, economy of land, production of higher yield and higher economic returns per unit area in a given season.

The main principle involved in selecting intercrops is that they should not be competitive with main crop for soil moisture, nutrients and solar radiation. Considerable work has been done in selecting suitable crops like lucerne, chicory, fenugreek and carrot which were found suitable for intercropping in fennel but there is no any information available for taking these crops as intercrops in *rabi* castor. Shortages of fodder and vegetables in the country have focused the attention on including vegetables and fodder crops in intercropping systems which have capacity to get more return per unit area as well as to improve the physical, biological and chemical properties of soil. As the wide space is available between two paired rows in which profitable short duration crop can be grown during early growth stages of crop as *rabi* intercrop which gives an additional income also.

MATERIALS AND METHODS

A field experiment was conducted during *rabi* season of the year 2013 at Agronomy Instructional Farm, Chimanbhai Patel College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar, District Banaskantha (North Gujarat). The field experiment was laid out in a Randomized Block Design with three replications. An experiment consists of nine treatments including sole crops and intercropping systems in additive series viz., T₁ : Castor sole, T₂ : Lucerne sole, T₃ : Chicory sole, T₄ : Fenugreek sole,

Comment [WU1]: ?

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

T₅ : Carrot sole, T₆ : Castor + Lucerne (1:2), T₇ : Castor + Chicory (1:2), T₈ : Castor + Fenugreek (1:2), and T₉ : Castor + Carrot (1:2) were allotted to each plot by random method. The soil of experimental plot was loamy sand in texture, the initial soil status of organic carbon (0.18%), available N (148 kg ha⁻¹) is low, available P₂O₅ (47 kg ha⁻¹), available K₂O (284 kg ha⁻¹) is medium and available sulphur (9.56 ppm ha⁻¹). The organic carbon (%), available N (kg ha⁻¹), available P₂O₅ (kg ha⁻¹), available K₂O (kg ha⁻¹) and available sulphur (ppm ha⁻¹) were estimated by Walkley and Black's rapid titration method (Jackson, 1973), Alkaline permanganate method (Jackson, 1973), Olsen's method (Jackson, 1954), Flame photometer method (Jackson, 1973) and Extraction with 1% NaCl (William and Steinbergs, 1959), respectively.

Among the intercrops, legumes and fodder crops were chosen because of being widely used in common diet of livestock. Lucerne and chicory are the most important green forage crops by virtue of their excellent forage quality, flavour, taste and softness. Fenugreek and carrot are also eaten as raw vegetable and seed production. Full dose of P₂O₅ and half dose of N were applied as basal at the time of sowing of *rabi* castor and remaining half dose of N was applied in two equal splits at 40 and 70 days after sowing. In lucerne and fenugreek full dose of N and P₂O₅ was applied as basal. While in carrot and chicory, full dose of P₂O₅ and half dose of N was applied as basal at the time of sowing and remaining half dose of nitrogen was applied in two equal split at 35 and 60 days after sowing. The castor, lucerne and chicory were given 10 irrigations each of 50 mm depth whereas five irrigations were applied to carrot and fenugreek crops. Two common light irrigations viz., first irrigation at the time of castor sowing and second irrigation at intercrop sowing were given to each crop for good germination and establishment of seedling. The recommended dose of fertilizer of *rabi* castor: 80-25-0, lucerne: 20-40-0, chicory: 150-30-0, fenugreek: 20-40-0 and carrot: 80-60-0 N-P₂O₅-K₂O kg ha⁻¹.

Random seed samples were drawn from the produce of each net plot to estimate the oil content of castor seed. The oil content in castor seed samples was determined by IBM DC/20 series, NMR (Nuclear Magnetic Resonance) analyzer. Oil yield (kg ha⁻¹) was worked out by using following formula

$$\text{Oil yield (kg ha}^{-1}\text{)} = \frac{\text{Oil (\%)} \times \text{seed yield (kg ha}^{-1}\text{)}}{100}$$

The crude protein content and crude fibre content were measured with the help of INSTALAB 600 NIR product analyzer and it was expressed as percentage on dry weight basis (Nortvedt *et al.*, 1998).

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

RESULTS AND DISCUSSION

Castor equivalent yield (Table 1) was significantly higher with intercropping of castor with lucerne at 1:2 rows ratio than rest of the treatments but it remained at par with castor + carrot and castor + chicory in 1:2 row ratio, because of additional advantage of intercrop yields and higher yield of castor with intercrops recorded with component crops in these treatments. Additionally efficient utilization of natural resources by component crops having different rooting pattern, canopy depth and nutrient requirement in intercropping system resulted in the highest castor equivalent yield. These finding ~~were~~ are in conformity with those reported by Porwal *et al.* (2006) and Patel *et al.* (2009). Land equivalent ratio (LER) indicated that 38-60 per cent higher yield advantage were found in castor with carrot, chicory, lucerne and fenugreek at 1:2 row ratio over their sole crops (Table 1). Castor + fenugreek, castor + lucerne, and castor + chicory, were equally effective and found significantly superior over rest of the intercrops and their respective sole crop system. The higher yield of castor and land utilization in intercropping system recorded higher LER as compared to sole crop. This was due to extra yield obtained from intercrops and makes the combination of higher advantageous over sole crops due to development of better complementary relationship. These results ~~were~~ corroborated with the finding of Chandra and Sujatha (2000) and Basith and Mohammad (2013).

Seed and stalk yields (kg ha^{-1}) were recorded higher by sole castor than rest of the intercropping treatments, which could be attributed to ~~higher and optimum plant densities~~ no competition offered by intercrops in sole cropping system. Among the intercropping treatments, castor + lucerne (1:2) and castor + fenugreek (1:2) were found equally efficient to castor sole as lucerne and fenugreek crops have ability to fix atmospheric N and supply it to the associated castor crop and suppressed the weeds and higher sunshine availability to castor. Lower seed and stalk yields were noticed under castor intercropping with carrot and chicory at 1:2 row ratios (Table 1). This might be due to higher competition offered by intercrops for natural resources like space, plant nutrients, moisture and incoming sun radiation. The findings are in agreement with the results reported by Dhimmer (2009) and Singh (2009).

Green forage and dry fodder yields of lucerne and chicory, seed and straw yield of fenugreek and root and green forage yields of carrot were reduced in intercropping systems as

compared to their respective sole cropping (Table1). Such variation could be ascribed due to ~~increase in plant densities since these were grown as intercropped with castor and higher~~ competition for natural resources like soil moisture, plant nutrients, space and sunlight which are responsible for less photosynthesis rate resulted in lower accumulation of photosynthates in seed and dry matter per plant when grown as intercrop in comparison to sole crop. These findings are in close vicinity with the results of Srilatha *et al.* (2002).

The sole crop and intercropping systems exerted their non-significant effect on oil content of castor (Table 1), but marginally higher oil content was recorded with castor grown as sole crop as compared to different intercropping systems. The decrease in oil content of castor grown in intercropping system might be due to sharing of natural resources like space, plant nutrient, moisture and incoming sun radiation. Similar findings were recorded by Singh (2009) and Singhet *et al.* (2013).

Lower values of crude protein content in lucerne and chicory and total soluble salts in carrot were recorded when they were grown as intercrop with castor at 1:2 row ratio (Table 1). Moreover, higher crude fibre content of sole crops (lucerne and chicory) was ascribed as compared to their respective intercropping systems *i.e.* lucerne, chicory and carrot (Table 1). This might be due to higher competition offered by intra row spacing for natural resources like space, plant nutrients, moisture and incoming solar radiation.

The sole crop and intercropping systems did not exert their significant effect on soil available organic carbon, nitrogen, phosphorus, potassium and sulphur after harvest of the crops (Table 2). However, slight improvement in organic carbon of soil was found in castor + lucerne intercropping (1:2 row ratio) at the end of cropping as compared to initial nutrients status of soil. This might be due to the nitrogen fixing behaviour of lucerne (leguminous crop). Similar results for intercropping systems in mustard + lucerne were also reported by Patel *et al.* (2007) and Singhet *et al.* (2013) in sandy loam soils of north Gujarat. Available N, P and K content increased in soil after harvest of crops might be due to the complementary effect of lucerne and fenugreek throughout the crop growth period. These results are in conformity with findings of Patel *et al.* (2007) and Singh *et al.* (2013). However, growing of castor as a sole crop as well as castor + carrot as intercropping reduced the nutrient status of available N, P₂O₅ and K over the initial fertility status of the soil.

CONCLUSION

On the basis of one year study, it can be concluded that the highest seed yields (2315 kg ha⁻¹) and stalk yield (2616 kg ha⁻¹) of castor were recorded when ~~they were~~it was grown as sole crops as compared to **intercrops**. Castor seed equivalent yield (CEY) (3297 kg ha⁻¹) was significantly higher when it was intercropped with lucerne than rest of the treatments. Oil content of castor seed did not showed significant difference due to different sole and intercropping treatments. Oil yield (~~kg ha⁻¹~~) of castor seed was significantly higher when it was grown as sole and significantly at par when it intercropped with lucerne and **fenugreek** than rest of the treatments and castor grown in intercropped did not effect on nutrient status in soil after harvest of crop on loamy sand soil of North Gujarat Agro-climatic condition.

REFERENCES

- Basith MA, Mohammad S. Evaluation of productivity and profitability of castor-based intercropping systems. Indian Journal of Dryland Agriculture Research and Development. 2013;28(1):19-25.
- Chand P, Sujatha M. Castor based intercropping systems – ~~A~~a review. Agriculture Reviews. 2000;21(4):244-248.
- Dhimmar SK. Effect on growth and yield of *rabi* castor on pulses intercropping under varying planting geometry. American-~~European~~European Journal of Scientific Research. 2009;4(3):165-168.
- Jackson ML. **Soil eChemical Analysis** Prentice Hall of India Pvt. Ltd., New Delhi, ~~pp~~ 1954;pp. 183-192.
- Jackson ML. Soil Chemical Analysis, Prentice Hall of India Pvt. Ltd., New Delhi. 1973.
- Nortvedt R, Torrisen OJ, Tuene S. Application of near infrared transmittance spectroscopy in the determination of fat, protein and dry matter in Atlantic halibut fillet. Chemometrics and Intelligent Laboratory Systems. 1998;42(1-2):199-207.
- Patel GN, Patel MK, Patel RM, Patel KM, Pathak HC. Economics of inter/mixed cropping of lucerne (~~Medicago~~*Medicago sativa*) in mustard (*Brassica juncea* L.) in Northern Agro-Climatic zone of Gujarat. Journal of Oilseeds Research. 2007;24(2):340-34.
- Patel RM., Patel GN, Solanki SS. Inter-relay cropping of castor in greengram under irrigated condition. Journal of Oilseeds Research. 2009;26(2):168-169.

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

191 Porwal MK Agarwal SK, Khokhar AK. Effect of planting methods and intercrops on
192 productivity and economics of castor (*Ricinus communis*) based cropping systems.
193 Indian Journal of Agronomy. 2006;51(4):274-277.
194 SEA. Annual Report 2012-13, Castor Crop Survey, Solvent Extractors Association of India.
195 2013;2012-13.
196 Singh I. Study on intercropping of castor, *Ricinus communis* L. under irrigated condition.
197 Journal of Oilseeds Research. 2009;26(2):170-171.
198 Singh J, Patel BS, Rathode BS. Feasibility of inter/mixed cropping system of mustard
199 {*Brassica juncea* (L.) Czern & Coss} with lucerne (~~*Medicago*~~ *Medicago sativa*).
200 Haryana Journal Agronomy. 2013;29(1&2):67-71.
201 Srilatha AN, Masthan SC, Mohammed S. Production potential of castor intercropping with
202 legumes under rainfed conditions. Journal of Oilseeds Research 2002;19(1):127-
203 128.
204 William DH, Steinberg A. Soil sulphur fraction as chemical index of available sulphur in
205 some Australian soil. Australian Journal of Agronomy Research. 1959;10(1):340-
206 352.

Formatted: English (United Kingdom)

Formatted: Font: Not Italic, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Table 1: Yield and quality of castor and inter crops as influence by different treatments

Treatments	Seed yield / green forage yield/ root yield (kg ha ⁻¹)	Stalk yield / dry or green fodder yield / straw yield (kg ha ⁻¹)	Castor equivalent yield (kg ha ⁻¹)	Land equivalent ratio (LER)	Oil content (%) in castor seed	Oil yield (kg ha ⁻¹)	Crude protein content/ total soluble salts (%) in carrot root	Crude fibre content (%)
Castor sole	2315	2616	2315	1.00	48.70	1127	-	-
Lucerne sole	19800	3400	1803	1.00	-	-	18.65	20.6
Chicory sole	30300	5419	1763	1.00	-	-	20.41	10.7
Fenugreek sole	816	1353	823	1.00	-	-	-	-
Carrot sole	16518	18704	2284	1.00	-	-	12.60	2.35
Castor + Lucerne (1:2)	2267 (10900)	2539 (1879)	3297	1.54	48.14	1091	18.13	20.4
Castor + Chicory (1:2)	1974 (18280)	2171 (3140)	3057	1.46	47.85	945	20.22	10.5
Castor + Fenugreek (1:2)	2154 (534)	2391 (958)	2731	1.60	48.05	1035	-	-

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Font: Font color: Auto, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted Table

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted ... [1]

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted ... [2]

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted ... [3]

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted ... [4]

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted ... [5]

Castor + Carrot (1:2)	1761 (9783)	1919 (15163)	3237	1.38	47.32	833	12.25	2.20
S.E.m.±	-	-	110	0.06	1.55	521.61	-	-
C.D. (0.05)	-	-	321	0.17	NS	158.54	-	-

*Data presented in parenthesis indicates intercrops values

Table 2: Effect of different treatments on residual soil fertility after harvest of the crops

Treatments	Nutrient status in soil after harvest				
	Organic carbon (%)	Available nitrogen (kg ha ⁻¹)	Available phosphorus (kg ha ⁻¹)	Available potash (kg ha ⁻¹)	Available sulphur (ppm)
Castor sole	0.19	141.45	45.68	280.64	10.23
Lucerne sole	0.21	158.45	51.26	286.42	10.09
Chicory sole	0.20	151.60	48.50	279.68	9.94
Fenugreek sole	0.20	155.26	49.71	282.34	10.57
Carrot sole	0.19	142.30	46.35	280.12	9.85
Castor+ Lucerne (1:2)	0.23	152.56	48.62	286.40	9.88
Castor+ Chicory (1:2)	0.22	151.85	46.21	284.20	10.13

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Font: Not Bold

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

▲ Castor+ Fenugreek (1:2)	0.21	148.62	46.89	283.61	9.95
▲ Castor+ Carrot (1:2)	0.20	142.17	44.15	276.56	9.82
▲ S.Em.±	0.008	4.13	1.60	7.57	0.35
▲ C.D. (0.05)	NS	NS	NS	NS	NS
▲ Initial soil nutrient status	0.17	143.50	47.50	282.00	9.23
▲					

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: Font: Not Bold, English (United Kingdom)

Formatted: English (United Kingdom)

Formatted: English (United Kingdom)

