

Original Research Article

Endophytic fungi associated with the crown-of-friar cactus (*Melocactus zehntneri* Britton & Rose) in the semiarid region of Brazil

ABSTRACT

Aims: The objective of this study was to identify the occurrence of endophytic fungi associated with the crown-of-friar cactus (*Melocactus zehntneri* Britton & Rose) in the Brazilian Semi-arid.

Methodology: The plants were collected in the municipality of Taperoá, State of Paraíba, Brazil and sent to the Phytopathology Laboratory, belonging to the Department of Plant Science and Environmental Sciences, Federal University of Paraíba in the city of Areia, Paraíba, Brazil. Based on the indirect isolation of the fragments of symptomatic tissues, fungal structures which were examined under optical microscopy and compared with specialized literature. It was carried out the removal of vegetative and reproductive structures of fungi cultivated in BDA medium at a temperature of 25 ± 2 ° C and a photoperiod of 12 hours of light.

Results: Fungal genera were identified *Colletotrichum* sp., *Curvularia* sp., *Fusarium* sp., *Pestalotopsis* sp. e *Phomopsis* sp. in plants of de crown-of-friar cactus collected in the state of Paraíba, these genera cause diverse symptomatology in this cactus, negatively affecting the crown, base and root of the plant.

Conclusion: The fungal genera *Colletotrichum*, *Curvularia*, *Fusarium*, *Pestalotopsis* and *Phomopsis* were found on *M. zehntneri* plants collected in the state of Paraíba, Brazil.

Keywords: Cactacea; pathogens; sanity

1. INTRODUCTION

The *Caatinga* biome dominates the semiarid region of northeastern Brazil and is characterized as a complex set of vegetal physiognomies, with an ample number of arboreal, arbustive and herbaceous species (MORO *et al.*, 2015). However, phytosanitation issues have been studied little in this biome and there is a need for strategic planning to avoid the loss of biodiversity due to the high degree of human interference in the form of agriculture, livestock farming and extractive activities (GONÇALVES *et al.*, 2016).

Cacti play a fundamental role in the ecology of the *Caatinga*. The genus *Melocactus* spp. has 37 xerophilic species that occur from Mexico to the state of Rio de Janeiro, with the highest

concentration found in northeastern region of Brazil, which is considered the center of primary diversity for the genus (MENEZES *et al.*, 2015). The *Caatinga* biome is home to several species of *Melocactus*. *M. zehntneri*, which is commonly known as the crown-of-friar cactus or Turk's cap cactus, is widely used by the local population in cooking, folk medicine (CORREIA *et al.*, 2018), decorative landscaping and as fodder (LOPES *et al.*, 2012).

Like *M. zehntneri*, other cacti are infected by fungal pathogens, such as the prickly pear (*Opuntia ficus-indica*), on which 15 fungal genera have been found, including *Fusarium*, *Colletotrichum* and *Pestalotiopsis* (SOUZA *et al.*, 2017). Few studies have been conducted on the etiology and epidemiology of diseases affecting the crown-of-friar cactus caused by pathogenic fungi. Therefore, the aim of the present paper was to identify the occurrence of endophytic fungi associated with the species.

2. MATERIAL AND METHODS

Plants were collected from the municipality of Taperoá, which is located in the central region of the state of Paraíba, Brazil, in the micro-region known as the Western Cariri (coordinates: 739,959EW and 9,202,794NS-MC-39) at an altitude of 532 meters. The climate is rainy tropical with a dry summer. The rainy season begins in January/February and ends in September, but can extend into October. After collection, the material was sent to the Phytopathology Laboratory for the Phytotechnic and Environmental Science Department of the Federal University of Paraíba.

M. zehntneri plants with typical symptoms of infection were cleaned and kept at room temperature ($25 \pm 2^{\circ}\text{C}$). Tissue fragments were sampled with the aid of a sterile surgical scalpel, with the removal of part of the infected tissue from the crown, base and root of the plant. The fragments were plated in potato-dextrose-agar (PDA) medium at 25°C with a 12-hour photoperiod in biochemical oxygen demand (BOD).

After seven days of incubation, the identification of fungi was performed beginning with the spatial separation of the pathogen promoted by the host tissues with the aid of an optic stereomicroscope. The structures were compared to descriptions found in the literature (SEIFERT *et al.*, 2011) for the determination of the occurrence of fungi associated with *M. zehntneri*.

3. RESULTS AND DISCUSSION

Under field conditions, necrosis of a dark coloration and yellowish spots were found in the regions of the crown, base and root of infected plants, with the consequent shedding of spines, which serve as a defense mechanism against animal herbivory (Figure 1).



Fig. 1. crown-of-friar cactus plants (*Melocactus zehntneri* Britton & Rose), collected in the municipality of Taperoá, State of Paraíba, showing symptoms of aculeous fall (thorns) and foraging of plants by animals.

According to Soares et al. (2008), environmental factors play a fundamental role in the epidemiological development of diseases and the predominance of high temperatures combined with low humidity throughout a large part of the year favors a number of diseases and their causal agents. Freire (2009) identified more than ten diseases caused by pathogens on the mandacaru cactus (*Cereus jamacaru*), particularly rot caused by *Lasiodiplodia theobromae*, which initiates infection at the apex of the plants, causing the darkening of infected tissues, followed by rot. The author stresses that, despite its apparent rusticity, the mandacaru and other native cacti of the *Caatinga* can be infected by numerous pathogens.

Fungal spores from *Colletotrichum* sp., *Pestalotopsis* sp. and *Fusarium* sp. were predominant in different regions of the crown, base and root of the plant (Figure 2A, B, C and D). It is possible that

some of these structures are able to cover injured areas, which is normal for the genus *Colletotrichum*.




Fig. 2. Fungus conidia associated with crown-of-friar cactus (*Melocactus zehntneri* Britton & Rose), where (A) *Colletotrichum* sp. (B) *Pestalotiopsis* sp. (C) *Fusarium* sp. (D) *Colletotrichum* sp.

The dominance of these fungi reduced the diversity of microbial agents. According to Suryanarayanan, Wittlinger and Faeth (2005), this is a common occurrence when plants in dry zones of the world are evaluated and is due mainly to environmental factors, such as low precipitation and low vegetation density. However, the findings diverge from data reported by Bezerra et al. (2012), who report high endophyte richness on different cactus species in the dry tropical Brazilian forest, attributing the findings to the high diversity of plants in the *Caatinga*.

Besides *Colletotrichum*, *Pestalotiopsis* and *Fusarium*, the genera *Curvularia* and *Phomopsis* were also identified. Gonçalves, Freire and Lima (2013) report the presence of *Colletotrichum* sp., *Fusarium* sp., *Nodulisporium* sp. and *Phomopsis* sp. colonizing mandacaru and the xique-xique (*Pilosocereus gounellei*) under natural conditions of the *Caatinga*.

The microflora was more intense in the regions of the crown and base of the plant, with a predominance of the genera *Fusarium* and *Pestalotiopsis*. However, only *Fusarium* was common to the infected regions of the crown, base and root. The infected area of the plant acquires a soft rot appearance and yellowish coloration, causing an exudate denominated gummosis and the shedding of spines, such that these areas become defenseless against animal herbivory. In areas where extensive livestock farming predominates, infected plants can suffer irreversible damage due to the foraging of animals, especially in periods of drought, when the herbaceous and arbustive strata become scarcer.

The genus *Colletotrichum* was found in regions of the crown and roots of the plants. This fungus causes the darkening of the organs of the plant and photosynthesizing area and the injured area is subsequently covered with a mass of dark spores. After a few days, the injury increases and the plant tissue becomes hard and dry (FREIRE et al., 2015). Barbosa et al. (2012) report that this pathogen causes injury to the surface of the cladodes in the prickly pear cactus, with a tendency to concentrate at the edges in the form of depressed dark spots and the emergence of fructifications (acervuli) in the injured area in the form of salient black dots.

4. CONCLUSION

The fungal genera *Colletotrichum*, *Curvularia*, *Fusarium*, *Pestalotopsis* and *Phomopsis* were found on *M. zehntneri* plants collected in the state of Paraíba, Brazil. These genera cause diverse symptoms in cacti, which underscores the need for further studies on the potential effect of these pathogens on *M. zehntneri*.

COMPETING INTERESTS

The authors declare that there is no conflict of interest.

REFERENCES

- Alfenas AC. Isolation of phytopathogenic fungi. In: Alfenas  Mafia RG, editors. Methods in phytopathology. 3rd ed. Viçosa: Editora UFV, 2016.
- Barbosa SR, Cavalcanti VALB, Lopes EB, Araújo E. Forage palm diseases, In: Lopes EB, Forage Palm: Cultivation, Current Usage and Perspective of Use in the Northeastern Semiarid. 2nd ed. About Us |.
- Bezerra JDP, Lopes DHG, Santos MGS, Svedese VM, Paiva LM, Almeida-Cortez JS, Souza-Motta CM. Wealth of endophytic micro-organisms in species of family Cactaceae. Bulletin of the Latin American and Caribbean Society of Cactaceae and other Succulents. 2012; 9 (2): 19-23.
- Correia D, Nascimento EHS, Gomes Filho A, Abelardo H, Lima MLB, Almeida JVF. Melocactus. 1 ed. Fortaleza: Embrapa Tropical Agroindustry; 2018.
- Freire FCO. Pathogens Associated with Mandacaru (*Cereus jamacaru* Dc.) In  State of Ceará. Technical Communiqué 148. 1 ed. Fortaleza: Embrapa Tropical Agroindustry; 2009.
- Freire KTLS, Araújo GR, Bezerra JDP, Barbosa RN, Silva DCV, Svedese VM, Paiva LM, Souza-Motta CM. Endophytic fungi of *Opuntia ficus-indica* (L.) Mill. (Cactaceae) healthy and infested by *Dactylopius opuntiae* (Cockerell, 1896) (Hemiptera: Dactylopiidae). Gaia Scientia. 2015; 9 (2): 104-110.
- Gonçalves FJT, Freire FCO, Lima JS. Endophytic fungi and their potential as producers of bioactive compounds. Essentia. 2013; 15 (1): 71-92.
- Gonçalves FJT, Freire FCO, Lima JS, Melo JGM, Câmara MPS. Pathogenicity of Endophytic Botryosphaeriaceae species of plants from the Caatinga of the state of Ceará in mango and umbu-cajá. Summa Phytopathol. 2016; 42 (1): 43-52.
- doi: <http://dx.doi.org/10.1590/0100-5405/2099>.
- Lopes EB. Forage palm: cultivation, current use and prospects of use in the Northeastern Semi-arid. 1st ed. João Pessoa, PB: Emepa / Faepa; 2012.
- Menezes  Ribeiro-Silva S. Cactaceae of Ceará, Brazil: Priorities for conservation. Gaia Scientia. 2015; 9 (2): 67-76.

doi: <https://doi.org/10.21707/gv.v9i2.24166>.

Moro MF, Macedo MB, Moura-Fé MM, Castro ASF, Costa RC. Vegetation, phytoecological units and landscape diversity in the state of Ceará. *Rodriguésia*. 2015; 66 (3): 717-743.

Soares AR. Infection of guavas by *Colletotrichum gloeosporioides* and *Colletotrichum acutatum* under different temperatures and periods of wetting. *Tropical Plant Pathology*. 2008; 33 (4): 265-272.

Seifert K, Morgan-Jones G, Gams W, Kendrick B. The Genera of Hyphomycetes. 1 ed. Netherlands: CBS-KNAW Fungal Biodiversity Centre Utrecht; 2011.

Souza AEF, Nascimento LC, Souza BO. Principal components of the intensity of squamous rot on prickly pear plantations in the Semiarid region of the state of Paraíba, Brazil. *Revista. Caatinga*. 2017; 30 (2): 370–376.

doi: <http://dx.doi.org/10.1590/1983-21252017v30n212rc>.

Suryanarayanan TS, Wittlinger SK, Faeth SH. Endophytic fungi associated with cacti in Arizona. *Mycological Research*. 2005; 109 (1): 635–639.

doi: <https://doi.org/10.1017/S0953756205002753>.