

ROLE OF ZINC IN PLANT NUTRITION- A REVIEW

Hafeez, B¹., Khanif, Y. M²., Saleem, M^{1*}

¹Agriculture Research Institute Tandojam-Pakistan

²Department of Land Management, Faculty of Agriculture, University Putra Malaysia

ABSTRACT

Zinc is plant micronutrient which is involved in many physiological functions its inadequate supply will reduce crop yields. Zinc deficiency is the most wide spread micronutrient deficiency problem, almost all crops and calcareous, sandy soils, peat soils, and soils with high phosphorus and silicon are expected to be deficient. Zinc deficiencies can affect plant by stunting its growth, decreasing number of tillers, chlorosis and smaller leaves, increasing crop maturity period, spikelet sterility and inferior quality of harvested products. Beside its role in crop production Zn plays a part in the basic roles of cellular functions in all living organisms and is involved in improving the human immune system, due to its insufficient intake, human body will suffer from hair and memory loss, skin problems and weakness in body muscles.

Key words: Zinc, Plant nutrition, Rice, Bioavailability, Human health

INTRODUCTION

Zinc is essential for the growth in animals, human beings, and plants it is vital to the crop nutrition as required in various enzymatic reactions, metabolic processes, and oxidation-reduction reactions. In addition, Zn is also essential for many enzymes which are needed for nitrogen metabolism, energy transfer and protein synthesis. Zinc deficiency not only retards growth and yield of plants, but it also has effects on human beings. More than 3 billion people worldwide are suffering from Fe and Zn deficiencies, and this condition is particularly widespread in areas where population is heavily dependent on an unvaried diet of cereal-based foods, in which Fe and Zn are stored almost exclusively in the husk, and are therefore lost during milling and polishing (Cakmak, 2002; Graham *et al.*, 2001).

1. Importance of Zn in Humans

Zinc deficiency is common in humans, animals and plants. More than 30% world's population suffers from Zn deficiency (Welch *et al.*, 2002). Zinc plays a part in the basic roles of cellular functions in all living organisms and is also involved in improving the human immune system. The optimum dietary intake for human adults is 15 mg Zn per day. Zinc acts as a catalytic or structural component in various body enzymes.

Unsatisfactory intake and improper absorption of Zn in the body may cause deficiency of Zn. Due to Zn deficiency; the human body will suffer from hair and memory loss, skin problems and weakness in body muscles. Further insufficient intake of Zn during pregnancy also causes stunted brain development of the fetus. Infertility has also been observed in Zn deficient men. Zinc deficiency may cause congenital diseases like Acrodermatitis enteropathica (Lukaski, 2004; Morley, 2004 and Zimmermann, 2001).

As per recommendations of the FAO/WHO (2002) and Hotz and Brown (2004) an average male need 11 mg of Zn daily while an average female needs 9 mg of Zn. During pregnancy

*Corresponding author E-mail: sarkisaleem@yahoo.com.

and lactation, the female needs 13 mg to 14 mg of Zn daily. Infants from 7 months to 3 years need 3 mg, 4 to 8 years need 5 mg and children from 9 to 13 years need 8 mg of Zn daily. Cakmak, 2002 reported that Zn is stored in the rice husks and grains and with the consumption of this cereal human zinc deficiency can be decreased. The foods rich in Zn are beef, pork, chicken, and breakfast cereals, nuts like roasted peanuts, almonds, walnuts, oats and dairy products like yogurt, cheese and milk.

2. Role of Zinc in Plants

The Zn plays very important role in plant metabolism by influencing the activities of hydrogenase and carbonic anhydrase, stabilization of ribosomal fractions and synthesis of cytochrome (Tisdale et al., 1984). Plant enzymes activated by Zn are involved in carbohydrate metabolism, maintenance of the integrity of cellular membranes, protein synthesis,, regulation of auxin synthesis and pollen formation (Marschner 1995). The regulation and maintenance of the gene expression required for the tolerance of environmental stresses in plants are Zn dependent (Cakmak 2000). Its deficiency results in the development of abnormalities in plants which become visible as deficiency symptoms such as stunted growth, chlorosis and smaller leaves, spikelet sterility. Micronutrient Zn deficiency can also adversely affect the quality of harvested products; plants susceptibility to injury by high light or temperature intensity and to infection by fungal diseases can also increase (Marschner 1995; Cakmak 2000). Zinc seems to affect the capacity for water uptake and transport in plants and also reduce the adverse effects of short periods of heat and salt stress (Kasim, 2007; Disante et al., 2010, Peck and McDonald, 2010, Tavallali et al., 2010). As Zn is required for the synthesis of tryptophan which is a precursor of IAA, it also has an active role in the production of an essential growth hormone auxin, (Alloway, 2004, Brennan, 2005). The Zn is required for integrity of cellular membranes to preserve the structural orientation of macromolecules and ion transport systems. Its interaction with phospholipids and sulphhydryl groups of membrane proteins contributes for the maintenance of membranes (Cakmak, 2000; Kabata-Pendias and Pendias, 2001; Alloway, 2004; Dang et al., 2010; Disante et al., 2010).

The Zn-finger transcription factors are involved in the development and function of floral tissues such as anthers, tapetum, pollen and pistil secretory tissues in many plant species, it is likely that VvZIP3 may play a key role in both flower and normal fruit development (Kobayashi et al; 1998, Sharma et al;1987). The alteration of VvZIP3 expression during flowering and fertilization can modify the distribution and availability of Zn, hence affects normal reproductive development (Kapoor et al; 2002).

3. Zinc Deficiency in Soil

Zinc deficiency can be found in every part of the world and almost all crops respond positively to application of Zn (Alloway, 2008). Normal soils inherit their trace elements which include Zn primarily from the rocks through geochemical and pedochemical weathering processes. Besides mineralogical composition of the parent material, the total amount of Zn present in the soil is also dependent on the type, intensity of weathering, climate and numerous other predominating factors during the process of soil formation (Saeed and Fox, 1977). Meanwhile, high pH and high contents of CaCO_3 , organic matter, clay and phosphate can fix Zn in the soil and give rise to the reduction of available Zn (Imtiaz, 1999). Soils derived from granite and gneiss can be low in total Zn (Krauskopf, 1972). Similarly, total Zn is low in highly leached, acid, sandy soils such as the ones found in many coastal areas. Quartz in the soil dilutes Zn from it because the reported concentrations of Zn in quartz are very low, which ranged from $1.0 \mu\text{g g}^{-1}$ to < 5 to $8 \mu\text{g g}^{-1}$ (Helmke *et al.*, 1977 and Brehler and Wedepohl, 1978). According to the Food and Agriculture Organization (FAO), about 30% of the cultivable soils of the world contain low levels of plant available Zn (Sillanpaa, 1990). The lowest Zn concentrations were always found in Spodosols ($28 \mu\text{g g}^{-1}$) and luvisols ($35 \mu\text{g g}^{-1}$), while higher levels were found in fluvisols ($60 \mu\text{g g}^{-1}$) and Histosols ($58 \mu\text{g g}^{-1}$) (Kiekens, 1995).

92 Generally, Zn deficiency is expected in calcareous soils, sandy soils, peat soils, and soils
93 with high phosphorus and silicon (Alloway, 2008, 2004). The submerged soils are well
94 recognized for the lack of Zn availability to the plants; particularly due to the reaction of Zn
95 with free sulphide (Mikkelsen and Shiou, 1977). Flooding and submergence bring about a
96 decline in available Zn because of the changes in pH value and the formation of insoluble Zn
97 compounds. Meanwhile, the insoluble Zn compounds formed are likely to be with Mn and Fe
98 hydroxides from the breakdown of oxides and adsorption on carbonates, specifically
99 magnesium carbonate. Under the submerged conditions for rice cultivation, Zn is
100 transformed into amorphous sesquioxide precipitates or franklinite; ZnFe_2O_4 (Sajwan and
101 Lindsay, 1988). Zinc deficiency causes multiple symptoms which usually appear 2 to 3
102 weeks after transplanting of rice seedlings, with leaves developing brown blotches and
103 streaks that may fuse to entirely cover older leaves, and plants remain stunted, whereas in
104 severe cases, the plants may die, while those which recover will show substantial delay in
105 maturity and reduction in yield (Yoshida and Tanka, 1969; Van Breemen and Castro, 1980
106 and Neue and Lantin, 1994).

107 **4. Zinc under Submerged Soil Conditions**

108 Zn deficiency is very common under flooded soil conditions. In acidic soils, Zn is precipitated
109 as $\text{Zn}(\text{OH})_2$ and as ZnS in sulfur-rich and alkaline soils. The availability and solubility of Zn
110 decreases while pH increases. The oxides of Mn and Zn along with CaCO_3 or MgCO_3 are
111 strongly absorbed by Zn under submerged condition. Whereas in calcareous soils, HCO_3^- is
112 the predominant anion, which mainly reduces Zn transport from root to shoot, but not so
113 much the Zn uptake by roots. Under anaerobic condition Zn forms an insoluble Zn-
114 phosphate. Under this condition plant roots will not take up the soluble Zn from the Zn
115 solution as required by the plant. Under submerged conditions, when organic acid
116 concentration increased, the Zn uptake is reduced and this effects the plant growth. The Zn
117 uptake is also reduced under acidic rhizosphere condition due to the release of H^+ from the
118 roots and the surplus intake of cat ions over anions. Under acidic rhizosphere conditions, Zn
119 is released from acid-soluble fractions (e.g., absorbed Zn, organic matter or $\text{Fe}(\text{OH})_3$) and
120 is available for plant uptake. Generally, rice plants absorb most Zn from solubilization in the
121 rhizosphere because the available Zn in soil is very low in flooding condition (Dobermann
122 and Fairhurst 2000).

123 **5. Soil Factors Associated with Zn Deficiency**

124 All types of soil may be affected by Zn including: loams, sands, clays (with all classification),
125 loess, alluvium, and soils formed from basalt, sandstone, granite, volcanic ash and many
126 other rocks. In general, soils of arid and semi arid regions and the slightly acidic, leached
127 soils of warm and tropical climates are most inclined to Zn deficiency, however, crops are
128 not equally susceptible to Zn deficiency and at the same soil some crops may suffer from Zn
129 deficiency while others are not affected (Takkar and Randhawa, 1978). Lindsay (1972);
130 Pendias and Pendias (1992) and Alloway (2008) reported that major Zn deficiency causes
131 include: (i) Soils of low Zn content (Parent material), (ii) soils with Restricted Zones, (iii) pH,
132 (iv) soils low in organic matter, (v) Microbially inactivated Zn, (vi) Cool soil temperature, (vii)
133 Plant species and genotypes (viii) High level of available phosphorus and (ix) Effects of
134 nitrogen. Zn deficiency problems may occurs in soils with the subsequent characters; (a)
135 strongly alkaline in reaction (b) high phosphorus status by application of phosphatic
136 fertilizers may reduce use of zinc (c) leached sandy soils (d) acid soils of low total Zn status
137 developed on highly weathered parent material (e) calcareous soil (f) peat and muck soils (g)
138 permanently wet (water logged) and (h) high bicarbonate and magnesium in soils or irrigated
139 water (Alloway, 2004, Benton 2003).

140 **5.1 Parent Material of Soils and Zn Content**

141 The amounts of Zn in unpolluted soils typically are lower than 125 ppm (Di Baccio et al.,
142 2003; Hussain et al., 2010). The major factors affecting the concentration of Zn in soils is the
143 concentration of Zn in soil parent material. The soils derived from gneisses and granites can
144 be low in total Zn and also those originating from sandstone and limestone had lower Zn

contents (Barak and Helmke, 1993 and Pendias and Pendias, 1992). Quartz (sand) in the soils also dilutes soil Zn as concentrations of Zn in quartz are very low which range between 1 - 8 $\mu\text{g g}^{-1}$ (Brehler and Wedepohl, 1978). Also total Zn is low ($< 30 \mu\text{g g}^{-1}$) in highly leached acid sands. Zinc deficiency may occur in such soils which are inherently low in Zn. The total Zn concentrations in soils vary between 10 to 300 $\mu\text{g g}^{-1}$ with an average of 50 $\mu\text{g g}^{-1}$ (Lindsay, 1972). But the average available Zn varied from 1 to 3 $\mu\text{g g}^{-1}$ (extracted by dithizone). The problem is that only a small amount of soil Zn is available to the crop because of one or more adverse factors. The remainder of the total Zn is fixed in the soil in an insoluble or unexchangeable form and difficult to make available to crop (Stahl and James, 1991).

5.2 Soil pH

Zinc availability is highly dependent on pH. When the pH is above 6, the availability of Zn is usually very low. The availability of Zn in alkaline soils is reduced due to lower solubility of the soil Zn. The concentration of Zn in the soil solution decreases from 10^{-4} ($6.5 \mu\text{g g}^{-1}$) to 10^{-10} M ($0.007 \mu\text{g L}^{-1}$) with an increase from pH 5 to pH 8 (Kiekens, 1995). Thus it is more probable that Zn deficiency will occur in alkaline rather than acidic soils. The solubility constant values of ZnCO_3 and hydroxides suggest that a soil having high pH would usually contain a small amount of available Zn. In the case of soils characterized by high contents of hydroxyl (OH^-) ions, it is difficult to get a crop response even to applied Zn. The lower availability of Zn under alkaline conditions is attributed to the precipitation of Zn as $\text{Zn}(\text{OH})_2$ (Shukla and Mittal, 1979) or ZnCO_3 (Saeed and Fox, 1977). The higher carbonate contents in alkaline soils also absorb Zn and hold it in an unexchangeable form (Udo *et al.*, 1970). All these factors contribute to the low availability of Zn at higher pH values. Liming of acidic soils increases pH and also the Zn fixing capacity, particularly in soils with high P levels (Alloway, 2004). The movement of Zn in limed soils is considerably lower than in acidic soils (Mortvedt and Giordano, 1967) so that absorption of Zn by the crop may be low. Liming can thus reduce the Zn uptake (Shukla and Moris, 1967) and induce Zn deficiency (Viets, 1966).

5.3 Soil Organic Matter

Low organic matter contents in soils give rise to Zn deficiency as it is observed that available Zn increases with increase in organic matter in soil. Soil organic matter is an important soil constituent which originates from decomposition of animal and plant products. The most stable organic compounds in soil are humic substances such as humic and fulvic acids. Both of these substances contain a relatively large number of functional groups (OH , COOH , SH) which have a great affinity for metal ions such as Zn^{2+} . Fulvic acids mainly form chelates with Zn over a wide pH range and increases the solubility and mobility of Zn (Kiekens, 1995). Simple organic compounds such as amino acids, hydroxy acids and also phosphoric acids are effective in complexing Zn, thus increasing its mobility and solubility in soils (Pendias and Pendias, 1992). An increase in the organic matter contents of a soil will increase its Zn availability; however, if the organic matter content in soil is too high, like in peat and muck soils, this can also contribute to Zn deficiency due to the binding of Zn on solid state humic substances (Katyal and Randhawa, 1983).

5.4 Soil Texture

Lighter textured soils (sands) contain low levels of Zn. Finer texture soils like clay have higher CEC values and therefore have highly reactive sites and can retain more Zn than lighter textured soils (Shukla and Mittal, 1979). Therefore heavier textured soils with larger CEC have higher capacities for Zn adsorption than light textured soils (Stahl and James, 1991). Consequently, Zn deficiency is more likely to occur in sandy than clayey soils. Clay soils adsorb Zn and this adsorption is controlled by CEC and pH (Ellis and Knezek, 1972). Nelson *et al.*, (1953) showed that a certain portion of the Zn adsorbed on the clay was not exchangeable but was acid soluble. This portion of Zn was not available to the plants. Reddy and Perkin (1974) found that kaolinite fixes less Zn than bentonite or illite. Thus clays such as bentonite and illite with higher CECs contribute to the fixing of Zn more strongly, thus making it unavailable to plants.

5.5 Phosphate Fertilizers

Soils with higher phosphate levels, either from native P or due to application of phosphate fertilizers, can cause Zn deficiency stress in crops (Alloway, 2008). Heavy application or prolonged use of phosphatic fertilizers reduces Zn uptake by plants. This effect may be due to the physiological imbalances within the plant (Olsen, 1972). Zinc deficiency due to phosphorus application is termed “P-induced Zn deficiency” (Singh *et al.*, 1986).

5.6 Soil Flooding

Zinc deficiency is more often associated with flooded soil than dry soils. For example, rice plants under submerged conditions suffer from Zn deficiency in calcareous soils. But wheat grown in the same soil following rice grows normally (Kausar *et al.*, 1976). Zinc deficiency due to flooding was a result of Zn reaction with free sulphide (Mikkelsen and Shiou, 1977). Under the submerged conditions of rice cultivation, Zn is changed into amorphous sesquioxide precipitates or franklinite; ZnFe_2O_4 (Sajwan and Lindsay, 1988 and Singh and Abrol, 1986). Thus a delay in Zn application until after flooding for rice minimizes Zn fixation by sesquioxides (Mandal and Mandal, 1986).

5.7 Soil Temperature

In warm and moist soils, Zn uptake was higher in rice than in maize (*Zea mays* L.) (Bauer and Lindsay, 1965). Temperatures below 16°C during growth caused decreased Zn uptake in maize tops (Ellis *et al.*, 1965). It appears that Zn deficiency was associated with cool and wet seasons. Soil temperature effects appear to be due to the rate of Zn mineralization (Takkar and Walker, 1993). Other factors which can cause Zn deficiency in plants are high light intensity and long day-lengths (Marschner and Cakmak, 1989). Besides the natural soil and environmental factors, soil management practices carried out by man often causes Zn deficiency beside this plants can also suffer from Zn deficiency under adverse climatic conditions such as drought or compaction (Alloway, 2008).

6 Zinc Interaction with Other Nutrients

Interactions occur between the micronutrients and some macronutrients. ‘Interaction’ may be defined as “an influence, a mutual or reciprocal action of one element upon another in relation to plant growth” (Olsen, 1972). Another factor is the differential response of plants to one element in combination with varying levels of a second element applied simultaneously i.e. the two elements combine to produce an added effect not due to each of them acting alone (Olsen, 1972). Such interactions may take place in the soil and within the plant. These interactions should be taken into account when providing adequate micronutrient supply to plants. Other nutrients may interact with Zn by affecting its availability from soils and its status in the plant throughout the growth process, especially Zn absorption, distribution or utilization. These interactions may enhance or reduce plant growth as a response to Zn. Where an interaction does occur, it is necessary for the diagnosis and treatment of Zn deficiency to identify the factors and its sites and modes of action (Loneragan and Webb, 1993). Some important interactions of Zn with other nutrients will be discussed below.

6.1 Phosphorus-Zn Interactions

The study of the interaction between P and Zn started in 1936 (Barnette *et al.*, 1936) and till now, this important plant growth disorder is still under investigation. The interaction is usually termed ‘P-induced-Zn deficiency’. This disorder in plant growth is associated with high levels of available P or with application of P to soil. The Zn deficiency symptoms can be prevented by the application of Zn fertilizers. The actual causal relationship and mechanisms are still not fully understood. In general, four possible causes have been considered responsible for P-induced-Zn deficiency. These include (i) a P-Zn interaction in soil; (ii) a slower rate of translocation of Zn from the roots to shoot; (iii) a simple dilution effect on Zn concentration in plant tops due to growth responses to P; (iv) a metabolic disorder within plant cells related to an imbalance between P and Zn (Olsen, 1972).

It was suspected that formation of an insoluble $\text{Zn}_3(\text{PO}_4)_2$ in the soil reduced the Zn concentration in soil to deficient levels. But these suspicions were disproved by Brown *et al* (1970) who observed that $\text{Zn}_3(\text{PO}_4)_2$ was a good source of fertilizer for sorghum. The

investigation of this precipitation as a mechanism that causes Zn deficiency continued till 1970. Carrol and Loneragan (1968) reported that maximal or near maximal yields were found with legumes at 0.05 μM Zn in flowing culture and with cereals at 0.01 μM . This evidence indicates that precipitation of $\text{Zn}_3(\text{PO}_4)_2$ is not involved in P-induced -Zn deficiency. Many researchers have reported that applied P accentuated Zn deficiency symptoms in plants (Loneragan *et al.*, 1979; Sharma *et al.*, 1968). The higher P levels in soil reduced the Zn concentrations in the plant tops and also reduced total Zn contents (Singh *et al.*, 1986; Clark, 1978). These scientists suggested that P-Zn antagonism existed in the roots of the plants. Other studies suggested that although P decreased the Zn concentrations in the tops, the total Zn contents either increased or remained the same (Boawn and Brown, 1968; Boawn and Leggett, 1968). The cause of this P-induced-Zn deficiency has been suggested to be due to interference by P with the uptake, translocation, or utilization of Zn (Adriano *et al.*, 1971).

6.2 Nitrogen-Zn Interactions

Zinc deficiency can be increased or ameliorated in plants with the application of nitrogen fertilizers. The interactions resulting from the effects of N application helps to promote plant growth and, to a lesser extent, in changing the pH of the root environment since application of N promotes the growth of plants, it is possible to find positive interactions between increasing levels of Zn and N fertilizers (Alloway, 2004). Chaudhry and Loneragan (1970) reported that wheat grown on N deficient soil with adequate levels of all nutrients except N and Zn, did not respond to Zn application in the absence of NH_4NO_3 fertilizer, however, a strong response to Zn application was observed in the presence of N fertilizer.

On the other hand, in soils low in Zn and high in fertility, N fertilizers have ameliorated (or intensified) Zn deficiency by affecting Zn absorption through changing pH (Viets *et al.*, 1957). As ammonium ions have an acidifying effect, ZnSO_4 application with concurrent dressings of N were very effective in controlling Zn deficiency where ZnSO_4 alone had no effect (Viets *et al.*, 1957). It was also observed that NH_4^+ salts inhibited Zn absorption from low Zn^{2+} concentration, in a short term study with wheat (Chaudhry and Loneragan, 1972b). Ammonium ions inhibited Zn^{2+} absorption more strongly than alkali and alkaline earth anions, but were competitive with alkali and alkaline earth cat ions. So NH_4^+ effect would be diminished by relatively high concentrations of competing ions in soil. Thus any direct effect of NH_4^+ on Zn absorption would disappear.

6.3 Macronutrient Cations-Zn Interaction

Macronutrient cations such as Ca, Mg and K inhibit the absorption of Zn by plants from solution. They need to be considered when interpreting the results of solution culture experiments involving Zn nutrition, however, in soil they seem to be less effective in the inhibition of Zn absorption compared to the effects of their salts on soil pH. Zinc concentrations were highest in legumes grown in solution culture at constant pH with the lowest Ca level at which the plants were not Ca deficient. Zinc concentrations progressively decreased with increasing Ca concentrations in solutions (Bell *et al.*, 1990). This finding that Ca inhibits Zn absorption was in accord with a short term study conducted by Chaudhry and Loneragan (1972). They found that increasing concentrations of Ca (NO_3)₂ from 0 mM to 40 mM inhibited the rate of Zn absorption by wheat seedlings in a non-competitive manner, however, higher Ca concentrations (100mM) had no additional effect on Zn absorption. This inhibition was attributed to Ca as varying the anions and had little effect on Zn absorption, whereas substituting other cat ions for Ca had similarly negative effect.

In soils, the effects of Ca compounds on Zn nutrition are variable, due to the effects of its salts on soil pH. Zinc concentrations in plants growing in soil treated with CaSO_4 (which decreased the soil pH from 5.6 to 4.8) increased slightly but decreased strongly when an equivalent amount of CaCO_3 was applied (which increases the soil pH from 5.7 to 6.6) (Wear and Evan, 1968). The macronutrient cat ions K, NH_4 and Mg all inhibited the rate of Zn absorption strongly from solutions of low Ca concentrations; with increasing Ca concentrations, the inhibitory effects weakened and in the case of two ions (K, Mg) tested at

sufficiently high Ca concentration (2.5-10 mM) eventually disappeared, suggesting that they operate through the same mechanism as Ca (Chaudhry and Loneragan, 1972).

6.4 Copper-Zn Interactions

Loneragan and Webb (1993) reported that Cu and Zn may interact in several ways: Zn strongly depresses Cu absorption, Cu competitively inhibits Zn absorption and Cu nutrition affects the redistribution of Zn within plants.

A very strong Cu-Zn antagonism has been observed in wheat growing on soils deficient in Cu and Zn (Kausar *et al.*, 1976). In N-Cu-Zn experiment, Chaudhry and Loneragan, (1970) found that N fertilizer increased grain yield in the absence of Zn and diluted Cu concentrations to deficiency levels in plant. Addition to that, Zn along with N fertilizer intensified the Cu deficiency so severely that grain yield was lower than in the control plants (without NH_4NO_3). In this case, Zn intensified Cu deficiency in plants by depressing Cu uptake. This may be a result from competitive inhibition of Zn on Cu absorption (Bowen, 1987). The competitive inhibition of Cu^{2+} ion on Zn^{2+} absorption has been established in short term studies (Giordano *et al.*, 1974). While Zn severely depressed Cu uptake by wheat, Cu did not depress Zn absorption in the same experiment. The reason for the difference in soil and solution culture results may be the form of these ions present in the soil and solution. In solution studies, the Cu and Zn were present as divalent ions whereas in most of the soils they are predominantly present as complex forms and a much higher proportion of Cu is complexed compared to Zn (Geering and Hodson, 1969). So Zn^{2+} activity would be much higher than Cu^{2+} activity at the absorbing sites making it an effective competitor in Cu absorption and making its absorption less sensitive to competition from Cu (Loneragan and Webb, 1993).

6.5 Iron-Zn Interaction

The interaction between Zn and Fe is also complex like P-Zn interaction. The increased application of Zn had little effect (Norvell and Welch, 1993) or decreased (Safaya, 1976) Fe concentrations in the shoot. In the same way, higher levels of Fe generally have only a depressive effect on Zn concentration in plant tissues (Zhang *et al.*, 1991b), although it has been shown to increase have no effect on or to decrease the rate of Zn absorption by plant roots (Giordano *et al.*, 1974). These conflicting reports are probably due to differences in experimental details, especially in plant species and the concentration, ionic state and complexation of Fe.

Iron (Fe^{2+}) at low concentrations (10 μM) had no effect on the rate of Zn absorption by wheat seedlings from solutions containing 1 or 10 μM Zn and 50 mM $\text{Ca}(\text{NO}_3)_2$ (Chaudhry and Loneragan, 1972b). But at higher concentrations (100 μM Fe^{2+}), and at concentrations likely to occur in flooded rice soils, Fe completely suppressed the Zn absorption by rice seedlings from a solution of 0.05 μM ZnCl_2 with no Ca (Giordano *et al.*, 1974). Iron deficiency increased Zn concentrations in shoots of plants (Agarwala *et al.*, 1979) and also the rate of Zn absorption in both dicotyledonous plants (Romheld *et al.*, 1982) and grasses (Zhang *et al.*, 1991b). In dicotyledonous plants, the mechanism for increasing Zn absorption is probably the acidification of the rhizosphere resulting from Fe deficiency (Marschner and Cakmak, 1989). For grasses, the release of phytosiderophores under Zn deficiency is responsible for the higher Zn absorption rate as phytosiderophores have enhanced the mobilization of Zn from calcareous soils (Zhang *et al.*, 1991b and Treeby *et al.*, 1989). In a similar way, under Zn deficient conditions, Fe accumulated in the shoots of Zn deficient navy beans and corn plants are possibly due to the acidification of the rhizosphere and the release of reductants and phytosiderophores ((Ambler and Brown, 1969 and Jackson *et al.*, 1962).

7. Dilution Effect

When the rate of plant growth is faster than the rate of uptake of a particular nutrient, the concentration of the nutrient is "diluted" in the plant (Singh *et al.*, 1986; Olsen, 1972). These researchers also showed that in wheat and bean the yield and total Zn content increased with P application while the Zn concentrations in plants decreased. A response in yield was

found for applied P, so a dilution effect on Zn is largely accountable for this effect. In general, this interaction occurs when the soil is deficient in P and/or slightly deficient in available Zn. The growth rate increases due to applied P but the rate of Zn uptake does not increase fast enough to maintain Zn concentration in plants. In some cases of P-induced-Zn deficiency the dilution effect only partially explains the data (Singh *et al.*, 1986). The applied P reduces the Zn concentration in the tops of the plant while the yield response to P is minimal (Singh *et al.*, 1986).

8. Distribution of Zn in Plant Roots and Tops

Zn is an essential micronutrient involved in a wide variety of physiological processes (Cakmak, 2000; Reeves and Baker, 2000; Doncheva *et al.*, 2001; Stoyanova and Doncheva, 2002; Di Baccio *et al.*, 2005; Broadley *et al.*, 2007). Zn uptake varies among plant species and is determined by the composition and concentration of the growth media. Zn uptake occur as divalent cation or as complexes with organic ligands and display a linear pattern with its concentration in the nutrient solution of soils (Kabata-Pendias and Pendias, 2001), roots load it via xylem to the shoot tissues (Broadley *et al.*, 2007). The Zn translocation to roots xylem occurs via symplast and apoplast but its high levels have also been detected in the phloem, denoting that this metal is translocated through both xylem and phloem tissues (Brennan, 2005; Broadley *et al.*, 2007; Haslett *et al.*, 2001).

9. Zinc Efficiency

Zn efficiency can be defined as “the ability of plants to maintain high yields in soils with low Zn availability”. Many mechanisms are perhaps involved in Zn efficiency (Rengel, 2001). Depending on the nature of experiments and plant species, the most significant mechanisms may be Zn utilization in tissues (Hacisalihoglu and Kochian, 2003) and Zn uptake (Genc *et al.*, 2006). Under Zn deficiency, Zn-efficient genotypes have a high activity of Cu/Zn anhydrase (Hacisalihoglu *et al.*, 2003a; Yu *et al.*, 1999; Cakmak *et al.*, 1997) and carbonic anhydrase (Hacisalihoglu *et al.*, 2004; Rengel, 1995). Zn efficiency and Zn uptake are very susceptible for plant growth and its total content in soil is influenced by several soil properties like pH, CaCO₃, organic matter content, crop, as well as cultivars and nutrient interactions in soil environment. There is no precise mechanism used in determining Zn efficiency is available so far; however, several crops have been evaluated for their Zn efficiency like: beans (Ambler and Brown, 1969; Hacisalihoglu *et al.*, 2001; Hacisalihoglu *et al.*, 2003b), wheat (Graham and Raangel 1993; Cakmak *et al.*, 1997a) and rice (Brown *et al.*, 1993; Clark, 1990; De Datta and Neue, 1993; Nand, 2002; Geo, 2007).

9.1 Effect of Zn Deficiency on Plants and Its Correction

Zinc is an essential micronutrient for plant growth and plays an important role in the catalytic part of several enzymes (Fageria, 2002) its deficiency will result in stunted growth. Many researchers observed that Zn is closely related to the nitrogen metabolism pathway of plants, thus causing a reduction in protein synthesis for Zn deficient plants. Zinc deficiency significantly affects the root system including root development (Fageria, 2004). Zinc deficiency affects the absorption of water and nutrients from soil and thus resulting in growth and yield reduction in the plant. Epstein and Bloom (2005) indicated that the flowering and fruiting process were greatly reduced under severe Zn deficiency.

Zinc deficiency is one of the major constraints in world food production. Identification of Zn-deficient areas, and causes would help in planning the appropriate strategies to correct these Zn deficiencies. Although Zn is widely used as a fertilizer, but efficient and economical methods to correct its deficiency on a long term basis and in a specific cropping system is desirable. Zinc deficiency can be corrected through the application of Zn fertilizers, recycling crop residues, natural organic manures and cultivation of Zn efficient genotypes (Singh, 2008). Zinc fertilizers are broadcast and sprayed onto topsoil, banded in the seedbed, applied as foliar sprays, used as seed treatment and in the case of transplanted rice seedlings, roots of these seedlings are dipped into Zn before transplanting. Zinc sulphate is the commonly used fertilizer compound (ZnSO₄·7H₂O containing 26% Zn, or ZnSO₄·H₂O

409 containing 37% Zn). Other Zn compounds are Zn chloride (ZnCl_2), Zn nitrate ($\text{Zn}(\text{NO}_3)_2$), Zn
410 oxide (ZnO), Zn oxy-sulphate and Zn-coated urea (Mortvedt and Gilkes 1993).

411 **9.2 Zinc Deficiency Symptoms in Plants**

412 Visual symptoms of Zn deficiency in plants are fairly characteristic and are relatively easy to
413 identify. These distinctive symptoms are useful for recognizing acute Zn deficiency and for
414 indicating Zn responsive soils, but not the hidden or marginal deficiencies. The most
415 common symptoms of Zn deficiency include: stunted growth, shortened internodes and
416 petioles, and small malformed leaves (little leaf) which results in the "rosette" symptom in the
417 early growth stages of dicotyledons and "fan shaped" stems in monocotyledons (Snowball
418 and Robson, 1986). The deficiency symptoms first appear on young leaves as Zn is
419 immobile under conditions of deficiency. These leaves remain small, cup upward and
420 develop interveinal chlorosis and necrotic spots on the upper leaf surfaces which later join to
421 each other to form brown necrotic and brittle patches. The necrosis is often more noticeable
422 on middle aged leaves which eventually wilt, bend and collapse (Brennan *et al.*, 1993). Zinc
423 deficiency is typically patchy, even within a single field and symptoms develop rapidly but
424 depend greatly on the degree of stress (Kubota and Allaway, 1972). Enzyme activity, like
425 ribonuclease activity or carbonic anhydrase activity can be used as an index for precise
426 information. This is particularly important in the initial stages of growth when micronutrient
427 requirements of plants are very low and the total contents of Zn fail to provide precise
428 information about the hidden deficiency. Zinc deficient plants are unthrifty, lack vigor; give
429 patchy appearance with short and thin stems. In young plants interveinal areas are with dark
430 brown necrotic lesions (Benton 2003).

431 The visible symptoms in rice are: wilting due to loss of turgidity in the leaves, basal chlorosis
432 of the leaves, delayed development of the plants, "bronzing" of the leaves and in some
433 cases death of the rice seedling (Neue *et al.*, 1998). Neue *et al.* (1998) stated that the
434 common symptoms of Zn deficiency in rice are: chlorosis in the mid rib at the base of the
435 youngest leaf within 2-4 weeks after sowing or transplanting and the appearance of brown
436 spots on the older leaves. The spots enlarge, coalesce and give the leaves a brown color.
437 Zinc deficient plants show stunted growth and reduced tillering. If the deficiency is not too
438 severe the plant may recover after 4-6 weeks but maturity is delayed and yields of
439 susceptible cultivars are reduced. The most noticeable symptom is the plant's loss of
440 turgidity, where plants fall over and float on the surface of the water. The basal leaves
441 become pale green and after 3-7 days the leaves become chlorotic. It is important to note
442 that visual symptoms of Zn deficiency in rice vary, to a certain extent, with soil type, cultivar
443 and growth stages. Symptoms can be mistaken for those of N, Mg, Mn or Fe deficiencies
444 which are often combined with Zn deficiency, making it difficult to distinguish between the
445 symptoms of the two. Therefore, plant analysis is required for confirmation (Dobermann and
446 Fairhurst, 2000; Alloway, 2004, 2008).

447 **10. Zinc Critical Levels in Plants**

448 Zinc deficiency is one of the major constraints in world food production. It is therefore
449 essential to identify the Zn-deficient areas, and the different causes of deficiency. It would
450 help in planning the appropriate strategies to correct these Zn deficiencies. Although Zn is
451 being used as a fertilizer, an understanding of efficient and economical methods to correct
452 Zn deficiency on a long term basis and in a specific cropping system is desirable. It was
453 reported by Katyal and Randhawa (1983) that there is a strong relationship between Zn
454 concentration in tissues with the growth and yield of crops. The critical limits of Zn in plants
455 indicates deficiency as suggested by Dobermann and Fairhurst (2000) are $< 10 \text{ mg kg}^{-1}$
456 definite Zn deficiency, $10\text{--}15 \text{ mg kg}^{-1}$ very likely, $15\text{--}20 \text{ mg kg}^{-1}$ likely and $>20 \text{ mg kg}^{-1}$
457 unlikely (sufficient). In most crop species leaf sufficiency range for Zn 15 to 50 ppm in the dry
458 matter of mature plants and in most cases 15 ppm Zn is considered as critical value (Benton
459 2003).

460 **11. Zinc Toxicity**

The threshold of Zn toxicity varies among plant species, time of exposure to Zn stress and composition of the nutrient growth medium. Plant growth inhibition extends in *E. maculata* and *E. urophylla* by five weeks after addition of 400-1600 mM ZnSO₄, whereas *Pisum sativum* became inhibited after 1000 µM Zn application (Soares et al., 2001, Doncheva et al., 2001). Photosynthesis is strongly affected in plants exposed to heavy metals excess (Prasad, 1999). High Zn concentrations in plants can cause phytotoxicity. The yield may be reduced when plant leaf Zn concentrations reaches about 300 - 1000 µg Zn g⁻¹. A typical phytotoxicity critical concentration is about 500 µg Zn g⁻¹ (Chaney, 1993). The best way to identify Zn deficiency in crops is the determination of Zn concentrations in tissues, however, the results should be interpreted in full recognition of the interaction of Zn with other nutrients because the deficiency of one nutrient may result in excess accumulation of other nutrients by a plant (Katyal and Rawdhawa, 1983).

12. Effect of Zn on Microbial Activity

Microorganism requires various nutrients for their growth and metabolism. Among the nutrients, Zn is an element present in the enzyme system as co-factor and mental activator of many enzymes (Vankatakrishnan et al., 2003). According to Vankatakrishnan et al. (2003); Baath (1992) and Doelman and Haanstra (1984), the solubilization of Zn might limit the growth of bacteria at higher levels (>13.60 mg kg⁻¹). Furthermore, cell growth as well as microbial populations and their activity in soil were badly affected under high levels of Zn.

13. Conclusion

Our extensive review of literature has shown that Zn is very essential plant nutrient for all types of crops. It is deficient in all parts of the globe with different types of soils. Under these conditions application of Zn fertilizer is necessary for healthy crop growth and higher yields. Soil and foliar applications of Zn fertilizer are recommended for correcting deficiencies. Soil dressings of Zn chelates, sulfates and oxides should be broadcast and mixed in the soil. Soil applied Zn had residual effects for subsequent crops but foliar sprays have no residual effect and fresh applications must be made to each crop.

REFERENCES:

- Adriano, D.C., Paulson, G.M. and Murphy, L. S. (1971). P-Fe and P-Zn relationship in corn seedlings as affected by mineral nutrition. *Agronomy Journal*, 63,36-39.
- Alloway, B.J. (2004). In *Zinc in Soil and Crop Nutrition*. International Zinc Association. Brussels, Belgium.
- Alloway, B.J. (2008). Micronutrients and crop production. In *Micronutrient Deficiencies in Global Crop Production*. © Springer Science Business Media B.V. pp. 1-39.
- Ambler, J.E. and Brown J.C. (1969). Cause of differential susceptibility to Zn deficiency in two varieties of navy beans. *Agronomy journal*, 61, 41-43.
- Baath, E. (1992). Measurement of heavy metal tolerance of soil bacteria using thymidine incorporation into bacteria extracted after homogenization-centrifugation. *Soil Biological biochemistry*, 24, 1167-1172.
- Barak, P. and Helmke P.A. (1993). The chemistry of Zinc. In *Zinc in Soils and Plants*. (ed.) A.D. Robin. Dordrecht: Kluwer Academic Publishers.
- Barnette, R.M., Camp. J.P., Warner J.D. and Gall, J.D. (1936). Use of zinc sulphate under corn and other field crops. *Fla. Agri. Exp. Sta. Bull.*, 293: 3.
- Bauer, A. and Lindsay W.L. (1965). The effect of soil temperature on the availability of indigenous soil zinc. *Soil Science Society of America. Proc.*, 29, 413-420.
- Bell, R.W., Kirk, G., Plaskell, D. and Loneragan J.F. (1990). Diagnosis of zinc deficiency in peanut by plant analysis. *Communication Soil Sci. Plant Analysis*, 21, 273-285.
- Benton, J.J. (2003) *Agronomic handbook; management of crops, soils and their fertility*. CRC press LLC. USA.
- Boawn, L.C. and Brown, J.C. (1968). Further evidence for a P/Zn imbalance in plants. *Soil Science Society of America. Proc.*, 32, 94-97.
- Boawn, L.C. and Leggett G.E. (1968). Phosphorus and zinc concentrations in Russett

- 514 Burbank potato tissue in relation to development of zinc deficiency symptoms. *Soil*
515 *Science Society of America. Proc.*, 28, 229-232.
- 516 Bowen, J.E. (1987). Physiology of genotypic differences in Zn and Cu uptake in rice and
517 tomato. Proceedings of 2nd International Symposium on Genetic Aspects of *Plant*
518 *Mineral Nutrition*.
- 519 Brennan, R. F. (2005). Zinc Application and Its Availability to Plants. Ph. D. dissertation,
520 School of Environmental Science, Division of Science and Engineering, Murdoch
521 University
- 522 Brehler, B., and Wedepohl, K.H. (1978). Zinc. In K.H. Wedepohl (ed). *Handbook of*
523 *Geochemistry* (p.125). Vol. II/3. Springer-Verlag, Berlin.
- 524 Brennan, R.F., Armour, J.D., and Reuter, J.D. (1993). Diagnosis of Zinc Deficiency. In
525 Robson, A. D. Kluwer (ed). *Zinc in soils and plants* (pp. 167-181). Dordecht:
526 Academic Publisher.
- 527 Brown, A.L., Krantz, B.A. and Edding, J.L. (1970). Zinc-phosphorus interaction as
528 measured by plant response and soil analysis. *Soil Science.*, 110, 415-420.
- 529 Brown, P.K., Cakmak, I., and Zhang, Q.L. (1993). Form and function of Zn plant. In
530 Robson A D. Kluwer (ed). *Zinc in soil and plants* (pp. 93-106). Dordrecht: Academic
531 Publishers.
- 532 Broadley, M. R., P. J. White, J. P. Hammond, I. Zelko and A. Lux. (2007). Zinc in plants.
533 *New Phytol.* 173:677–702.
- 534 Cakmak, I. (2002). Plant nutrition research priorities to meet human needs for food in
535 sustainable ways. *Plant Science*, 247, 3-24.
- 536 Cakmak, I., Ekiz, H., Yilmaz, A., Torun, B., Koleli, N., Gultekin, I., Alkan, A., and Eker,
537 S. (1997). Differential response of rye, triticale, bread and durum wheats to zinc
538 deficiency in calcareous soils. *Plant and Soil*, 188, 1–10.
- 539 Cakmak, I. (2000). Role of zinc in protecting plant cells from reactive oxygen species. *New*
540 *Phytol*, 146, 185–205.
- 541 Cakmak, I. (2002). Plant nutrition research priorities to meet human needs for food in
542 sustainable ways. *Plant Science*, 247, 3-24.
- 543 Carrol, M.D. and Loneraga, J.F. (1968). The relevance of solution cultural studies to the
544 absorption of Zn from soils. In transcation of 9th international congress of soil
545 science. vol. 2. pp. 191-202. *International society of soil science and Angus and*
546 *Robertson*, Sydney
- 547 15th Federal Convention.vol.1. Australian water and wastewater Association,
548 Queensland, Australia.
- 549 Chaudhry, F.M. and Loneragan, J.F. (1970). Effect of nitrogen, copper and zinc
550 fertilizers on the copper and zinc nutrition of wheat plants. *Australian Journal of*
551 *Agriculture Res.*, 21, 865-879.
- 552 Chaudhry, F.M., and Loneragan J.F. (1972). Zinc absorption by wheat seedlings. I.
553 Inhibition by hydrogen ions and micronutrient cat ions. *Soil Science Society of*
554 *America. Proc.*, 36, 327-331.
- 555 Clark, R.B. (1978). Differential response of maize inbreeds to Zn. *Agronomy Journal*,
556 70, 1057-1060.
- 557 Clark, R.B. (1990). Physiology of cereals for mineral nutrient uptake use and efficiency.
558 In V.C. Baligar and R.R. Duncan (Ed). *Crops as enhancers of nutrient use* (pp. 131-
559 209). San Diego: Academic press
- 560 Dang, H., R. Li, Y. Sun, X. Zhang and Y. Li. (2010). Absorption, accumulation and
561 distribution of zinc in highly-yielding winter wheat. *Agr. Sci. China* 9(7):965-973.
- 562 Disante, K. B., D. Fuentes and J. Cortina. (2010). Response to drought of Zn-stressed
563 *Quercus suber* L. Seedlings. *Env. Exp. Bot.* 70:96-103.
- 564 De Datta, S.K and Neue, H.U. (1993). Success in rice improvement for poor soils. In:
565 *workshop on adaptation of plants to soil stress*, Lincoln. proceedings. Lincoln:
566 University of Nebraska, pp. 248-268.

567 Di Baccio, D., R. Tognetti, L. Sebastiani and C. Vitagliano. (2003). Responses of *Populus*
568 *deltoides* x *Populus nigra* (*Populus* x *euramericana*) clone I-214 to high zinc
569 concentrations. *New Phytol.* 159:443-452.

570 DiBaccio, D., S. Kopriva, L. Sebastiani and H. Rennenberg. (2005). Does glutathione metabolism
571 have a role in the defence of poplar against zinc excess? *New Phytol.* 167:73-80.

572 Doncheva, S., Z. Stoyanova and V. Velikova. (2001). Influence of succinate on zinc toxicity
573 of pea plants. *J. Plant Nutr.* 24(6):789-804.

574 Dobermann, A. and Fairhurst, T. (2000). Rice: Nutritional Disorders and Nutrient
575 Management. Potash and Phosphate Institute and Potash and Phosphate Institute
576 of 12. References 145 Canada (PPI/PPIC) and International Rice Research Institute
577 (IRRI), Singapore and Makati City, the Philippines.

578 Doelman, P. and Haanstra L. (1984). Short-term and long-term effects of cadmium,
579 chromium, copper, nickel, lead and zinc on soil microbial respiration in relation to
580 abiotic soil factors. *Plant Soil*, 79:317-327 Dordrecht, the Netherlands: Kluwer
581 Academic Publisher, 413-423.

582 Ellis, B.G. and Knezek B.D. (1972). Adsorption Reactions of Micronutrients in Soils. In
583 Mortvedt, J., Giordano J. and Lindsay W.L. (ed), *Micronutrient in Agriculture* (pp. 59-
584 78). Soil Science Society of America. Madison, Wis.

585 Ellis, B.G., Davis, J.F. and Judy W.H. (1965). Effect of method of incorporation of Zn in
586 fertilizer on zinc uptake and yield of pea beans (*Phaseolus vulgaris*). *Soil Science*
587 *Society of America*. Proc. 29, 635-636.

588 Epstein and Bloom. (2005). *Mineral Nutrition of Plants: Principles and Perspectives*.
589 Sinauer Assoc.

590 Fageria, N.K. (2002). Influence of micronutrients on dry matter yield and interaction with
591 other nutrients in annual crops. *Pesq. Agropec. Bras.*, 37, 1765-1772.

592 FAO/WHO (2002). *Human vitamin and mineral requirements* - Report of a joint
593 FAO/WHO expert consultation - Bangkok, Thailand, FAO, Rome. Chapter 16. Zinc.
594 pp 257-270.

595 Gao, X.P. (2007). Bioavailability of Zinc to Aerobic Rice. PhD thesis, Wageningen
596 University, Wageningen, The Netherlands.

597 Geering, H.R. and Hodson, J.F. (1969). Micronutrient cation complexes in soil solution.
598 *Soil Science Society of America*. Proc. 33, 54-59.

599 Genc, Y., McDonald, G.K. and Graham, R.D. (2006). Contribution of different
600 mechanisms to zinc efficiency in bread wheat during early vegetative stage. *Plant*
601 *Soil*, 281, 353-367.

602 Giordano, M., Noggle J.C. and Mortvedt J.J. (1974). Zinc uptake by rice, as affected by
603 metabolic inhibitors and competing cations. *Plant Soil*, 41, 637-646.

604 Graham, R.D. and Rangel, Z. (1993). Genotypic Variation in Zn uptake and Utilization
605 by Plants. In A.D. Robson (ed). *Zn in soil and plants* (pp. 107-114), Dordrecht, the
606 Netherlands.

607 Graham, R.D., Welch, R.M and Bouis, H.E. (2001). Addressing micronutrients
608 malnutrition through enhancing the nutritional quality of staple foods principles,
609 perspectives and knowledge gaps. *Advanced Agronomy*. 70, 77-142.

610 Hacisalihoglu, G. and Kochian, L.V. (2003). How do some plants tolerate low levels of
611 soil zinc? Mechanisms of zinc efficiency in crop plants. *New Phytologist*, 159, 341-
612 350.

613 Hacisalihoglu, G., Hart J.J. and Kochian L.V. (2001). High and low-effinity zinc transport
614 systems and their possible role in zinc efficiency in bread wheat. *Plant Physiology*,
615 125, 456-463.

616 Hacisalihoglu, G., Hart, J.J., Vallejos, C.E. and Kochian, L.V. (2004). The role of
617 shoot localized processes in the mechanism of Zn efficiency in common bean.
618 *Planta*, 218, 704-711.

619 Hacisalihoglu, G., Hart, J.J., Wang, Y., Cakmak, I., and Kochian L.B. (2003). Zinc

620 efficiency is correlated with enhanced expression and activity of Cu/ Zn superoxide
 621 dismutase and carbonic anhydrase in wheat. *Plant Physiology*, 131, 595-602.
 622 Haslett, B. S., R. J. Reid and Z. Rengel. (2001). Zinc mobility in wheat: uptake and
 623 distribution of zinc applied to leaves or roots. *Ann. Bot.* 87:379–386.
 624 Helmke P. A. Koons R.D., Schomberg P. J. and Iskandar I. K. 1977. Determination of
 625 trace element contamination of sediments by multielement analysis of the clay-size
 626 fraction. *Environmental. Science Technology*. 11, 984-989.
 627 Hotz, C and K. H. Brown (eds.) (2004) Assessment of the Risk of Zinc Deficiency in
 628 Populations and Options for its Control. *Food and Nutrition Bulletin*, 25 (Supplement
 629 2): S91-S204.
 630 Hussain S., M. A. Maqsood and Rahmatullah. (2010). Increasing grain zinc and yield of
 631 wheat or the developing world: A Review. *Emir. J. Food Agric.* 22(5):326-339.
 632 Imtiaz, M. (1999). *Zn deficiency in cereals*. PhD Thesis Reading University, U.K.
 633 IRRI (2000). Nutritional Disorders and Nutrient Management in Rice. *Inter. Rice Res.*
 634 *Ins. Manila, Philippines* .
 635 Jackson, M.L. (1962). *Soil Chemical Analysis*. London: Constable and Company Ltd.
 636 Katyal, J.C. and. Randhawa, N.S. (1983). Micronutrients FAO Fertilizer and Plant
 637 Nutrition Bulletin 7. Rome: Food and Agriculture Organization of the United Nations.
 638 Kausar, M.A., Chaudry, F.M., Rashid, A., Latif, A. and Alam, S.M. (1976). Micronutrient
 639 availability to cereals from calcareous soils. I. Comparative Zn and Cu deficiency
 640 and their mutual interaction in rice and wheat. *Plant and Soil*, 45, 397-410.
 641 Kabata-Pendias, A. and H. Pendias. (2001). Trace elements in soils and plants, CRC Press,
 642 Boca Raton - London - New York – Washington D.C.
 643 Kasim, W. A. (2007). Physiological consequences of structural and ultra-structural changes
 644 induced by Zn stress in *Phaseolus vulgaris*. I. Growth and Photosynthetic apparatus.
 645 *Int. J. Bot.* 3(1):15-22.
 646 Kapoor S, Kobayashi A, Takatsuji H (2002) Silencing of the Tapetum-Specific Zinc Finger
 647 Gene TAZ1 Causes Premature Degeneration of Tapetum and Pollen Abortion in
 648 Petunia. *Plant Cell Online* , 14(10):2353-2367.
 649 Kiekens, L. (1995) Zinc in Heavy Metals. In B.J. Alloway (Ed.). *Soils*. London: Blackie
 650 Academic and Professional
 651 Kobayashi A, Sakamoto A, Kubo K, Rybka Z, Kanno Y, Takatsuji H (1998). Seven zinc-
 652 finger transcription factors are expressed sequentially during the development of
 653 anthers in petunia. *Plant J* 13:571-576
 654 Krauskopf K. B. 1972. Geochemistry of Micronutrients. *In* Micronutrients in Agriculture.
 655 (Eds.) Mortved J. J., Goirdano P. M. and Lindsay W. L. Soil Science Society
 656 America., Inc. Madison, Wisconsin USA.
 657 Kubota, J. and Alloway, W.H. (1972). *In* Micronutrients in Geographic Distribution of
 658 Trace Metal Problems.
 659 Lindsay, W.L. (1972). Zinc in soil and plant nutrition. *Advance Agronomy*, 24, 147-188.
 660 Loneragan, J.F. and Webb M.J (1993). Interactions between Zn and other Nutrients
 661 affecting the Growth of Plants. In A.D. Robson (ed). Zinc in soils and plants (p.151).
 662 Kluwer Academic Publisher, Dordrecht.
 663 Loneragan, J.F., Grove T.S., Robson A.D., and Snowball, K. (1979). Phosphorus
 664 toxicity as a factor in zinc phosphorus interaction. *Soil Science Society of America.*
 665 *J.*, 43, 966-972.
 666 Lukaski, H.C. (2004) Vitamin and mineral status: effects on physical performance.
 667 *Nutrition*, 20: 632–644.
 668 Maharana, D.P., Sarengi, S.K. Singh, N.R.B., Ali M.H. (1993). Proceeding of the
 669 workshop on micronutrients. 22-23 January 1992. Bhubaneswar, India. pp. 228-238.
 670 Mandal, L.N and Mandal, B. (1986). Zinc fraction in soils in relation to Zn nutrition of low
 671 land rice. *Soil Science*, 142, 141-148.
 672 Marschner, H. and Cakmak, I. (1989). High light intensity enhances chlorosis and

- necrosis in the leaves of zinc, potassium and manganese deficient bean (*Phaseolus vulgaris* L.) plants. *Plant Physiology*, 134, 308-315.
- Marschner, H. (1995). Mineral nutrition of higher plants (2nd ed.). London: Academic Press.
- Tisdale, S.L., Nelson, W.L. and Beaten, J. D. 1984. Zinc In soil Fertility and Fertilizers. Fourth edition, pp. 382-391. Macmillan Publishing Company, New York.
- Mikkelsen, D.S. and Shiou, K. (1977). Zinc fertilization and behaviour in flooded soils. Spec. Publ. No. 5 Comm. Agric. Bur., Farnham Royal. p. 59. Mineral Stresses. In A.R. Yeo and T.J. Flowers (ed). *Approaches to Crop Improvement* (pp. 175- 200) . Berlin: Springer-Verlag.
- Morley, J.E. (2004) The top 10 hot topics in aging. *J Gerontol.*, 59: 24–33.
- Mortvedt, J. J. and Giordano P. M. (1967). Zinc movement in soils from fertilizer granules. *Soil Science Society of America. Proc.*, 31, 606.
- Mortvedt, J. J., and Gilkes, R. J. (1993). Zinc fertilisers. In A. D. Robson (Ed.), *Zinc in soils and plants* (pp. 33–44). Dordrecht: Kluwer Academic Publishers.
- Nand, F. (2002). Screening method of low land rice genotypes for Zn uptake efficiency. *Scientia Agricola*, 58, 623-626.
- Nand, R., and Ram, N. (1996). Amelioration of Zinc stress by farmyard manure in a rice-wheat-cowpea system. *Acta-Agronomica-Hungarica*, 44(1), 35-39.
- Nelson W. L., Mehlich, A. and Winters, E. (1953). The Development, Evaluation and use of Soil Tests for Phosphorus Availability. In Pierr W.H. and A.G. Norman (Eds.) *Soil and Fertilizer Phosphorus in Crop Nutrition* (pp. 153-158). New York; Agrono. Monogr. Acad. Press.
- Neue, H.U, Quijano, C., Senadhira, D., Setter, T. (1998). Strategies for dealing with micronutrient disorders and salinity in lowland rice systems. *Field Crops Research*, 1998, 56, 139-155.
- Norvell, W.A. and Welch, R.M. (1993). Growth and nutrient uptake by barley :Studies using an N(2-Hydroxyethyle) ethylenedinitrilotriacetic acid buffered nutrient solution technique. I. Zinc ion requirements. *Plant Physiology*, 101, 619-625.
- Pendias A.K. and Pendias, H. (1992). *Trace Elements in Soil and Plants* (2nd edition). Boca Raton, Florida: CRC Press.
- Peck, A.W. and G. K. McDonald. (2010). Adequate zinc nutrition alleviates the adverse effects of heat stress in bread wheat. *Plant Soil* 337:355- 374.
- Prasad, M. N. V. (1999). Trace Metals. In: M. N. V. Prasad and J. Hagemeyer (Eds). pp 207-249. Heavy metal stress in plants: from molecules to ecosystems. Springer. Berlin - New York.
- Rajan, A.R. (1993). Relative utilisation of different zinc carries in rice (*Oryza sativa* L.). *Indian Journal of Agriculture Chemistry*, 26(1), 1-4.
- Reddy, M.R. and H.F. Perkin. (1974). Fixation of Zn by clay minerals. *Soil Science of America. Proc.*, 38, 229-230.
- Rengel, Z. (1995) Carbonic anhydrase activity in leaves of wheat genotypes differing in Zn efficiency. *Journal of Plant Physiology*, 147, 251-256.
- Reeves, R. D. and J. M. Baker. (2000). Metalaccumulating plants. In: H. Raskin and B. D. Ensley (Eds.) pp 193–230. *Phytoremediation of Toxic Metals: Using Plants to Clean Up the Environment*. John Wiley & Sons Inc., London.
- Rengel, Z. (2001). genotypic differences in micronutrient use efficiency in crops. *Comm. Soil Science and Plant. Analysis.*, 32, 1163-1186.
- Romheld, V., Marschner H. and Kramer, D. (1982). Response to Fe deficiency in roots of “Fe-efficient” plant. *Journal of Plant Nutrition*, 5, 489-498.
- Saeed, M. and Fox, R.L. (1977). Relation between suspension pH and Zn solubility in acid and calcareous soils. *Soil Science*, 124, 199-204.
- Safaya, N. M. (1976). Phosphorus-Zinc interaction in relation rate of phosphorus, zinc, copper, manganese and iron in corn(*Zea mays* L.). *Journal of Soil Science Society America*, 71, 132-136.

- 726 Sajwan, K.S. and Lindsay, W.L. (1988). Effect of redox, zinc fertilisation and incubation
727 time on DTPA-extractable zinc, iron and manganese. *Commun. Soil Science and*
728 *Plant Analysis*, 19, 1-11.
- 729 Sharma PN, Chatterjee C, Sharma CP, Agarwala SC (1987). Zinc deficiency and anther
730 development in maize. *Plant Cell Physiol* 28(1):11-18.
- 731 Sharma, K. C., Karantz, B.A, Brown, A.L. and Quick, J. (1968). Interaction of Zn and P
732 in the tops and roots of corn and tomatoes. *Agronomy Journal*, 60, 453-456.
- 733 Soares, C. R. F. S., P. H. Graziotti, J. O. Siqueira, J. G. De Carvalho and F. M. S. Moreira.
734 (2001). Toxidez de zinco no crescimento e nutrição de *Eucalyptus maculata* e
735 *Eucalyptus urophylla* em solução nutritiva. *Pesq. Agropec. Bras.* 36(2):339-348.
- 736 Shukla, U.C. and Mittal, S.B. (1979). Characterization of zinc application in some soils
737 of India. *Journal of Soil Science Society of America*, 43, 905-908.
- 738 Shukla, U.C. and Moris H.D. (1967). Relative efficiency of several zinc sources for corn.
739 *Agronomy Journal*, 59, 200.
- 740 Sillanpaa, M. (1990). Micronutrients Assessment at the Country Level. An international
741 Study FAO Soils Bulletin 63. Food and Agriculture Organization of the United
742 Nations
- 743 Singh, J.P., Karamonas, R.E. and Stewart, J.W.B. (1986). Phosphorus-induced zinc
744 deficiency in wheat on residual phosphorus plots. *Agronomy Journal*, 78, 668-675.
- 745 Singh, M.V. (2008). Micronutrients Deficiencies in Crops and Soils in India. In B.J.
746 Alloway (ed). *Micronutrient Deficiencies in Global Crop Production* (p.93-125).
747 Springer Science+Business Media B.V.
- 748 Singh, M.V. and Abrol, I.P. (1986). Transformation and movement of zinc in an alkali
749 soil and their influence on the yield and uptake of zinc by rice and wheat crops. *Plant*
750 *Soil*, 94, 445-449.
- 751 Snowball, K. and Robson, A.D. (1986). Symptoms of Nutrient Deficiencies: Lupins.
752 University of Western Australia Press, Nedlands Australia.
- 753 Stahl, R.S. and James B.R. (1991). Zinc sorption by B Horizons Soils as a function of
754 pH. *Journal of Soil Science Society of America*, 55, 1592-1597.
- 755 Stoyanova, Z. and S. Doncheva. (2002). The effect of zinc supply and succinate treatment
756 on plant growth and mineral uptake in pea plant. *Bras. J. Plant Physiol.* 14(2):111-
757 116
- 758 Takkar, P.N. and Randhawa, N.S. (1978). Micronutrients in Indian Agriculture. *Fertility*
759 *News*, 23, 3-26.
- 760 Takkar, P.N. and Walker, C. (1993). The Distribution and Correction of Zinc Deficiency.
761 In A.D. Robson (ed). *Zinc in Soils and Plants* (pp. 51). London: Kluwer Academic
762 publisher.
- 763 Tavallali, V., M. Rahemi, S. Eshghi, B. Kholdebarin and A. Ramezani. (2010). Zinc
764 alleviates salt stress and increases antioxidant enzyme activity in the leaves of
765 pistachio (*Pistacia vera* L. 'Badami') seedlings, *Turk. J. Agr. Forest* 34(4):349-359.
- 766 Treeby, M., Marschner, H. and Romheld, V. (1989). Mobilisation of iron and other
767 micronutrients from a calcareous soil by plant born microbial and synthetic metal
768 chelator. *Plant and Soil*, 114, 217-226.
- 769 Udo, E.J., Bhon, L.H. and Tukker, T.C. (1970). Zinc adsorption by calcareous *Journal of*
770 *Soil Science Society of America. Proc*, 34, 405-407.
- 771 Van Breemen, N., and Castro, R.U. (1980). Zinc deficiency in wetland rice along a
772 toposequence of hydromorphic soils in the Philippines. II. Cropping experiment.
773 *Plant and Soil*, 57, 215-221.
- 774 Vankatakrishnan, S.S, Sudlayandy, R.S, Savariappan, A.R. (2003). Assessing in vitro
775 solubilization potential of different zinc solubilizing Bacteria (ZSB) isolates. *Brazilian*
776 *J. Microbiol.*34:121-125.
- 777 Viets, F.G. (1966). Zinc Deficiency in Soil Plant System. In A.S. Prasad, C. Charles,
778 Thomas Springfield II (ed). *Zinc Metabolism*.

779 Viets, F.G., Boawn, L.C. and Crawford C.L. (1957). The effect of nitrogen and types of
780 nitrogen carrier on plant uptake indigenous and applied zinc. *Journal of Soil Science*
781 *Society of America*, 21, 197-201
782 Wear, J.I. and Evan, C.E. (1968). Relationship of zinc uptake by corn and sorghum to
783 soil zinc measured by three extractants. *Journal of Soil Science Society of America*.
784 *Proc.*, 32, 543-546.
785 Welch, R.M. (2002). The impact of mineral nutrients in food crops on global human
786 health. *Plant and Soil*, 247, 83-90.
787 Yu, Q., Worth, C. and Rengel, Z. (1999). Using capillary electrophoresis to measure
788 Cu/Zn superoxide dismutase concentration in leaves of wheat genotypes differing in
789 tolerance to zinc deficiency. *Plant Science*, 143, 231-239.
790 Zhang, F., Romheld, V. and Marschner, H. (1991). Diurnal rhythm release of
791 phytosiderophore and uptake rate of zinc in Fe-efficient wheat. *Soil Science and*
792 *Plant Nutrition*, 37, 671-678.
793 Zimmermann, M. (2001) Micronutrients in Health and Disease. Georg Thieme Verlag,
794 Stuttgart.
795
796