

Soil Fertility Status of Cassava Fields in South Western Nigeria

Salami B. T^{*}, Sangoyomi T. E

Department of Crop Production, Soil and Environmental Management,
Faculty of Agriculture, Bowen University, Iwo, Nigeria.

ABSTRACT

Aims: Current information on nutrient status of south western Nigeria soils is needed to develop appropriate integrated nutrient management packages for sustainable cassava production within the area. This study is designed to provide information on fertility status of some soils of the area.

Study design: A field survey.

Place and Duration of Study: The study was carried out in March 2009 within two agro ecological zones of Osun state, south western Nigeria.

Methodology: A field survey of 33 farmers' fields in two agro ecological zones of Osun state namely; Iwo zone and Oshogbo zone was undertaken. Representative soil samples (0-20cm) were obtained from each field. Physical and chemical properties of soil samples were determined in the laboratory following established methods. Statistical analyses of soil data was carried out using appropriate techniques.

Results: The soils of the area are acidic (pH in water range: 5.4 -6.4) and 79% of the fields are deficient in soil organic matter. Nitrogen and phosphorus are below established critical limits for cassava production in half of the fields; exchangeable cations (calcium, magnesium and potassium) are present in adequate amounts in most soils. No significant differences (P .05) were observed between the zones for soil properties measured.

Conclusion: Farmer acceptable strategies for improving nutrients availability (particularly nitrogen and phosphorus), organic matter and overall soil health through use of fertilizers, organic materials and multipurpose legumes among other options are required for sustained cassava production.

Keywords: [Nitrogen, phosphorus, mehlich soil test, micro nutrients, fertilizer, organic material]

1. INTRODUCTION

Cassava (*Manihot esculenta*) is an important staple crop for more than 700 million people in the developing world, about 500 million of whom reside in Africa (FAO, 2003; El-Sharkawy, 2004). The majority of production is in Africa and Nigeria ranks as the largest producer of cassava worldwide, harvesting over 35 million tonnes of fresh roots from 3.1 million hectare of land (FAO, 2011).

Cassava production in Nigeria is mainly in the hands of small scale farmers under rain fed conditions. The crop plays a vital role in the food security of the rural economy of Nigerians because of its ability to tolerate drought and give reasonable yield in soils of low fertility (Ezedinma et al., 2006), hence the name 'poor man's crop'. Therefore, cassava is most often grown in marginal soils and comes last in a crop rotation cycle (Nweke et al., 2002; Adjei-Nsiah et al., 2004). Population pressure on land has resulted in reduced fallow periods, more intensive land use and increasing problems of soil infertility (Rao and Mathuva, 2000). In

addition, fertilizer is scarce and priced much higher than the average farmer can afford thereby effectively limiting its use in cassava cultivation (FAO, 2005). Constant depletion and little or no external addition of plant nutrients impoverish the soil, resulting in declining yields and soil degradation. Average farm yield of cassava in Nigeria is lower (10.8 t ha^{-1}) compared to India ($> 27 \text{ t ha}^{-1}$) (FAO, 2011). In East Africa, using integrated management practices increased average yields of cassava up to 20 t ha^{-1} on farmers' fields and 60% of the increase was attributed to fertilizer use (Fermont, 2009). Therefore, soil fertility is a major constraint in cassava production (Spittel and Van Huis, 2000) and must be overcome if the Federal government of Nigeria backed "cassava initiative" for rural industrial development and economic empowerment will be realized.

An important step in achieving this is an inventory providing valuable information on the current fertility status of cassava growing soils and appropriate zonal specific soil nutrient management options. This study was designed to begin an assessment of the fertility status of soils under cassava in Osun state, Nigeria.

2. MATERIAL AND METHODS

2.1 Study Site

Osun state (7.5000° N , 4.5000° E) in south western Nigeria has a tropical climate with distinct wet and dry seasons. It is characterized by a bimodal rainfall pattern and 7- 8 months of rainfall, permitting the cultivation of long growing crops like cassava in a predominantly rain fed system. The mean annual temperature varies between 21°C and 31°C and annual rainfall ranges from 800 mm in the Savannah agro-ecology to 1,500 mm in the rain forest belt. The population of the state was estimated at slightly below 3.5 million from the last population census in 2006. More than half the population of the state is directly or indirectly involved in farming.

Cassava is an important crop occupying 69, 353 ha of land or 45% of the total land area cultivated in the state (Osun State Government, 2012). The state is divided into three agricultural zones based on its agro-ecological and cultural characteristics namely; Iwo, Oshogbo and Ife/Ijesha. Although cassava is grown in all three zones, it is relatively less important in Ife/Ijesha zone where tree based farming of cocoa (*Theobroma cacao*) and kola nut (*Cola nitida*) mainly predominates. Arable crop farming is predominant in the other two zones and cassava is an important component in cropping systems. Therefore, Iwo and Oshogbo zones were selected for the survey (Figure 1). A brief description of each of the zones studied is given as follows:

Iwo zone (Forest/savannah transition)

This zone is characterized by derived savanna vegetation with rainfall averaging $1100 \text{ mm per annum}$. The soils of the area are mainly derived from undifferentiated basement complex rocks with pebble beds and to a lesser extent coarse granite parent material. The topography is mainly undulating plains, dissected plains and patches of nearly level to gently undulating plains.

Farmlands may be acquired by inheritance, purchase, leasing etc. Typically, farm size ranges between 0.5-2 ha; farmers may have multiple farm plots (fragmented farm holdings). Land management options commonly practiced within the study area include the use of cover crops, multiple cropping, crop rotations, growing nitrogen fixing legumes and fallow. With shrinking land resource, fallow periods have become progressively shorter in length and farmlands are intensively cultivated for up to a decade or more before being allowed to fallow. Inorganic fertilizer use in Osun state is estimated at 11.3 kg ha^{-1} (Salimonu, 2008).

Comment [U1]: Or Salimonu. It is not the same in the References section.

Arable crops cultivated include maize (*Zea mays*), cassava, okra (*Hibiscus esculentus*) and chilli pepper (*Capsicum* sp.).

Oshogbo zone (Savanna zone)

This zone is located at the northern part of the state and it is characterized by savanna vegetation. The topography is mainly gently undulating plains with undulating dissected plains. Annual rainfall is about 800 mm per annum. Average farm size is between 0.5 and 1.5 ha, and farmlands are typically inherited, being passed on to succeeding generations. They may also be purchased outright or leased, payment being cash or a share of the farm produce.

Apart from the use of fertilizer, other soil fertility management options adopted within the area include crop rotation, intercropping legumes with other complementary crops, use of organic manures, mulching with organic residues and short duration bush fallows less than four (4) years in length. The soils of the area are derived mainly from quartzite and basement complex rocks. The main crops cultivated are maize, cassava, okra, soybean (*Glycine max*) and cowpea (*Vigna unguiculata*).




Figure 1. Map of Osun State, Nigeria with study area shaded. Inset is map of Nigeria with Osun state highlighted

2.2 Soil Sampling and Laboratory Analyses

Cassava is a major crop grown for food and cash in both zones. In most of the sites selected for soil sampling, cassava was grown as a sole crop. On a few sites, it was intercropped with maize, okra or chilli peppers. The location of the sites was arrived at in conjunction with the

state agricultural development program (ADP) personnel. A host of factors were considered namely; cassava production potentials of the locations, accessibility by road and approximate distribution of soil types. A total of 33 sites were sampled. The survey was carried out in March 2009.

Soil samples were taken from fields in the following manner. Four cores were randomly sampled using a soil auger at 0-20 cm soil depth, and bulked to form a composite. The composites were air dried and ground to pass through a 2 mm sieve.

The soil samples were analyzed for texture (Bouyoucos, 1962), pH in water (1:1) and KCl (1:1). Total N was determined by the Micro-Kjeldahl method (Jackson, 1962) and available P was by Bray's P1 method (Bray and Kurtz, 1945). Exchangeable bases (K, Ca and Mg) and micronutrients (Cu and Zn) were extracted with Mehlich-3 solution (Mehlich, 1984). The ratio of soil to extraction solution used was 1:10.

The exchangeable acidity was determined by extracting 2g of soil with 20 ml of the extract and determined by titration with 0.01N ~~NaOH~~ ~~NaOH~~ using phenolphthalein indicator. Organic carbon (C) was determined using wet Walkey Black dichromate digestion method (Nelson and Sommers, 1982). Organic matter (OM) was derived from data on organic C as follows:

OM = Organic C*1.7

2.3 Statistical Analysis and Data Presentation

Descriptive statistics (mean, minimum and maximum values) of soil physical and chemical properties are presented. One way analysis of variance by soil zone was carried out on measured soil properties. Correlation and regression analyses were performed on the soil data. All computations were carried out using Microsoft Excel.

3. RESULTS AND DISCUSSION

3.1 Acidity, Organic Matter and Macronutrients Status of Soils of Iwo and Oshogbo Zones

Soil pH in water varied among sites from 5.4 to 6.4 with a coefficient of variation of 4.3% (Table 1). The soils were moderately to slightly acidic. Cassava is tolerant of soil acidity (Howeler, 2002) and so acidity would not severely constrain production since all of the soils were within adequate levels of pH for cassava production (Table 2).

Two key inter-related properties to be considered in soil fertility maintenance and build up are the soil organic matter and soil organic carbon (SOC). Organic matter improves soil chemical properties in three ways: as a net source of carbon and nutrients, increases cation exchange capacity and stimulates biological activity (Asadu et al., 1997). Soil organic carbon is the major component of soil organic matter and plays a vital role in plant nutrient supply, determines response to N and P fertilizer and improves soil physical structure and processes (Motavalli et al., 1994). Declining SOC indicates soil degradation with resulting decline in crop yields (Sanchez et al., 1997). Tropical soils ~~have~~ typically ~~have~~ low levels of SOC, < 20 to 30 g kg⁻¹ (Bationo et al., 2007). In this study, the SOC of all soils ranged between 4.4 and 26.6 g kg⁻¹ (Table 1), while 86% of the soils had SOC less than 20 g kg⁻¹. Soil organic matter ranged from 0.7% to 4.5%, while 14% of the soil samples had organic matter above 4%. For a critical level of 3.0% organic matter for crop production in Nigeria (Akinrinde

and Obigbesan, 2000), 79% of the soils had low levels of organic matter (equally distributed among both zones) and a fifth of the soils (21%) had > 3.0% organic matter.

Total N values of the soils ranged from 0.5 g kg⁻¹ to 2.8 g kg⁻¹. Total N in fifty percent of the soils was lower than the critical value of 0.15% (Agboola and Corey, 1973), obtained for maize in some southwestern Nigerian soils. Linear regression analysis showed that a difference of 1 g kg⁻¹ in organic matter resulted in a difference of 0.5 g kg⁻¹ in total N (standard error = 0.066; r = 0.90). Building up the soil organic matter levels is a key to increasing soil N content and largely determines the fertility of soils of the tropics (Buresh and Giller, 1998; Yao et al., 2010).

Phosphorus extracted by Bray's P 1 solution varied widely, and ranged from 0.1 to 40.0 mg kg⁻¹ of P in the soil. Fifty percent of the soil samples were above the critical level for cassava production (8 mg kg⁻¹ P₁ from Table 2).

Table 1. Selected chemical properties of soils from Iwo and Oshogbo zones of Osun state, Nigeria (results combined and presented by zone), 2009.

Soil property	Measure of average	All soil samples	Iwo soils	Oshogbo soils
pH (water)	Mean	5.9	5.7	6.0
	Range	5.4-6.4		
pH (KCl)	Mean	5.4	5.5	5.5
	Range	5.0-5.8		
Organic carbon (g kg ⁻¹)	Mean	14.9	14.3	15.5
	Range	4.4-26.7		
Organic matter (%)	Mean	2.5	2.4	2.6
	Range	0.7-4.5		
Nitrogen (g kg ⁻¹)	Mean	1.5	1.4	1.6
	Range	0.5-2.8		
Phosphorus (mg kg ⁻¹)	Mean	12.2	10.5	13.9
	Range	0.1-40.0		
Exch. Acidity (cmol _c kg ⁻¹)	Mean	0.4	0.4	0.4
	Range	0.3-0.6		
Calcium (cmol _c kg ⁻¹)	Mean	2.6	2.9	2.3
	Range	0.7-7.2		
Magnesium (cmol _c kg ⁻¹)	Mean	1.5	1.3	1.7
	Range	0.5-4.1		

Formatted Table

Comment [U2]: Indicate the units;ii

Potassium ($\text{cmol}_c \text{ kg}^{-1}$)	Mean	0.4	0.3	0.4
	Range	0.2-0.7		

Table 2. Approximate soil fertility indices according to nutrient requirements for cassava and maize

Soil property	Method	Medium/critical level(range)	Country	Source
pH H ₂ O	(1:1)	4.6-7.8		CIAT (1979)
Organic matter		3%	Nigeria	Akinrinde and Obigbesan (2000)
P	Bray 1	8 mg kg ⁻¹	Nigeria	Kang et al. 1980
Ca*	Mehlich-3	0.3 $\text{cmol}_c \text{ kg}^{-1}$	Malawi	Wendt, 1995
Mg*	Mehlich-3	0.6 $\text{cmol}_c \text{ kg}^{-1}$	Malawi	Wendt, 1995
K	Mehlich-3	0.2 $\text{cmol}_c \text{ kg}^{-1}$	Malawi	Snapp, 1998
Zn	Mehlich-3	1.0 mg kg ⁻¹	Malawi	Wendt, 1995
Cu	Mehlich-3	0.5 mg kg ⁻¹	Malawi	Wendt, 1995

*Values for Ca and Mg divided by 200 and 120 respectively to convert from mg dm^{-3} to $\text{cmol}_c \text{ kg}^{-1}$.

The Mehlich-3 (M-3) soil extractant was used in the determination of basic cations and micronutrients. It is an efficient method for soil nutrient determination, especially in acid and neutral soils, since it can simultaneously extract multiple nutrients, saving time and money. The test method has been considerably researched with different crops and calibrated with other soils test methods, so that there is considerable information in literature (Hylkema, 2011). However, there exists no soil test calibration or interpretation for cassava, therefore general soil test calibration and interpretations for maize were used.

The mean of extractable K was 0.4 $\text{cmol}_c \text{ kg}^{-1}$ when data for both zones were combined, 0.3 $\text{cmol}_c \text{ kg}^{-1}$ for Iwo zone and 0.4 $\text{cmol}_c \text{ kg}^{-1}$ for Oshogbo zone (Table 1). Using a tentative M-3 K critical value of 0.2 $\text{cmol}_c \text{ kg}^{-1}$ for upland soils of Malawi (Table 2), K was at adequate levels in most soils. Some locations in Oshogbo had relatively higher K values and this might be due to the particular parent materials from which these soils were derived since these locations do not have a history of better fertilizer applications than other study locations (Sangoyomi and Ayandiji, unpublished data). Extractable Ca varied widely, and the range

was from 0.7 to 7.2 cmol_c kg⁻¹. Although mean values for Iwo and Oshogbo soils differed substantially, all soils had greater than the critical limit proposed for extractable Ca (Tables 1 and 2).

Extractable Mg ranged from 0.5 to 4.1 cmol_c kg⁻¹, with higher mean values in Oshogbo (Table 1). Magnesium appears to be sufficient throughout most of the study area, as 93% of the soils had Mg values above the proposed critical level of 0.6 cmol_c kg⁻¹ (Table 2). The study however reveals isolated areas of Mg deficiency. Response of cassava to Mg application in acid soils of southern Nigeria has been reported (Kang, 1984).

3.2 Particle Size Analysis and Micronutrient Status of Soils from Iwo and Oshogbo Zones

Studies on soil micronutrients levels for crop production are less commonly reported in literature. Tentative soil Zn and Cu critical values for maize production in upland soils of Malawi have been proposed (Wendt, 1995). Zinc in soil ranged from 2.9 mg kg⁻¹ to 11.0 mg kg⁻¹ (Table 3). Using the M-3 test, Chude et al (2003) proposed a tentative soil Zn level of 1.29 mg kg⁻¹, above which soil Zn is adequate for maize production in Samaru, northern Nigeria, only slightly higher than was obtained for some soils of Malawi (Table 2). Zinc appears sufficient within the study area. Soil Zn showed significant positive correlations with soil pH and extractable cations at P = .05 (Table 4).

Table 3. Micronutrients (copper, zinc) and physical properties of soils from Iwo and Oshogbo zones of Osun state, Nigeria (results combined and presented by zone), 2009.

Soil property	Measure of average	All soil samples	Iwo soils	Oshogbo soils
Copper (mg kg ⁻¹)	Mean	2.0	1.9	2.1
	Range	1.2-2.8		
Zinc (mg kg ⁻¹)	Mean	6.5	5.4	7.5
	Range	2.9-11.0		
Particle size (g kg ⁻¹)				
Sand	Mean	826	823	828
	Range	736-912		
Silt	Mean	129	131	128
	Range	40-214		
Clay	Mean	45	46	44
	Range	40-60		

Table 4. Correlations of zinc and copper with selected soil properties

Property	Zn	Cu
pH (H ₂ O)	0.62*	0.28 ^{ns}
pH (KCl)	0.72**	0.63*
Exch. Acid	-0.44 ^{ns}	-0.75**
Ca	0.65**	0.49 ^{ns}
Mg	0.60*	0.43 ^{ns}
K	0.57*	0.46 ^{ns}

Note: ^{ns}, * and ** denote not significant ($P < 0.05$), significant at ($P < 0.05$) and ($P < 0.01$) levels of significance, respectively.

The mean value of Cu was 2.0 mg kg⁻¹ overall, and mean values of 1.9 mg kg⁻¹ and 2.1 mg kg⁻¹ were recorded for Iwo and Oshogbo zones respectively (Table 1). Soil Cu ranged from 1.2 mg kg⁻¹ to 2.8 mg kg⁻¹. Table 3 presents correlation results of Cu with some soil properties. Copper content of soil significantly correlated with exchangeable acidity and pH (KCl), but not with pH (H₂O). The variation in soil Cu attributable to the two variables was 67% in both zones. Available Cu in these soils appears to be influenced by a combination of the acidity of the soil solution and reserved acidity in the soil colloids. The regression equation was {Cu = -0.33 + (0.72 * pHKCl X) + (-3.5 * Exch. Acid X)}. Available Cu was adequate in all soils.

Sand was the dominant particle fraction, comprising between 75 and 91% of total soil particle fraction. Mean values for sand in Iwo and Oshogbo were similar (Table 1). These results are typical of soils within this region generally described as sandy in texture and well drained (Salako et al., 1999). From soil texture analysis, most of the soils fall into the sandy loam class, with few exceptions; a lone field from Iwo zone classified as sand and another one from Oshogbo zone classified as loamy sand. All soils consisted of less than 10% clay by proportion.

3.3 Comparison of Soil Physical and Chemical Properties between Iwo and Oshogbo Zones

Statistical analysis ($P = .05$) indicated no significant differences between both zones for soil pH, organic matter, macronutrients (N, P, K, Ca, Mg) and exchangeable acidity. Similarly, differences in soil particle fractions, available Zn and extractable Cu were small between zones and did not prove statistically significant.

3.4 Soil Fertility Status, Fertilizer Use and the Quest for Increased Sustainable Cassava Production in Osun state

The success of the “cassava initiative” program is hinged on efficient and stable production and marketing channels to serve both domestic industries and export markets. To achieve this both land area under cassava and yield per hectare must increase. Even at the current rate of 2 million ha under cassava, a modest increase in national yield average to 15 t ha⁻¹ would increase production by 8 million tonnes per annum. Push factors such as government support, new varieties, better farming practices and farmer motivation are some typically cited means of increasing yields (Philips et al., 2004).

As high yielding cassava cultivars are increasingly adopted by farmers, nutrient mining accelerates and multiple nutrient deficiencies become increasingly common. Fermont et al. (2007) observed that when cassava tuber yield increased from 10 t ha⁻¹ when using farmer traditional varieties to 30 t ha⁻¹ with improved varieties, amounts of N, P and K removed per hectare tripled thereby aggravating the rapid depletion of soil nutrient stocks.

The International Institute of Tropical Agriculture (IITA), Nigeria, recommends application of any of the following fertilizers on a one hectare cassava farm: (1) NPK 15:15:15– 12 bags of 50 kg each; NPK 20:10:10– 9 bags of 50 kg each and NPK 12:12:17–15 bags of 50 kg each. This translates to about a fifth of the production cost (minus fixed expenditure) incurred on the farm for a projected yield of at least 25 t ha⁻¹ (IITA, 2005). Fertilizer use in Nigeria is quite low, estimated at 13 kg ha⁻¹ (FMARD, 2010), and of this projected amount, crops like maize are more likely to receive a greater proportion. In reality farmers hardly apply fertilizer to cassava (Nweke, 1994). Fertilizer procurement in Nigeria is plagued by a host of problems including poor distribution channels, inadequate supply, adulteration and high cost (Liverpool-Tassie et al., 2010). In addition, farmers are often unwilling to take the risk of incurring added cost from fertilizer which might not translate to monetary profit. Any of a combination of different factors such as time and number of weeding, pests and disease incidence, variety used and weather conditions (especially rainfall) could affect cassava yield. Agbaje and Akinlosotu (2004) observed that fertilizer application did not increase cassava tuber yield in seasons with excessive or inadequate rainfall at the active growth stage.

Since fertilizer use in cassava production is minimal at the moment, flexible approaches to fertilizer use must be developed. A blanket fertilizer recommendation often fails to take into consideration differences in resource endowment (soil type, labor capacity, climate risk) nor make allowances for dramatic changes in input/output price ration (Bationo et al., 2006), thereby deterring farmers from fertilizer application. Among cereal farmers in dry regions of West Africa, a method of applying small amounts (micro-dose) of fertilizer strategically to individual planting stations has been adopted. Fertilizer used is a third of the recommended rates and yield increased by 43% than with the earlier recommended fertilizer broadcasting methods (Tabo et al., 2007). Bationo and Buerkert (2001) explained that small amounts of fertilizers are more affordable to farmers, give an economically optimum result and increase nutrient uptake efficiency. The place of micro-dosing in cassava production needs to be assessed.

Combined use of inorganic and organic fertilizers for sustainable cassava production warrants being examined. Results from preliminary trials such as Ayoola and Makinde (2007) suggest that yield from organic and inorganic fertilizer combinations would not be inferior to sole inorganic fertilizer. Organic technologies that evaluate presently discarded crop residues, industrial wastes and other readily available organic materials together with mineral fertilizers for soil replenishment in cassava production are needed. Of particular interest and potential benefits are studies that focus on improving fertilizer use efficiency in organic and inorganic fertilizer combinations, soil N and P replenishment and building up the

Formatted: Justified

Formatted: Superscript

Comment [U3]: In the Reference section say Tassie, which one is ok?

Comment [F4]: Or Buerkert. Buerkert is the the References section

soil organic carbon stock through proper crop residue management. The potential contributions of legumes introduced (as relay, inter crop or rotation) in cassava systems and N fertilizer saving should be addressed. Quick growing legume planted fallows are options to be exploited. An added benefit of organic materials addition is that in sufficient amounts, it acts much like lime to reduce soil acidity and enhance nutrient availability in acid soils (Bessho and Bell,1992-1981). Farmers within the study area rarely apply lime to their soils although it is recommended (IITA, 2005).

For P, studies utilizing local phosphate rocks such as Ogun rock phosphate which is available in commercial quantities within south western Nigeria are needed. These studies would need to compare P use efficiency of cassava using phosphate rock or commercial fertilizers as P sources.

In as much as a single set of recommendation would not do for the diverse agricultural environments and economic conditions that prevail within the study area, technologies that explore composite approaches need to be further developed with the farmer fully participating. Here, extension officers allow farmers access to appraise and choose appropriate and cost-effective technologies suitable for their unique circumstances (Snapp et al., 1998; Gruhn et al., 2000).

Chemical soil testing is completely alien to traditional agriculture, but must come into forefront in commercial cassava production as soil reserves become depleted and multiple nutrient deficiencies show up. Traditional soil tests extract either one or a small group of nutrients, so that test for an extended range of nutrients requires multiple individual tests at high cost to farmers. However, the Mehlich-3 soil test provides a rapid and cost effective means of measuring up to 13 nutrients (B, Ca, Co, Cu, Fe, K, Mg, Mn, Mo, P, Na, S, and Zn) simultaneously (Walton and Alle, 2004). The test would greatly reduce cost and simplify soil test procedures. Therefore, studies focused on calibrating the soils tests for southern Nigeria soil types are needful.

4. CONCLUSION

Soils of Iwo and Oshogbo zones are mainly sandy loam and moderately to slightly acidic. The major soil constraints identified in this study are low levels of available P and organic matter, with the latter influencing a (which influences a host of soil physical, chemical and biological attributes e.g. available nitrogen supply, and cation exchange capacity.) and phosphorus. Innovative strategies for reducing inorganic fertilizer use by combining with organic means of fertilization are required for farmers' appraisal. The set of technologies adopted by farmers would vary and represent the most appropriate and cost effective methods for each farm. Finally, recent results on soil test calibration interpretation for cassava in these zones are required.

ACKNOWLEDGEMENTS

This study was carried out with funds provided by the Staff Development Grant, Bowen University, Iwo, Nigeria. The authors acknowledge the help of Mr S. Adio, Senior Research Officer, Osun state Agricultural development Program (OSSADEP) in identifying sites and gaining access to farmers' fields.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES.

- Adjei-Nsiah, S., Leeuwis, C., Giller, K.E., Sakyi-Dawson, O., Cobbina, J., Kuyper, T.W., Abekoe, M., Van der Werf, W. (2004). Land tenure and differential soil fertility management practices among native and migrant farmers in Wenchi, Ghana: implications for interdisciplinary action research. *NJAS – Wageningen Journal of Life Sciences*, 52(3-4), 331–348.
- Agbaje, G. O., Akinlosotu, T. (2004). Influence of NPK fertilizer on tuber yield of early and late planted cassava in a forest alfisol of south-western Nigeria. *African Journal of Biotechnology* 3 (10), 547-551.
- Agboola, A.A., Corey, R.B. (1973). The relationship between soil pH, organic matter, available P, exchangeable K, Ca, Mg and nine elements in the maize tissue. *Soil Science*, 115(5), 367-375.
- Akinrinde, E. A., Obigbesan, G. O. (2000). Evaluation of the fertility status of selected soils for crop production in five ecological zones of Nigeria. *Proceedings of the 26th Annual Conference of Soil Science Society of Nigeria*, Ibadan, Nigeria, pp. 279-288.
- Asadu, C.L.A., Diels, J., Vanlauwe, B. 1997. A comparison of the contributions of clay, silt, and organic matter to the effective CEC of soils of sub-Saharan Africa. *Soil Science* 162, 785-794.
- Ayoola, O.T., Makinde, E.A. (2007). Complementary Organic and Inorganic Fertilizer Application: Influence on Growth and Yield of Cassava/maize/melon Intercrop with a Relayed Cowpea. *Australian Journal of Basic and Applied Sciences*, 1(3), 187-192.
- Bationo, A., Buerkert, A. (2001). Soil organic carbon management for sustainable land use in the Sudano-Sahelian West Africa. *Nutrient Cycling in Agroecosystems* 61, 31-142.
- Bationo, A., Hartemink, A., Lungu, O., Naimi, M., Okoth, P., Smaling, E., Thiombiano, L. (2006). African Soils: Their Productivity and Profitability for Fertilizer Use. Background paper for the Africa Fertilizer Summit. 9-13 June 2006 in Abuja, Nigeria.
- Bationo, A., Kihara, J., Vanlauwe, B., Waswa, B., Kimetu, J. (2007). Soil organic carbon dynamics, functions and management in West African agro-ecosystems. *Agricultural Systems*, 94(1), 13-25.
- Bessho, T., Bell, L. C. (1992). Soil solid and solution phase changes and mung bean response during amelioration of aluminum toxicity with organic matter. *Plant Soil* 140, 183–196
- Bouyoucos, C.J. (1962). Hydrometer method improved for making particle size analysis of soil. *Agron. J.*, 54, 464-465.
- Bray, R. H., Kurtz, L.T. (1945). Determination of total, organic and available forms of phosphorus in soils. *Soil Sci.*, 59, 39-45.
- Buresh, R.J., Giller, K.E. (1998). Strategies to replenish soil fertility in African smallholder agriculture. In Waddington, S.R., Murwira, H.K., Kumwenda, J.D.T., Hukwa, D., Tagwira, F. eds. *Soil fertility research for maize-based farming systems in Malawi and Zimbabwe*. Harare: Soil Fertility Network and CIMMYT Zimbabwe. pp 13–19.
- Chude, V.O., Iwuofor, E.N.O., Amapu, I.Y., Pam, S.G., Yusuf, A.A. (2003). Response of maize to zinc fertilization in relation to Mehlich III extractable zinc. In B. Badu-Apraku et al. (ed.) *Proc. of a regional maize workshop IITA Cotonou, Benin Republic. 14–18 May 2001. WECAMAN/IITA*, pp 201-207
- CIAT (1979). *Annual Report for 1978*. CIAT Cali Colombia

Comment [F5]: This section must be in alphabetic and chronological order. There are some mistakes.

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Light Green

- 394 [El-Sharkawy, M.A. \(2004\). Cassava biology and physiology. Plant Mol. Biol. 56, 481-501](#)
- 395 [Ezedinma, C.I., Okafor, C., Asumugha, G.N., Nweke, F. \(2006\). Trends in farm labour](#)
 396 [Productivity and implications for cassava industrialization in Nigeria. Proceeding of](#)
 397 [the 40th Annual conference of the Agricultural society of Nigeria held at NRCRI](#)
 398 [Umudike, Abia State. Oct.16th - 20th, 2006, pp 109-111.](#)
- 399 [FAO \(2000\). FAOSTAT. Statistics division of the food and collections. Subset= Agriculture.](#)
 400 [Agricultural Organization,http://faostat.fao.org/](#)
- 401 [FAO \(2003\). FAOSTAT, FAO statistical database Agriculture, Food and Agricultural](#)
 402 [Organization http://faostat.Fao.org](#)
- 403 [FAO \(2005\). A review of cassava in Africa with country case studies on Nigeria, Ghana, the](#)
 404 [United Republic of Tanzania, Uganda and Benin volume 2. 357 pages](#)
- 405 [FAO \(2011\). FAOSTAT | © FAO Statistics Division 2011 | 12 July 2011](#)
 406 [\(http://faostat.fao.org/site/567\).](#)
- 407 [Fermont, A.M. \(2009\). Cassava and soil fertility in intensifying smallholder farming systems](#)
 408 [of East Africa. Published doctoral dissertation. Wageningen University and Research](#)
 409 [Centre, Wageningen.](#)
- 410 [Fermont, A.M., Obiero, H.M., van Asten, P.J.A., Baguma, Y., Okwuosa, E. \(2007\).](#)
 411 [Improved cassava varieties increase the risk of soil nutrient mining: an ex-ante](#)
 412 [analysis for western Kenya and Uganda. Advances in Integrated Soil Fertility](#)
 413 [Management in sub-Saharan Africa: Challenges and Opportunities. pp 511-520.](#)
- 414 [FMARD \(2010\). Federal Ministry of Agriculture and Rural Development: Annual Report](#)
- 415 [Gruhn, P., Goletti, F., Yudelman, M. \(2000\). Integrated Nutrient Management, Soil Fertility,](#)
 416 [and Sustainable Agriculture: Current Issues and Future Challenges. International](#)
 417 [Food Policy Research Institute 2033 K Street, N.W. Washington, D.C. 20006 U.S.A.](#)
 418 [September. Food, Agriculture, and the Environment Discussion Paper 32.](#)
- 419 [Howeler, R.H. \(1990\). Phosphorus requirements and management of tropical roots and](#)
 420 [tuber crops. In: Proceedings of the Symposium on P Requirements for Sustainable](#)
 421 [Agriculture in Asia and Oceania. March 6-8 1989. IRRI, Los Banos, Philippines pp](#)
 422 [427-444.](#)
- 423 [Howeler, R.H. \(2002\). Cassava mineral nutrition and fertilization. In: Hillocks, R.J., Thresh,](#)
 424 [M.J., Bellotti, A.C. \(Eds.\). Cassava: Biology, Production and Utilization. 115-147](#)
- 425 [Hylkema, A. \(2011\). Haiti soil fertility analysis and crop interpretations for principal crops in](#)
 426 [the five WINNER watershed zones of Intervention. Thesis report, Soil and Water](#)
 427 [Science, University of Florida.](#)
- 428 [IITA \(2005\). http://www.cassavabiz.org/about/ICP. Accessed on 1st August 2012.](#)
- 429 [Jackson, M.C. \(1962\). Soil Chemical Analysis. Prentice Hall Inc. Englewood Cliffs, New](#)
 430 [Jersey, USA.](#)
- 431 [Kang, B.T., Islam, R., Sanders, F.E., Ayanaba, A. \(1980\). Effect of phosphate fertilization](#)
 432 [and inoculation with VA-mycorrhizal fungi on performance of cassava \(Manihot](#)
 433 [esculenta crantz\) grown on an alfisol. Field Crops Research, 3, 83-94.](#)
- 434 [Kang, B.T. \(1984\). Potassium and magnesium responses of cassava grown in Ultisol in](#)
 435 [southern Nigeria. Fertilizer Research 5, 403-410.](#)
- 436 [Liverpool-Tasie, L.S.O., Olaniyan, B., Salau S., Sackey, J. \(2010\). A Review of Fertilizer](#)
 437 [Policy Issues in Nigeria. International Food Policy Research Institute \(IFPRI\), NSSP](#)
 438 [working paper 19, 29pp.](#)

Formatted: Font color: Green

Formatted: Font color: Green

Comment [F6]: This reference is not referred in the text. Delete it.

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Comment [F7]: This reference is not referred in the text. Delete it.

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font color: Light Green

Formatted: Font color: Green

Formatted: Font color: Green

441	Mehlich, A. (1984). Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant.	Formatted: Font color: Green
442	Commun. Soil Sci. Plant Anal., 15, 1409-1416.	
443	Motavalli, P.P., Palm, C.A., Parton, W.J., Elliot, E.T., Frey, S.D. (1994). Comparison of	Formatted: Font color: Green
444	laboratory and modelling simulation methods for estimating soil carbon pools in	
445	tropical forest soils. Soil Biology & Biochemistry 26, 935-944.	
446	Nelson, D.W., Sommers, L.E. (1982). Total carbon, organic carbon, and organic matter, p.	Formatted: Font color: Green
447	539-580. In A.L. Page et al. (ed.) Methods of soil Analysis. Part 2. 2nd ed. Agron.	
448	Monogr. 9. ASA and SSSA, Madison, WI	
449		
450	Nweke, F.I. (1994). Processing potentials for cassava production growth. COSCA Working	Formatted: Font color: Green
451	paper No 11. Collaborative study on Cassava in Africa, International Institute of	
452	Tropical Agriculture, Ibadan.	
453		
454	Nweke, F.I., Spencer, D.S.C., Lynam, J.K. (2002). The cassava transformation. Africa's	Formatted: Font color: Green
455	best-kept secret, Michigan State University Press, East Lansing, Mich, USA.	
456	Osun sState Government. (2012). www.osunstate.gov.ng. (Accessed 1st August 2012).	Formatted: Font color: Green
457	Phillips, T., Taylor, D.S., Sanni, L., Akoroda, M. (2004). A cassava industrial revolution in	Formatted: Font color: Green
458	Nigeria: the potential for a new industrial crop. IFAD/FAO, Rome, Italy. 43 pp.	
459		
459	Rao, M.R., Mathuva, M.N. (2000). Legumes for improving maize yields and income in semi-	Formatted: Font color: Green
460	arid Kenya. Agric. Ecosy. Environ. 78: 123-137.	
461		
461	Reichert, J.M., Albuquerque, J.A., Kaiser, D.R., Reinert, D.J., Urach, F.L., Carlesso, R.	
462	(2009). Estimation of water retention and availability in soils of Rio Grande do Sul.	
463	Revista Brasileira de Ciência do Solo (Impresso) 33, 1547-1560	Comment [F8]: This reference is not referred in the text. Delete it.
464	Salako, F. K., Babalola, O., Hauser, S., Kang, B. T. (1999). Soil macroaggregate stability	Formatted: Font color: Green
465	under different fallow systems and cropping intensities in southwestern Nigeria.	
466	Geoderma 91: 103-123.	Formatted: Font color: Green
467	Sanchez, P.A., Shepherd, K.D., Soule, M.J., Place, F.M., Mukwunye, A.U., Bursch, R.J.,	
468	Kwesiga, F.R., Izac, A.-M.N., Ndiritu, C.G., Woome, P.L. (1997). Soil fertility	
469	replenishment in Africa: an investment in natural resource capital. In: Bursch, R.J.,	
470	Sanchez, P.A., Calhoon, F. (Eds.), Replenishing soil fertility in Africa. Soil Science	
471	Society of America, Madison, WI, pp. 1-46.	
472	Snapp, S. S. (1998). Soil nutrient status of smallholder farms in Malawi. Communications in	Formatted: Font color: Light Green
473	Soil Science and Plant Analysis, 29(17), 2571-2588.	
474	Salimonu, K. K. (2008). Access to fertilizer subsidy among food crop farmers in Osun State,	Formatted: Font color: Green
475	Nigeria. International Journal of Agricultural Economics & Rural Development 1(2),	
476	1-8.	
477	Snapp, S.S., Mafongoya, P.L., Waddington, S. (1998). Organic matter technologies to	Formatted: Font color: Green
478	improve nutrient cycling in smallholder cropping systems of southern Africa. Agric.	
479	Ecosyst. Environ. 71, 187-202.	
480	Spittel, M.C., Van Huis, A. (2000). Effect of cassava mosaic disease, soil fertility, plant	Formatted: Font color: Green
481	spacing and their interactions on cassava yields in Zanzibar. Int. Journal of Pest	
482	46(3)-2000, 187-193.	
483	Tabo, R., Bationo, A., Gérard, B., Ndjounga, J., Marchal, D., Amadou, B., Anou, M.C.,	Formatted: Font color: Green
484	Sogodogo, D., Taonda, J.B.S., Hassane, O., Diallo, M.K., Koala, S. (2007).	
485	Improving cereal productivity and farmers' income using a strategic application of	
486	fertilizers in West Africa. In: Bationo et al., (ed) Advances in integrated soil fertility	
487	management in Sub-Saharan Africa: challenges and opportunities, pp 201-208,	
488	Springer, NL.	
489	Walton, K., Alle, D. (2004). Mehlich No. 3 Soil Test - The Western Australian Experience.	Formatted: Font color: Green
490	SuperSoil 2004: 3rd Australian-New Zealand Soils Conference, 5-9 December	

- 2004, University of Sydney, Australia. Published on CDROM. Website www.regional.org.au/au/asssi/
- Wendt, J.W. (1995). Evaluation of the Mehlich 3 soil extractant for upland Malawi soils. *Communications in Soil Science and Plant Analysis* 26 (5-6), 687-702.
- Yao, M.K., Angui, P.K.T., Konaté, S., Tondoh, J.E., Tano, Y., Abbadie, L., Benest, D. (2010). Effects of land use types on soil organic carbon and nitrogen dynamics in mid-west Côte d'Ivoire. *European Journal of Scientific Research* (40), 211-222.
1. Adiei-Nsiah S, Leeuwis C, Giller KE, Sakvi-Dawson O, Cobbina J, Kuyper TW. et al. Land tenure and differential soil fertility management practices among native and migrant farmers in Wenchi, Ghana: implications for interdisciplinary action research. *NJAS—Wageningen Journal of Life Sci.* 2004;52(3-4):331-348.
 2. Agbaje GO, Akinlosotu T. Influence of NPK fertilizer on tuber yield of early and late-planted cassava in a forest alfisol of south-western Nigeria. *African Journal of Biotechnology*. 2004;3(10): 547-551.
 3. Agboola AA, Corey RB. The relationship between soil pH, organic matter, available P, exchangeable K, Ca, Mg and nine elements in the maize tissue. *Soil Science*. 1973;115(5):367-375.
 4. Akinrinde EA, Obigbesan GO. Evaluation of the fertility status of selected soils for crop production in five ecological zones of Nigeria. *Proceedings of the 26th Annual Conference of Soil Science Society of Nigeria*, Ibadan, Nigeria. 2000:279-288.
 5. Asadu CLA, Diels J, Vanlauwe B. A comparison of the contributions of clay, silt, and organic matter to the effective CEC of soils of sub-Saharan Africa. *Soil Science*. 1997;162:785-794.
 6. Ayoola OT, Makinde EA. Complementary organic and Inorganic fertilizer application: Influence on growth and yield of cassava/maize/melon intercrop with a relayed cowpea. *Australian Journal of Basic and Applied Sciences*. 2007;1(3):187-192.
 7. Bationo A, Buerkert A. Soil organic carbon management for sustainable land use in the Sudano-Sahelian West Africa. *Nutrient Cycling in Agroeco*. 2001;61:131-142.
 8. Bationo A, Hartemink A, Lungu O, Naimi M, Okoth P, Smaling E. et al. African soils: Their productivity and profitability for fertilizer use. Background paper for the Africa Fertilizer Summit. Abuja, Nigeria. 2006
 9. Bationo A, Kihara J, Vanlauwe B, Waswa B, Kimetu J. Soil organic carbon dynamics, functions and management in West African agro-ecosystems. *Agricultural Systems* 2007;94(1):13-25.
 10. Bessho T, Bell LC. Soil solid and solution phase changes and mung bean response during amelioration of aluminum toxicity with organic matter. *Plant Soil*. 1992;140:183-196.
 11. Bouyoucos GJ. Hydrometer method improved for making particle size analysis of soil. *Agron. J.* 1962;54:464-465.
 12. Bray RH, Kurtz LT. Determination of total, organic and available forms of phosphorus in soils. *Soil Sci.* 1945;59: 39-45.
 13. Buresh RJ, Giller KE. Strategies to replenish soil fertility in African smallholder agriculture. In Waddington SR, Murwira HK, Kumwenda JDT, Hikwa D, Tagwira F. eds. *Soil fertility research for maize-based farming systems in Malawi and Zimbabwe*. Harare: Soil Fertility Network and cimmyt-Zimbabwe. 1998;13-19.
 14. Chude VO, Iwuafor ENO, Amapu IY, Pam SG, Yusuf AA. Response of maize to zinc fertilization in relation to Mehlich III extractable zinc. In Badu-Apraku B. et al. (ed.) *Proc. of a regional maize workshop IITA Cotonou, Benin Republic. 14-18 May 2001. WECAMAN/IITA*. 2003:201-207.
 15. CIAT. Annual Report for 1978. CIAT Cali Colombia.1979.
 16. El-Sharkawy MA. Cassava biology and physiology. *Plant Mol. Biol.* 2004;56:481-501.

Formatted: Font color: Green

Formatted: Font color: Green

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

- 543 17. Ezedinma CI, Okafor C, Asumugha GN, Nweke F. Trends in farm labor productivity
544 and implications for cassava industrialization in Nigeria. Proceeding of the 40th
545 Annual conference of the Agricultural society of Nigeria held at NRCRI Umudike,
546 Abia State. Oct.16th - 20th. 2006;109-111.
- 547 18. FAO. FAOSTAT, FAO statistical database Agriculture. Food and Agricultural
548 Organization. 2003. Available: <http://faostat.fao.org>.
- 549 19. FAO. A review of cassava in Africa with country case studies on Nigeria, Ghana, the
550 United Republic of Tanzania, Uganda and Benin. Vol. 2. 2005.
- 551 20. FAO. FAOSTAT | © FAO Statistics Division. 2011. Accessed 29 July 2012.
552 Available: <http://faostat.fao.org/site/567>.
- 553 21. Fermont AM. Cassava and soil fertility in intensifying smallholder farming systems
554 of East Africa. Published doctoral dissertation. Wageningen University and Research
555 Centre, Wageningen. 2009.
- 556 22. Fermont AM, Obiero HM, van Asten PJA, Baguma Y, Okwuosa E. Improved
557 cassava varieties increase the risk of soil nutrient mining: an ex-ante analysis for
558 western Kenya and Uganda. Advances in Integrated Soil Fertility Management in
559 sub-Saharan Africa: Challenges and Opportunities. 2007:511-520.
- 560 23. FMARD. Federal Ministry of Agriculture and Rural Development: Annual Report.
561 Nigeria. 2010.
- 562 24. Gruhn P, Goletti F, Yudelman M. Integrated nutrient management, soil fertility and
563 sustainable agriculture: Current issues and future challenges. International Food
564 Policy Research Institute 2033 K Street, N.W. Washington, D.C. 20006 U.S.A.
565 September. Food, Agriculture, and the Environment Discussion Paper 32. 2000.
- 566 25. Howeler RH. Cassava mineral nutrition and fertilization. In: Hillocks RJ, Thresh MJ,
567 Bellotti AC, editors. Cassava: biology, production and utilization. 2002;115-147.
- 568 26. Hylkema A. Haiti soil fertility analysis and crop interpretations for principal crops in
569 the five WINNER watershed zones of Intervention. Thesis report, Soil and Water
570 Science, University of Florida. 2011.
- 571 27. IITA. <http://www.cassavabiz.org/about/CP>. 2005. Accessed: 1 August 2012.
- 572 28. Jackson MC. Soil Chemical Analysis. Prentice Hall Inc. EngleWood Cliffs, New
573 Jersey, USA. 1962.
- 574 29. Kang BT. Potassium and magnesium responses of cassava grown in Ultisol in
575 southern Nigeria. Fertilizer Res. 1984;5:403-410.
- 576 30. Kang BT, Islam R, Sanders FE, Ayanaba A. Effect of phosphate fertilization and
577 inoculation with VA-mycorrhizal fungi on performance of cassava (*Manihot esculenta*
578 crantz) grown on an alfisol. Field Crops Res. 1980;3:83-94.
- 579 31. Liverpool-Tasie LSO, Olaniyan B, Salau S, Sackey J. A Review of Fertilizer Policy
580 Issues in Nigeria. International Food Policy Research Institute (IFPRI), NSSP
581 working paper 19. 2010.
- 582 32. Mehlich A. Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant.
583 Commun. Soil Sci. Plant Anal. 1984;15:1409-1416.
- 584 33. Motavalli PP, Palm CA, Parton WJ, Elliot ET, Frey SD. Comparison of laboratory
585 and modeling simulation methods for estimating soil carbon pools in tropical forest
586 soils. Soil Biology & Biochem. 1994;26:935-944.
- 587 34. Nelson DW, Sommers LE. Total carbon, organic carbon, and organic matter.
588 Methods of soil analysis. Part 2. 2nd edition. Agron. Monogr. 9. ASA and SSSA,
589 Madison, WI. 1982:539-580.
- 590 35. Nweke FI. Processing potentials for cassava production growth. COSCA Working
591 paper No 11. Collaborative study on cassava in Africa, International Institute of
592 Tropical Agriculture, Ibadan. 1994.
- 593 36. Nweke FI, Spencer DSC, Lynam JK. The cassava transformation. Africa's best-kept
594 secret, Michigan State University Press, East Lansing, Mich, USA. 2002.

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt, Font color: Green

Formatted: Indent: Left: 0.2", Hanging: 0.3"

Field Code Changed

Formatted: Font: (Default) Arial, 10 pt

Formatted: Hyperlink, Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt, Font color: Black

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

- 595 37. Osun State Government. Available: www.osunstate.gov.ng. (Accessed 1st August
596 2012). 2012.
- 597 38. Phillips T, Taylor DS, Sanni L, Akoroda M. A cassava industrial revolution in Nigeria:
598 the potential for a new industrial crop. IFAD/FAO, Rome, Italy. 2004.
- 599 39. Rao MR, Mathuva MN. Legumes for improving maize yields and income in semi-arid
600 Kenya. *Agric. Ecosy. Environ.* 200;78:123-137.
- 601 40. Sanchez PA, Shepherd KD, Soule MJ, Place FM, Mukwunye AU, Bursch RJ. Et al.
602 Soil fertility replenishment in Africa: an investment in natural resource capital.
603 Bursch RJ, Sanchez PA, Calhoon F, editors. Replenishing soil fertility in Africa. Soil
604 Science Society of America, Madison, WI. 1997;1-46.
- 605 41. Snapp SS. Soil nutrient status of smallholder farms in Malawi. *Communications in*
606 *Soil Science and Plant Analysis*. 1998;29(17): 2571 -2588.
- 607 42. Salimonu KK. Access to fertilizer subsidy among food crop farmers in Osun State,
608 Nigeria. *International Journal of Agricultural Economics & Rural Dev.* 2008;1(2):1-8.
- 609 43. Snapp SS, Mafongoya PL, Waddington S. Organic matter technologies to improve
610 nutrient cycling in smallholder cropping systems of southern Africa. *Agric. Ecosyst.*
611 *Environ.* 1998;71:187-202.
- 612 44. Spittel MC, Van Huis A. Effect of cassava mosaic disease, soil fertility, plant spacing
613 and their interactions on cassava yields in Zanzibar. *Int. Journal of Pest.*
614 2000;46(3):187-193.
- 615 45. Tabo R, Bationo A, Gérard B, Ndjéunga J, Marchal D, Amadou B, et al. Improving
616 cereal productivity and farmers' income using a strategic application of fertilizers in
617 West Africa. *Advances in integrated soil fertility management in Sub-Saharan Africa:*
618 *challenges and opportunities*. Springer, NL. 2007; 201-208.
- 619 46. Walton K, Alle D. Mehlich No. 3 Soil Test - The Western Australian Experience.
620 SuperSoil 2004. 3rd Australian New Zealand soils conference, 5 – 9 Dec. 2004,
621 University of Sydney, Australia. 2004. Accessed 29 July 2012. Available:
622 http://www.regional.org.au/au/asssi/supersoil2004/s9/oral/1615_waltonks.htm.
- 623 47. Wendt, JW. Evaluation of the Mehlich 3 soil extractant for upland Malawi soils.
624 *Communications in Soil Science and Plant Analysis*. 1995;26(5-6):687-702.
- 625 48. Yao MK, Angui PKT, Konaté S, Tondoh JE, Tano Y, Abbadie L, et al. Effects of land
626 use types on soil organic carbon and nitrogen dynamics in mid-west Côte d'Ivoire.
627 *European Journal of Scientific Res.* 2010;40:211-222.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Indent: Left: 0.2", Hanging: 0.3"

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial