

Original Research Article

Enzymatic and Nutritive Evaluation of Zinc and Organic Fertilization on Field Maize Yields in Islamabad - Pakistan

ABSTRACT

Low fertility and organic matter contents in soil limit the nutrient bio-availability for crops, potentially leading to poor nutrition in the animals and human beings that consume those crops. This field study on maize production employed integration of organic and chemical fertilizers to determine their impact on plant enzymatic activities in relation to crop production, and nutrient dynamics in plants and soil. The treatments compared were: control (no application of N from any source); 100% N from fertilizer (FN); 75% FN + 25% N from organic manure (ON); and 50% FN + 50% ON. Further, all these treatments were superimposed by three levels of zinc (Zn) fertilizer, viz., 0, 4, 8 kg Zn ha⁻¹. Conjunctive use of FN and ON with the ratio of 75+25 along with Zn application at the rate of 4 kg ha⁻¹ rendered the highest maize grain yield as compared to sole N application from chemical fertilizer. Similarly, enzymatic activities of nitrate reductase (NR) and superoxide dismutase (SOD) were significantly affected with combined use of N and Zn. Nitrogen, phosphorus and zinc contents in diagnostic leaves of maize were also higher in organically substituted treatments.

Keywords: Nutrient deficiencies; chemical nitrogen fertilizer; organic manure; conjunctive use; nitrate reductase; superoxide dismutase.

22 1. INTRODUCTION

23 Pakistani soils exhibit multiple nutrient deficiencies including that of zinc (Zn) due to their alkaline-
 24 calcareous nature and low organic matter content, nutrient mining by previous intensive cropping, and
 25 inadequate and imbalanced fertilizer use. Low soil organic matter in cultivated lands is a major cause of
 26 soil degradation; making these soils infertile and thus not capable of sustaining crop production (Diacono
 27 and Montemurro 2010). Nutrient mining has also raised concerns about the sustainability of agricultural
 28 systems. Continuous use of imbalanced synthetic fertilizers over time can lower crops yield (Sortino *et al.*
 29 2013) and also contaminates the environment (Hu and Qi 2013). Achieving a balance between nutrient
 30 requirements of the plants and nutrient supplies from the soil is essential for attaining and maintaining
 31 high crop yields and good soil fertility. Manures enhance the bio-availability of macro- and micro-
 32 nutrients, viz., N, P, K and Zn etc., and thus raise the concentration of these nutrients in plant tissues
 33 (Masti *et al.* 2003). Degraded lands can be ameliorated by conjunctive use of organic manures and
 34 inorganic fertilizers (Saleque *et al.* 2004). Appropriate blending of chemical fertilizers and organic
 35 manures varies with land-use pattern, agro-ecology, and socio-economic circumstances. Similarly, an
 36 organo-mineral combined fertilizer blend could be a promising alternative for sustaining the crop yields
 37 (Deeks *et al.* 2013). Improved management of crop residues / livestock manure and nutrients may also
 38 reduce the environmental impact of agriculture on climate change (Stavi and Lal 2013).

39 Nutrient deficiencies directly affect enzymatic activities in plants. Zinc deficiency in crops is a common
 40 disorder in most of the cropped lands in Pakistan (Rafique *et al.* 2006). It is mainly due to calcareous-
 41 alkaline soils, low organic matter, nutrient depletion, and injudicious fertilizer application. Zinc is a
 42 cofactor for one class of superoxide dismutase enzymes, which are important for detoxifying reactive
 43 oxygen species (ROS) and controlling free radicals (Cakmak 2000; Wang and Jin 2005), especially under
 44 drought stress. Similarly nitrate reductase is an important enzyme of N metabolism: its activity in plants is

inducible by nitrate (Shankar *et al.* 2001). Therefore, addressing N deficiency by the use of nitrate affects the activity of nitrate reductase in plants. Under rain-fed conditions maize production is lower than in irrigated areas, which is mainly due to water stress and poor soil fertility status. Continuous additions of organic amendments to cropland increase soil fertility, organic carbon, microbial biomass and enzymatic activities, and enhance crop yields (Diacono and Montemurro 2010). With these understandings, a field experiment on integrated nutrient management was conducted on rain-fed maize.

This paper reports the maize crop response (yield, nutrient uptake and enzyme activity) to nitrogen and zinc fertilizers application along with organic manure. It was hypothesized that there will be differences in these crop attributes between organic-N fertilizer combinations and sole mineral N fertilizer treatments. A major objective of this study was to reduce the reliance on chemical fertilizers and enhance the use of organic amendments for sustainable soil health and crop production.

2. MATERIALS AND METHODS

The field experiment was conducted using maize cv. Swan at the research farm of National Agricultural Research Centre (NARC), Islamabad (latitude 33° 43' N, longitude 73° 04' E, and altitude 490 m). The soil is classified in the Nabipur soil series (coarse loamy, mixed, hyperthermic Udic Ustochrept). It is deep, well drained, and moderately calcareous developed on level to nearly level deposition of the flood plain. It is located in a sub-humid to humid zone with medium to high rainfall, with annual rainfall ranging from 517 to 1550 mm. Mean maximum temperature during summer ranges from 33 to 40°C (WRRI 2010). Analysis revealed that original field soil was alkaline (pH, 7.8), calcareous (CaCO₃ equiv., 4.3%), low in organic matter (0.50%), and deficient in NO₃-N (3.5 mg kg⁻¹), available P (3.0 mg kg⁻¹), with extractable K (80 mg kg⁻¹) and Zn (0.27 mg kg⁻¹).

The experiment comprised of two factor treatments, viz., nitrogen sources combinations and zinc fertilization. The N application was made in main plots, while Zn was applied in sub-plots. Experimental design was split-plot under randomized complete block design (RCBD) with three replications. Plot size for main-plot treatments was 5 m × 9 m and each main-plot was divided into three equal sub-plots measuring 3 m × 5 m. Treatments detail is as follows:

Main plots - Nitrogen combinations

Control ≡ No application of N from any source

FN100 ≡ Whole N from chemical fertilizer (FN)

FN75+ON25 ≡ 75% FN + 25% N from organic manure (ON)

FN50+ON50 ≡ 50% FN + 50% ON

Sub plots - Zinc levels

Zn0 ≡ No application of zinc fertilizer

Zn4 ≡ Zinc fertilizer application at the rate of 4 kg ha⁻¹

Zn8 ≡ Zinc fertilizer application at the rate of 8 kg ha⁻¹

Nitrogen was applied at the rate of 120 kg ha⁻¹ as recommended dose for the maize crop in rainfed areas using different combinations of organic (farm yard manure) and fertilizer N (urea) sources. Fertilizer N (FN) was applied in two splits, viz., at seed bed preparation and at booting stage. Farm yard manure analysis showed 1.85% N, 0.43% P and 1.14% K contents. It was applied before planting at the rate of 3.3 t ha⁻¹ for replacing 50% of recommended mineral N, and at the rate of 1.82 t ha⁻¹ for 25% replacement of N fertilizer. Phosphorus fertilizer was equally applied at the rate of 50 kg ha⁻¹ in all the treatments at the time of sowing.

2.1 Soil and Plant Analysis

Before sowing the crop, composite soil samples were collected from 0-15 cm depth, air dried, ground and passed through 2 mm sieve. Soil samples were analyzed for pH (McLean 1982), AB-DTPA P

(Soltanpour and Workman 1979), electrical conductivity (ECe), (Rhoades 1982), organic matter (Nelson and Sommer 1982) NH₄OAC extractable K (Knudsen *et al.* 1982), NO₃-N and Zn (Soltanpour and Workman 1979). At blooming stage, recently fully matured leaves from selected plants were collected. The leaves were washed, cleaned, air dried and then oven dried at 65°C to a constant weight. The samples were ground, sieved and stored in small plastic bags for determination of N, P and Zn. Nitrogen was determined using the Kjeldahl method as described by Jackson (1973). Phosphorus was determined by wet digestion as described by Benton *et al.* (1991). Zinc was determined using atomic absorption spectrometer as described by Wright and Stuczynski (1996).

2.2 Measurement of Enzyme Activities

Samples of second youngest fully expanded leaves at booting stage (diagnostic leaves) were taken, immediately frozen in liquid N, ground to make fine powder, and refrigerated at –80°C. Activity of superoxide dismutase (SOD) was assayed spectrophotometrically at 560 nm by measuring its ability to inhibit photochemical reduction of nitro blue tetrazolium (Dhindsa *et al.* 1981). For nitrate reductase activity, fresh leaf samples were weighed and shifted to plastic vials. In each vial, 2.5 mL phosphate buffer (pH 7.5), 0.2 M KNO₃ and 5% isopropanol solutions were added, and samples were incubated for 2 h in dark at 30°C. After incubation, 1% sulphanilamide and 0.2% NED-HCl (N-1-naphthylethylene-diamine dihydrochloride) were added, and the vials kept for 20 min for colour development. Absorbance in samples was measured spectrophotometrically at 540 nm as described by Jaworski (1971).

2.3 Statistical Analysis

Data on growth, yield and nutrient uptake attributes of the crop were subjected to analysis of variance (Sokal and Rohlf 1997). Data showing significant difference at $P \leq 0.05$ were put to comparison of treatment means by least significant difference (LSD) test. All the data were processed using MSTAT-C software for statistical analysis.

113 3. RESULTS AND DISCUSSION

114 3.1 Maize Grain Yield

115 Pakistani soils exhibit multiple nutrient deficiencies including that of Zn (Rafique *et al.* 2006) due to their
 116 alkaline-calcareous nature and low organic matter content, nutrient mining with intensive cropping, and
 117 inadequate and imbalanced fertilizer use. Effect of experimental treatments on maize grain yield is shown
 118 in table 1. Treatment with a combination of fertilizer and organic N (FN75+ON25) produced significantly
 119 higher grain yield as compared to the FN50+ON50 combination and sole chemical N source (FN100). It
 120 shows a synergistic effect of organic and inorganic N sources. Secondly, the results of the study also
 121 showed positive interaction of N sources and Zn fertilization on maize productivity. The yield increase
 122 might be due to enhanced enzymatic activities with the application of Zn, and due to balanced and
 123 sustained supply of nutrients from organic manure. In case of 100% chemical N, the nutrients availability
 124 might not be sufficient for longer period. (Yadav *et al.* 2006) reported that compared to their sole
 125 application, integrated use of N, FYM and Zn is the best choice in term of crop yield, nutrient uptake, and
 126 economic return and benefit cost ratio. Shaheen *et al.* (2010) also observed enhanced crop yield and
 127 nutrient uptake through synergistic and improved fertilization practices.

128 3.2 Nitrogen Concentration in Diagnostic Leaf of Maize

129 Nitrogen concentration in the maize diagnostic leaf (third leaf below tassel) is given in table 2. Compared
 130 to that in control, N application significantly increased N concentration. Regarding the comparison of N
 131 sources, organic treatments, viz., FN75+ON25 and FN50+ON50 showed higher N contents than with sole
 132 N application (FN100). It reflects that N was available for long period to plants from organically
 133 substituted treatments. It might be due to synergistic effect of organic and inorganic sources on
 134 mineralization and sustained supply of essential nutrient: in various studies on integrated plant nutrient
 135 management (IPNM), it has been observed that integration of N sources, viz., organic and inorganic

origin increased NPK concentration in alfalfa and maize (Lioveras *et al.* 2004; Sial *et al.* 2007). Under the combined application, the decomposition process is enhanced by microbial activity, and energy is readily available from carbon for release of nutrients. Sial *et al.* (2007) also recorded higher N and P concentration in maize leaves and grain, when mineral N was applied in combination with organic sources.

Zinc fertilization also enhanced N concentration significantly in the maize diagnostic leaf over control, but Zn application rates of 4 and 8 kg ha⁻¹ were statistically similar. As referenced below, different physiological processes are inhibited in plants due to Zn deficiency. As a result, plant growth and yield were severely affected. Zinc plays fundamental role in critical cellular functions like protein metabolism, gene expression, bio membranes integrity, and IAA metabolism (Marschner 1995). It has also been reported that zinc application significantly increased protein and nitrogen content of maize (Tahir *et al.* 2009; Adiloglu and Saglam 2005).

Organically substituted N treatments along with Zn4 or Zn8 showed higher N contents in the diagnostic leaf as compared to that in sole N treatments with zinc, viz., FN100+Zn4 and FN100+Zn8. Overall, there was positive interactive effect on N content in maize. Verma *et al.* (2005) also recorded positive interactive effect of N and Zn on the growth, yield, quality and nutrient uptake of fodder sorghum.

3.3 Phosphorus Concentration in Diagnostic Leaf of Maize

Effects of N and Zn treatments on P concentration in the diagnostic leaf of maize are given in table 3. Nitrogen and Zn treatments significantly affected P contents. The highest P content was recorded in FN75+ON25 treatment, which was statistically higher than that with FN50+ON50 or FN100. Similar to N, Zn treatments have also increased the P content in the diagnostic leaf. Zinc application at 4 and 8 kg ha⁻¹ gave significantly higher P content than in control but were statistically alike with each other. Akinrinde *et al.* (2006) also reported that Zn application increased the P contents in maize tissues.

Interaction of N and Zn treatments was highly significant with respect to P concentration in diagnostic leaf samples. The highest P contents were recorded in organically substituted combinations of N and Zn, viz., FN75+ON25+Zn4, FN75+ON25+Zn8, FN50+ON50+Zn4 and FN50+ON50+Zn8. These treatments were statistically superior to all others including that of 100% fertilizer N application. These findings support the phenomena of higher nutrient availability under combined use of organic and inorganic fertilizers. Balanced application of plant nutrients and IPNM enhance crop yields (Dong *et al.* 2006).

3.4 Zinc Concentration in Diagnostic Leaf of Maize

Influence of N and Zn application on Zn content in diagnostic leaf is reflected in table 4. Zinc contents in diagnostic leaf were significantly increased with N and Zn application. The treatment containing 50% N gave significantly higher Zn content than with FN75+ON25 and FN100, which were statistically alike. Similarly, Zn application also affected Zn content of plant tissues compared to that in control. All three levels of Zn application differed statistically. Maximum contents were observed in Zn8 level, which were significantly higher than in Zn4 level. Interactions of source of N and Zn were also found significant as related treatments with relative levels of organic sources of N. The highest Zn concentration was observed in FN50+ON50+Zn8 followed by FN50+ON50+Zn4, which are statistically similar but differed with all other combinations of Zn and N fertilization. Other researchers also reported that zinc content and uptake by maize increased with Zn application and attributed it to better root growth (Obrador *et al.* 2002). Concentrations of N and Zn were also increased in maize plants with FYM application (Gondek and Mazur 2005). Akinrinde *et al.* (2006) also found that application of cow manure + ZnSO₄ produced the highest plant shoot biomass and gave the highest Zn uptake by maize.

3.5 Activity of Superoxide Dismutase

Superoxide dismutase (SOD) is an antioxidant enzyme, which counters stress in plants. Zinc is the structural part of SOD, so activity was determined in leaves at booting stage to assess the effect of Zn

application on SOD. Data regarding SOD activity is shown in table 5. The effect of N application on SOD activity was significant over control. However, N sources were statistically at par w.r.t. SOD activity. The highest SOD activity was recorded in Z8 followed by Z4. The result is in line with the finding of (Alguacil *et al.* 2006), who reported that compost addition significantly enhanced the SOD and peroxidase (POD) activities in the shoots of *J. oxycedrus*. Zinc application at 8 kg Zn ha⁻¹ gave significantly higher activity than that with 4 kg Zn ha⁻¹ and control. Interaction of Zn and N treatments was also significantly positive. The highest SOD activity was recorded in FN50+ON50+Zn8 followed by FN75+ON25+Zn8, and FN75+ON25+Zn4 treatments, being statistically similar.

Organic manure treatments with Zn significantly increased the SOD activity. It might be due to the structural role of Zn in SOD enzyme providing positive effect on the activity of SOD. Activity of SOD in wheat leaves significantly increased in NO₃ and NH₄ treated plants as compared to N deficient wheat plants (Polesskaya *et al.* 2004). Zn is a cofactor of 300 different enzymes and proteins, and it is involved in cell division, nucleic acid metabolism and protein synthesis. Zinc application enhances the activities of antioxidant enzymes (Cakmak 2000). Wang and Jin (2005) also reported that Zn application raised the activity of SOD in maize leaves. Zinc enhances the SOD activity and reduces MDA contents, which is beneficial to counter the free radicals (Gao *et al.* 2001).

3.6 Nitrate Reductase Activity

Nitrate reductase is an important enzyme of N metabolism, its activity in plants is inducible by its substrate (NO₃). So, anything that affects nitrate availability will influence nitrate reductase activity (Pervaiz *et al.* 2002). Nitrogen application significantly enhanced nitrate reductase activity (NRA) over control even when N sources significantly differ in NRA with each other (table 6). The highest NRA was recorded in FN50+ON50, where N was applied in equal ratio from organic and inorganic sources. Similar to N application, Zn also significantly affected the NRA. Zinc application at 8 kg ha⁻¹ showed the highest

NRA, which was significantly higher than with 4 kg Zn ha⁻¹. Zinc fertilization showed a positive effect on NRA. Maximum NRA was observed in manure supplemented treatments with Zn application.

Soil microbial biomass and enzymatic activities increase with integrated nutrient management (INM), which ultimately activates the micronutrients bio-availability (Tejada *et al.* 2008). Interactive effect of N sources and Zn was also significant in both crops. The highest NRA was recorded in FN50+ON50+Zn8 followed by FN50+ON50+Zn4 and FN75+ON25+Zn8 treatments. Alguacil *et al.* (2006) also found that under drought conditions, only inoculation with exotic arbuscular mycorrhizae fungi increased the shoot and root NRA by 188 and 38%, respectively as compared to the plants without inoculation or compost application. Nardi *et al.* (2004) also reported that NRA correlated positively with organic N, and humus extract of FYM treated soil contained 42% higher NRA as compared to control. Soil microbial biomass also increases enzymatic activities in soil.

4. CONCLUSIONS

This study compared different combinations of organic and mineral nitrogen fertilizers with and without zinc application for maize growth, yield, nutrient uptake and enzymatic activities. Farmyard manure (25% N-basis) + 4 kg Zn ha⁻¹ performed better than N fertilizer alone (100%) for maize production. Integration of organic and inorganic sources of nitrogen improved N and P status in soil possibly because the farmyard manure also contained some quantities of nutrients other than nitrogen. Activity of antioxidative enzyme (superoxide dismutase) was enhanced with the application of zinc and FYM, indicating improved stress tolerance in plants. Further, enhanced activity of nitrate reductase under integrated N fertilization indicated balanced supply of nitrogen to the crop.

The study led to the conclusion that the synergistic use of nitrogen sources (FYM and chemical fertilizer at 25:75 N ratio) is advantageous over sole application of mineral fertilizer. Farm yard manure and zinc fertilization further enhance the crop growth, yield and antioxidative enzyme activity for stress tolerance.

228 Twenty percent increased maize yield with the above mentioned IPNM strategy makes the system
 229 economically incentive based.

230 LITERATURE CITED

231 Adiloglu S, Saglam MT. The effect of increasing nitrogen doses on the zinc content of maize plant in the
 232 soils of different properties. Pak J Biol Sci. 2005;8:905-909.

233 Akinrinde EA, Olubakin OA, Omotoso SO, Ahmed AA. Influence of zinc fertilizer, poultry manure and
 234 application levels on the performance of sweet corn. Agri J. 2006;1:96-103.

235 Alguacil M, Caravaca F, Diaz-Vivancos P, Hernandez JA, Roldan A. Effect of arbuscular mycorrhizae
 236 and induced drought stress on antioxidant enzyme and nitrate reductase activities in *Juniperus*
 237 *oxycedrus* L. grown in a composted sewage sludge-amended semi-arid soil. Plant Soil.
 238 2006;279:209-218.

239 Benton J, Benjamin W, Harry AM. Analysis Hand Books: A practical sampling, preparation and
 240 interpretation guide. Micro-macro Publishing, Inc. USA; 1991. pp 130.

241 Cakmak I. Possible role of zinc in protecting plant cells from damage by reactive oxygen species. New
 242 Phytologist. 2000;146:185-205.

243 Deeks LK, Chaney K, Murray C, Sakrabani R, Gedara S, Le MS, Tyrrel S, Pawlett M, Read R, Smith
 244 GH. A new sludge-derived organo-mineral fertilizer gives similar crop yields as conventional
 245 fertilizers. Agron Sustain Dev. 2013;33(3):539-549.

246 Dhindsa RS, Dhindsa P, Thorpe TA. Leaf senescence: correlated with increased levels of membrane
 247 permeability and lipid peroxidation and decreased levels of superoxide dismutase and catalase.

- 248 Exp Bot. 1981;32:93-101.
- 249 Diacono M, Montemurro F. Long-term effects of organic amendments on soil fertility. A review. Agron
250 Sustain Dev. 2010;30:401-422.
- 251 Dong J, Hengsdijk H, Dai TB, De Boer W, Jing Q, Cao WX. Long-term effects of manure and inorganic
252 fertilizers on yield and soil fertility for a winter wheat-maize system in Jiangsu, China.
253 Pedosphere. 2006;16:25-32.
- 254 Gao Z, Lin B, Zhou W. Effect of zinc nutrition on endogenous plant hormones and free radicals in spring
255 maize. Plant Nutr Fert Sci. 2001;1:424-428.
- 256 Gondek K, Mazur BF. The effect of mineral treatment and the amendments by organic and organomineral
257 fertilizers on crop yield, plant nutrient status and soil properties. Plant Soil Environ. 2005;51:34-
258 45.
- 259 Hu C, Qi Y. Effective microorganisms and compost favor nematodes in wheat crops. Agron. Sustain.
260 Dev. 2013;33:573-579.
- 261 Jackson, ML Soil Chemical Analysis. Prentice Hall of India Limited, New Delhi; 1973.
- 262 Jaworski EG. Nitrate reductase assay in intact plant tissues. Biochem Biophys Res Commun. 1971;43:
263 1274-1279.
- 264 Knudsen D, Peterson GA, Pratt PF. Lithium, sodium and potassium. In: Page AL, Miller RH, Keeney DR
265 (eds) Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties, 2ndEd. Soil Sci
266 Soc Amer, Madison, WI; 1982. pp 115-246.

- 267 Lioveras J, Aran M, Villar P, Ballesta A, Arcaya A, Vilanova X, Delgado I, Munoz F. Effects of swine
268 slurry on Alfalfa production and on tissue and soil nutrient concentration. *Agron J.* 2004;96:986-
269 991.
- 270 Marschner H. Mineral Nutrition of Higher Plants, (2nd Ed). Academic Press, London; 1995. pp 889.
- 271 Masti T, Lithourgidis AS, Gagianas AA. Effect over of injected liquid cattle manure on growth and yield
272 of winter wheat and soil characteristics. *Agron J.* 2003;95:592-599.
- 273 McLean EO. Soil pH and lime requirement. In: Page AL, Miller RH, Keeney DR (eds) *Methods of Soil*
274 *Analysis. Part 2 - Chemical and Microbiological Properties* (2nd Ed). Agronomy. 1982;9:199-223.
- 275 Nardi S, Morari F, Berti A, Tosoni M, Giardini L. Soil organic matter properties after 40 years of
276 different use of organic and mineral fertilizers. *Europ J Agron.* 2004;21:357-367.
- 277 Nelson DW, Sommer LE. Total carbon, organic carbon, and organic matter. In: Page AL (ed) *Methods of*
278 *Soil Analysis.* 2nd Ed. Amer. Soc. Agron. Madison, WI; 1982. pp 539-579.
- 279 Obrador A, Alveraz JM, Fernandez MD, Lopez-valdivia LM. Changes with time of zinc forms in a acid, a
280 neutral and a calcareous soil amended with three organic zinc complexes. *Aust J Soil Res.*
281 2002;40:139-148.
- 282 Pervaiz Z, Afzal M, Xi X, Xiaoe Y, Ancheng L. Relationship among nitrate reductase activity, growth,
283 mineral contents and salt tolerance of wheat cultivars under saline condition. *Pak J Soil Sci.*
284 2002;21:67-72.
- 285 Polesskaya OG, Kashirina EI, Alekhina ND. Changes in the activity of antioxidant enzymes in wheat
286 leaves and roots as a function of nitrogen source and supply. *Russ J Plant Physiol.* 2004;51:615-

- 287 620.
- 288 Rafique E, Rashid A, Ryan J, Bhatti AU. Zinc deficiency in rainfed wheat in Pakistan: Magnitude, spatial
 289 variability, management, and plant analysis diagnostic norms. *Commun Soil Sci Plant Anal.*
 290 2006;37:181-197.
- 291 Rhoades JD. Soluble salts. In: Page A. et al. (ed) *Methods of Soil Analysis. Part 2. Agronomy.*
 292 1982;9:167-178.
- 293 Saleque MA, Abedin MJ, Bhuijan NI, Zaman SK, Panaullah GM. Long term effect of inorganic and
 294 organic fertilizer sources on yield and nutrient accumulating of low land rice. *Field Crops Res.*
 295 2004;86:53-65.
- 296 Shaheen A, Naeem MA, Jilani G, Shafiq M. Integrated soil management in eroded land augments the
 297 crop yield and water use efficiency. *Acta Agri Scand B - Plant Soil Sci.* 2010;60(3):274-282.
- 298 Shankar N, Khan SR, Srivastava HS. The response of nitrate reductase activity and nitrate assimilation in
 299 the maize roots to growth regulators at acidic pH. *Biologia Plant.* 2001;44:599-601.
- 300 Sial RA, Chaudhary EH, Hussain S, Naveed M. Effect of organic manures and chemical fertilizers on
 301 grain yield of maize in rain- fed area. *Soil Environ.* 2007;26:130-133.
- 302 Soltanpour PN, Workman S. Modification of the $\text{NH}_4\text{HCO}_3\text{DTPA}$ soil test to omit carbon black.
 303 *Commun Soil Sci Plant Anal.* 1979;10:1411-1420.
- 304 Sokal RR, Rohlf FJ. *Biometry: The Principles and Practice of Statistic in Biological Research.* WH
 305 Freeman, New York; 1997.

- Sortino O, Dipasquale M, Montoneri E, Tomasso L, Avetta P, Prevot AB. 90 % yield increase of red pepper with unexpectedly low doses of compost soluble substances. *Agron Sustain Dev.* 2013;33:433-441.
- Stavi I, Lal R. Agriculture and greenhouse gases, a common tragedy. A review. *Agron Sustain Dev.* 2013;33:275-289.
- Tahir M, Fiaz N, Nadeem MA, Khalid F, Ali M. Effect of different chelated zinc sources on the growth and yield of maize (*Zea-mays* L.). *Soil Environ.* 2009;28:179-183.
- Tejada M, Gonzalez JL, Garcia-Martinez AM, Parrado J. Application of green manure and green manure composted with beet vinasse on soil restoration: Effect on soil properties. *Bioresour Technol.* 2008;99:4949-4957.
- Verma S, Subehia SK, Sharma SP. Phosphorus fractions in an acid soil continuously fertilized with mineral and organic fertilizers. *Biol Fertil Soils.* 2005;41:295-300.
- Wang H, Jin JY. Photosynthetic rate, chlorophyll florescence parameters, and lipid peroxidation of maize leaves as affected by zinc deficiency. *Photosynthetica.* 2005;43:591-596.
- Wright RJ, Stuczynski T. Atomic absorption and flame emission spectrometry. In: Sparks DL (ed.) *Methods of Soil Analysis, Part 3. Chemical Methods.* SSSA; 1996. pp. 65-90.
- WRRI. Meteorological data for temperature, humidity, and average rainfall for Islamabad. Water Resources Research Institute, NARC, Islamabad, Pakistan; 2010.
- Yadav RP, Singh P, Aggarwal RK, Yadav MK. Improved technology for sustainable production of maize (*Zea mays* L.) in Shivalik foothills of Himachal Pradesh. *Indian J Dryland Agri Res Dev.*

326 2006;21:1-6.

327

328 **Tables**329 **Table 1: Maize grain yield (t ha⁻¹)**

330

Treatment	Nitrogen Source		Zn Levels (kg ha ⁻¹)			
	Input (%)		Zn1	Zn2	Zn3	Means (N)
	Mineral	Organic	0	4	8	
N1	00	00	2.51 g	3.07 f	3.28 e	2.95 D
N2	100	00	4.27 d	4.88 b	4.89b	4.68 C
N3	75	25	4.88 b	5.18 a	5.05 ab	5.04 A
N4	50	50	4.66 c	5.01ab	5.00 b	4.89 B
Means (Zn)			4.08 B	4.54 A	4.55A	

331

332 **LSD and CV values with different N sources and Zn levels**

333

Treatments		
N source	0.13	
Zn levels	0.08	
Nitrogen x Zinc	0.164	
C.V. (%)	2.20	

334

335 * Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike
 336 at $p \leq 0.05$.

337 ** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike
 338 at $p \leq 0.01$.

339 .

340

341

342

Table 2: Nitrogen concentration (g/100 g) in maize leaf

Treatment	Nitrogen Source		Zinc Levels (kg ha ⁻¹)			
	Input (%)		Zn1	Zn2	Zn3	Means (N)
	Mineral	Organic	0	4	8	
N1	00	00	0.95 f	0.97 f	0.98 f	0.97 C
N2	100	00	1.50 e	1.56 d	1.58 d	1.55 B
N3	75	25	1.707 e	1.75 abc	1.783 a	1.75 A
N4	50	50	1.72b c	1.77 ab	1.797 a	1.76 A
Means (Zn)			1.47 B	1.51 A	1.53 A	

LSD and CV values with different N sources and Zn levels

Treatments		
N source	0.036	
Zn levels	0.027	
Nitrogen x Zinc	0.055	
C.V. (%)	1.67	

* Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.05$.

** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.01$.

Table 3: Phosphorus concentration in maize leaf (g/100 g)

Treatment	Nitrogen Source Input (%)		Zinc Levels (kg ha ⁻¹)			
	Mineral	Organic	Zn1	Zn2	Zn3	Means (N)
			0	4	8	
N1	00	00	0.18 e	0.19 e	0.18 e	0.183 C
N2	100	00	0.213 d	0.257 ab	0.24 bc	0.237 B
N3	75	25	0.243 cd	0.26 bc	0.26 ab	0.254A
N4	50	50	0.23 abc	0.24 ab	0.25 a	0.24B
Means (Zn)			0.217B	0.237A	0.232A	

LSD and CV values with different N sources and Zn levels

Treatments		
N source	0.0115	
Zn levels	0.0086	
Nitrogen x Zinc	0.017	
C.V. (%)	8.78	

* Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.05$.

** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.01$.

Table 4: Zinc concentration (mg kg⁻¹) in leaf

Treatment	Nitrogen Source		Zinc Levels (kg ha ⁻¹)			
	Input (%)		Zn1	Zn2	Zn3	Means (N)
	Mineral	Organic	0	4	8	
N1	00	00	15NS	20	24	19C
N2	100	00	16	22	27	21B
N3	75	25	19	26	27	24A
N4	50	50	20	25	30	25A
Means (Zn)			17C	23B	27A	

LSD and CV values with different N sources and Zn levels

Treatments		
N source	1.68	
Zn levels	1.62	
Nitrogen x Zinc		
C.V. (%)	8.19	

* Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.05$.

** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.01$.

NS Means of N sources, Zn levels or their interactions have statistically non significant difference at $p \leq 0.05$.

Table 5: SOD (U g⁻¹ fresh wt) in maize leaf

Treatment	Nitrogen Source		Zinc Levels (kg ha ⁻¹)			
	Input (%)		Zn1	Zn2	Zn3	Means (N)
	Mineral	Organic	0		8	
N1	00	00	20.41 d	28.13 c	37.16 a	28.57 B
N2	100	00	25.76 c	33.52 b	37.93 a	32.40 A
N3	75	25	27.58 c	39.50 a	38.04 a	35.04 A
N4	50	50	28.81 c	36.78 a	38.57 a	34.72 A
Means (Zn)			25.64 C	34.48 B	37.92 A	

LSD and CV values with different N sources and Zn levels

Treatments		
N source	3.146	
Zn levels	1.601	
Nitrogen x Zinc	3.201	
C.V. (%)	5.66	

* Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.05$.

** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.01$.

Table 6: Nitrate reductase activity ($\mu\text{mole g}^{-1}$ fresh wt) in maize leaf

Treatment	Nitrogen Source		Zinc Levels (kg ha^{-1})			
	Input (%)		Zn1	Zn2	Zn3	Means (N)
	Mineral	Organic	0		8	
N1	00	00	1.02 h	1.04 h	1.03 h	1.03 D
N2	100	00	1.42 g	1.46 f	1.48 e	1.45 C
N3	75	25	1.51 d	1.55 c	1.56 bc	1.54 B
N4	50	50	1.57 b	1.58 d	1.60 a	1.58 A
Means (Zn)			1.38 C	1.41 B	1.42 A	

LSD and CV values with different N sources and Zn levels

Treatments		
N source	0.036	
Zn levels	0.008	
Nitrogen x Zinc	0.017	
C.V. (%)	1.54	

* Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.05$.

** Means of N sources, Zn levels or their interactions bearing same letter(s) are statistically alike at $p \leq 0.01$.