

# Soil Fertility as a Predictor of the Geospatial Distribution of Forest Species in Natural Regeneration in Brazil

## ABSTRACT

The relationship between soil attributes and spatial distribution variability of tree and shrub vegetation, specifically those in process of natural regeneration, can be an important tool for understanding the ecology of populations and communities while enabling the recommendation of species that can be used in restoration programs of degraded areas, as in this area from Atlantic Forest of Brazil. Thus, this work aimed to study soil chemical attributes as indicators of spatial distribution of forest species in natural regeneration with higher absolute density in the Lowlands Dense Ombrophilous Forest fragment in Pernambuco, Brazil. For soil chemical characterization, samples were collected at depths of 0.0-0.10 and 0.11-0.20 m. The natural regeneration species sampling was performed in 40 subunits of 25 m<sup>2</sup>, implemented in 40 sampling units of 250 m<sup>2</sup>. The individuals with diameter at breast height (DBH) < 15 cm and height ≥ 1.0 m were measured. Thus, the ten natural regeneration species with the highest Absolute Density were defined. Geostatistical analysis was used to characterize the spatial variability of forest species and soil attributes. *Tovomita mangle* may be indicated for soils with different chemical characteristics and natural fertility; *Brosimum rubescens* and *Inga capitata* for acidic soils with low natural fertility, but with moderate levels of exchangeable Al; *Talisia retusa* should be recommended for alic soils and *Caraipa densifolia* for non-alic soils; and *Anaxagorea dolichocarpa* and *Protium arachouchini* for high natural fertility soils.

**Keywords:** Atlantic forest, soil chemical attributes, soil/forest relationship, soil fertility, spatial distribution of vegetation.

## 1. INTRODUCTION

Atlantic Forest biome in Brazil is a continuous forest formation along the Brazilian coast region, extending from the northeast to the south. But, in general, as the colonization and occupation of Brazilian territory occurred initially in coastal regions, the biome was devastated, reducing its area to disjoint fragments of forest, mainly located in inhospitable sites of top discontinuous topography, slopes and hills [1].

So Atlantic Forest biome was reduced to approximately 12% of its original area in the state of Pernambuco, Brazil. It is represented by small forest fragments, isolated and surrounded by sugarcane monoculture, as is the case in the municipality of this study's region, which preserves only 10% (about 50.55 km<sup>2</sup>) of its original forest cover [2]. Some studies on these forest fragments have been carried out [3-6], adding important information about these remnants. Such information can contribute to the maintenance of the native flora populations of the biome, as well as subsidize recovery actions for degraded areas.

In this context, the soil is an important factor that interferes with the formation and understanding of the landscape. It has a fundamental role in the environment, providing mechanical support and nutrients for the development of vegetation. It also has direct or indirect relations with the different phytophysiognomies of a region, which allows natural occurrence of different forest formations, even in homogeneous regions in relation to other environmental factors [7, 8].

33 In order to evaluate the spatial variability of vegetation distribution as a function of soil attributes,  
34 geostatistical techniques can be used, which allows the interpretation of the data based on the natural  
35 variability structure of the evaluated attributes [9]. The use of geostatistical techniques in forest areas is  
36 very advantageous because it considers that the data obtained in the sample units are associated with  
37 their space location [10]. Therefore, it is possible to estimate variables in non-sampled areas using  
38 interpolators, when they present spatial dependence.

39  
40 The application of geostatistics techniques allows modeling and describing the spatial variability of  
41 vegetation distribution and soil attributes, which helps the elaboration of maps with a desirable level of  
42 detail, for a better understanding of these relationships [11].

43  
44 Several studies of spatial and temporal variation of soil properties have been conducted in Brazil [12],  
45 especially in planted forests, with the main objective of optimizing the management of production  
46 processes [11, 13, 14]. Some works were also developed in native forests [15-19]. However, studies on  
47 the relationship between soil attributes and variability of vegetation spatial distribution, especially in  
48 natural regeneration process, are scarce and can be important for understanding the ecology of  
49 populations and communities, subsidizing conservation strategies and/or sustainable management of tree  
50 species [20] in the Atlantic Forest biome.

51  
52 Thus, the objective of this **research** was to evaluate soil chemical attributes as indicators of geospatial  
53 distribution of forest species in natural regeneration with higher absolute density in a Lowlands Dense  
54 Ombrophilous Forest fragment in Pernambuco, Brazil. Thus, aiming to contribute to the recommendation  
55 of forest species in programs to recover degraded areas **from** Brazilian Atlantic Forest, under different soil  
56 conditions.

57

## 58 **2. MATERIAL AND METHODS**


59

### 60 **2.1 Study area**

61

62 The study was carried out in a fragment of the Lowlands Dense Ombrophilous Forest [21], with  
63 approximately 79 ha in Sirinhaém, Pernambuco, Brazil. Located under the following geographical  
64 coordinates: UTM 25L 259089 and 9053293; 259604 and 9053741; 259727 and 9052723; 259920 and  
65 9052956, with an average altitude of 63 m (Fig. 1). According to Köppen's classification, the region  
66 presents an AM monsoon climate [22], with an annual average temperature of 25.6 °C.

67



68  
69 **Fig. 1. Geographic location of the Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil.**

70  
71  
72 The rainfall data of Pernambuco State Agency for Water and Climate - APAC recorded an annual rainfall  
73 of about 1,800 mm [23]. Soils found in the region are Yellow **Oxisol**, Yellow **Ultisol**, Red-Yellow **Ultisol**,  
74 Gray **Ultisol**, **Entisol**, **Inceptisol** and **Fluvent** [24].

75  
76  
77 **2.2 Soil chemical attributes**

78  
79 For soil chemical characterization of the forest fragment, four simple samples were collected and  
80 homogenized, giving rise to a composite sample. They were sampled in 40 plots (10 m x 25 m) that were  
81 distributed systematically in the fragment. Samples were collected at two depths (0.0-0.10 m and 0.11-  
82 0.20 m). Soil sampling was concentrated where sampling of the natural regeneration of the species was  
83 carried out.


84  
85 The  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Al}^{3+}$  were extracted by 1.0 mol L<sup>-1</sup> KCl solution and determined by titration. P, K<sup>+</sup>, Fe,  
86 Cu, Zn and Mn were extracted by Mehlich-1 solution. P was determined by spectrophotometry, K<sup>+</sup> by  
87 flame photometry and Fe, Cu, Zn and Mn by atomic absorption spectrophotometry. Potential acidity  
88 (H+Al) was extracted by 0.5 mol L<sup>-1</sup> calcium acetate solution and determined by titration, and the total  
89 organic C (TOC) determination was performed by oxidation using the K dichromate method. With the  
90 results of these chemical analyzes, the sum of bases (SB), base saturation (V), saturation by Al (m),

91 effective cation exchange capacity ( $CEC_{\text{effective}}$ ), and potential cation exchange capacity ( $CEC_{\text{potential}}$ ) were  
92 all calculated [25].

### 93 94 **2.3 Natural regeneration**

95  
96 For the sampling of shrub-tree species of natural regeneration, 40 subunits of 25 m<sup>2</sup> (5 m x 5 m) were  
97 systematically allocated. These subunits were implemented on the right side of 40 sample units of 250 m<sup>2</sup>  
98 (10 m x 25 m), previously allocated in a permanent form to study the adult floristic composition of shrub-  
99 tree community, equidistant by 25 m and interspersed to the right and left (Fig. 2).

100



101  
102

103 **Fig. 2. Schematic diagram of the plots distribution in a fragment of the Lowlands Dense**  
104 **Ombrophilous Forest, Pernambuco, Brazil.**

105  
106 Natural regeneration studies were established based on the level of inclusion [26], with adaptations [27].  
107 The individuals with diameter at breast height (DBH) <15 cm and height ≥1 m were measured. The  
108 identification of species was done according to the APG (Angiosperm Phylogeny Group III) classification  
109 system [28]. With the data, ten natural regeneration species with the highest Absolute Density (AD) were  
110 defined using the following expression [29]:

$$AD = \frac{n_i}{A}$$

111 AD is the absolute density (ind. ha<sup>-1</sup>); n is the number of individuals of i species; and A is the sample area  
112 in hectares.

### 113 114 **2.4 Geostatistical procedures**

115  
116 Initially, a principal component analysis (PCA) was performed between the ten natural regeneration  
117 species with highest AD and the soil chemical attributes in the 0.0-0.10 m depth layer. This was done in  
118 order to discard variables that presented the lowest factor loads. After this procedure, geostatistical  
119 analysis was used to characterize the spatial variability of forest species and selected soil attributes.

120

121 Geostatistical procedures and correlations between soil attributes and geospatial distribution of the forest  
122 species were performed in the 0.0-0.10 m depth layer, as the species were in process of natural  
123 regeneration with the majority of the root system concentrated in the superficial layer. Besides, nutrient  
124 concentrations are higher on the soil surface.

125  
126 Kolmogorov-Smirnov test [30] was used to test the hypothesis of normality of the data, and geostatistical  
127 analysis was used to characterize the spatial variability [31]. Under the theory of intrinsic hypothesis, the  
128 experimental semivariogram was estimated by the equation [32]:

$$\hat{\gamma}(h) = \frac{1}{2N(h)} \sum_{i=1}^{N(h)} [Z(x_i) - Z(x_i+h)]^2$$

129  $\hat{\gamma}(h)$  is semivariance value for a distance  $h$ ;  $N(h)$  is the number of pairs involved in semivariance  
130 calculation;  $Z(x_i)$  is the value of  $Z$  attribute in position  $x_i$ ;  $Z(x_i+h)$  is the value of  $Z$  attribute separated by a  
131 distance  $h$  from the position  $x_i$ .

132  
133 Mathematical model with the calculated values of the semivariance was adjusted (spherical, exponential  
134 and Gaussian) and coefficients for the semivariogram were defined (nugget effect,  $C_0$ ; structural variance,  
135  $C_1$ ; sill,  $C_0 + C_1$ ; and range,  $a$ ). The nugget effect is the semivariance value for a distance greater than  
136 zero and smaller than the shortest sampling distance, which represents the random variation component;  
137 sill is the semivariance value at which the curve stabilizes over a constant value. When sill and nugget  
138 effect are found at similar levels, one has the pure nugget effect, or completely random behavior; and  
139 range is the distance from origin to where the sill reaches stable values, expressing the distance at which  
140 samples are not correlated [32].

141  
142 Semivariograms exam [33] was used in order to determine the spatial dependence (SD). In case of doubt  
143 among more than one model for the same semivariogram, was chose the best coefficient of determination  
144 ( $R^2$ ).

145  
146 The degree of spatial dependence of the variables was classified [34]. Semivariograms that had a nugget  
147 effect of less than or equal to 25% of the sill were considered to have strong spatial dependence,  
148 moderate when they were between 25% and 75%, and weak when they were higher than 75%.

149  
150 The kriging is the interpolated surface of each variable, which shows its spatial distribution. From kriging,  
151 it is possible to identify the location and extent of the extreme values, homogeneity degree of the area,  
152 and the highest gradient directions [35]. Using sampling optimization maps, information is obtained to  
153 better understand the spatial distribution pattern, and to define different distribution zones of forest  
154 species and soil chemical attributes. The maps of spatial distribution of studied variables were presented  
155 with five regular intervals of specific values for each variable, allowing a better distribution understanding,  
156 especially those with small intervals between maximum and minimum values.

157  
158 Pearson linear correlation coefficient [30] was used to evaluate the degree of correlation between soil  
159 chemical attributes and spatial distribution of forest species. In addition were also used analyses of the  
160 kriging maps of species distribution and variability of soil chemical attributes. Forest species or chemical  
161 attributes that showed pure nugget effect were disregarded.

### 162 3. RESULTS

#### 163 3.1 Predominant species in natural regeneration of the forest fragment

164  
165 Estimated absolute density of the natural regeneration of Lowlands Dense Ombrophilous Forest studied  
166 fragment was of 9,680 ind. ha<sup>-1</sup>. The ten species of highest AD represented 47.8% of the total sampled  
167 individuals in the area, with *Brosimum rubescens* having 1,500 ind. ha<sup>-1</sup> (Table 1). *Thyrsodium*  
168 *spruceanum* (580 ind. ha<sup>-1</sup>) was also one of the most abundant species in the study of natural  
169 regeneration, followed by *Tovomita mangle* (Table 1).

172  
173  
174  
175

**Table 1. Forest species of natural regeneration of higher Absolute Density (AD) and botanical families in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil**

Forest species	Family	AD (Ind. ha <sup>-1</sup> )
<i>Brosimum rubescens</i> Taub.	Moraceae	1,500 ± 82.17
<i>Thyrsodium spruceanum</i> Benth.	Anacardiaceae	580 ± 36.37
<i>Tovomita mangle</i> G. Mariz	Clusiaceae	560 ± 21.45
<i>Anaxagorea dolichocarpa</i> Sprague & Sandwith	Annonaceae	340 ± 14.24
<i>Eschweilera ovata</i> (Cambess.) Miers	Lecythidaceae	340 ± 12.91
<i>Protium arachouchini</i> March.	Burseraceae	280 ± 11.36
<i>Caraipa densifolia</i> Mart.	Calophyllaceae	280 ± 18.14
<i>Talisia retusa</i> R.S. Cowan	Sapindaceae	260 ± 10.75
<i>Inga capitata</i> Desv.	Fabaceae	250 ± 10.29
<i>Protium heptaphyllum</i> (Aubl.) Marchand	Burseraceae	240 ± 10.07

176  
177  
178  
179  
180  
181  
182  
183  
184  
185

### 3.2 Soil chemical attributes of the forest fragment

The forest **fragment soil** showed predominance of the sand fraction (Table 2), with small variations between the depths. However, they are medium textured soils, presenting a sandy clay loam class of textures in the depth of 0.0-0.10 m, and loam clay in the depth of **0.11-0.20 m** (Table 2).

**Table 2. Soil chemical attributes in depth of the fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil**

Soil attribute	Depth (m)	
	0.0-0.10	0.11-0.20
pH (H <sub>2</sub> O)	3.88 ± 0.23	4.15 ± 0.23
P (mg dm <sup>-3</sup> )	1.33 ± 0.52	1.20 ± 0.40
Ca <sup>2+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	0.47 ± 0.21	0.22 ± 0.13
Mg <sup>2+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	0.64 ± 0.32	0.52 ± 0.23
K <sup>+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	0.07 ± 0.04	0.05 ± 0.03
Al <sup>3+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	1.41 ± 0.36	1.22 ± 0.25
(H+Al) (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>1</sup>	6.10 ± 1.75	4.68 ± 1.39
TOC(g kg <sup>-1</sup> ) <sup>2</sup>	25.2 ± 0.88	18.0 ± 0.53
SB <sup>3</sup>	1.18 ± 0.39	0.79 ± 0.29
CEC <sub>effective</sub> (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>4</sup>	2.59 ± 0.42	2.01 ± 0.32

CEC <sub>potential</sub> (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>5</sup>	7.28 ± 1.74	5.47 ± 1.36
m (%) <sup>6</sup>	54.44 ± 11.85	60.70 ± 10.74
V (%) <sup>7</sup>	16.21 ± 7.09	14.44 ± 7.25
Fe (mg dm <sup>-3</sup> )	79.85 ± 26.82	75.98 ± 27.45
Cu (mg dm <sup>-3</sup> )	0.61 ± 1.79	0.43 ± 0.71
Zn (mg dm <sup>-3</sup> )	0.82 ± 0.88	0.61 ± 0.51
Mn (mg dm <sup>-3</sup> )	0.52 ± 0.65	0.45 ± 0.41
Total Sand (g kg <sup>-1</sup> )	481.60 ± 6.96	432.90 ± 5.50
Coarse Sand (g kg <sup>-1</sup> )	384.80 ± 6.46	335.90 ± 4.91
Fine Sand (g kg <sup>-1</sup> )	96.80 ± 1.31	97.10 ± 1.58
Silt (g kg <sup>-1</sup> )	252.70 ± 6.21	270.80 ± 8.54
Clay (g kg <sup>-1</sup> )	265.70 ± 4.95	296.30 ± 7.70
Textural class	Sandy clay loam	Loam clay

<sup>1</sup>Potential acidity; <sup>2</sup>Total organic carbon; <sup>3</sup>Sum of bases; <sup>4</sup>Effective cation exchange capacity; <sup>5</sup>Potential cation exchange capacity; <sup>6</sup>Saturation by aluminum; <sup>7</sup>Base saturation.

The soil of the fragment was classified of low natural fertility, due to its dystrophic character, considering base saturation (V) as a soil fertility indicator, which represents the sum of Ca<sup>2+</sup>, Mg<sup>2+</sup> and K<sup>+</sup> in relation to CEC<sub>potential</sub> (Table 2).

Low pH values prevailed at different depths of the soil of the fragment: 3.88 and 4.15 at depths of 0.0-0.10 and 0.11-0.20 m, respectively, as well as high (H+Al) concentration, characterizing high acidity (Table 2). It should be noted that the potential acidity (H+Al) was predominantly formed by H<sup>+</sup> ions, because the Al<sup>3+</sup> concentrations represented only 23.1% of the potential acidity (Table 2). The highest levels of Ca<sup>2+</sup>, Mg<sup>2+</sup>, P, K<sup>+</sup>, Al<sup>3+</sup>, (H+Al), SB, CEC<sub>effective</sub>, CEC<sub>potential</sub>, and V were concentrated in the first 0.10 m depth.

Saturation by aluminum (m) was lower in the superficial layer (54.44%) due to higher base saturation (V) in this layer (16.21%), and m was higher in subsurface layer (60.70%), where V was lower (14.44%) (Table 2). Due to high Al exchangeable concentration, higher than 50%, the soil was classified as alic.

### 3.3 Geospatial variability of soil chemical attributes and forest species distribution

Geospatial variability of soil chemical attributes and forest species of higher AD in natural regeneration, except for *Protium heptaphyllum*, presented spatial dependence (Tables 3 and 4).

**Table 3. Models and parameters of semivariograms of soil chemical attributes in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil**

Attribute	Mean	SDV <sup>6</sup>	Model	C <sub>0</sub> <sup>7</sup>	(C <sub>0</sub> + C <sub>1</sub> ) <sup>8</sup>	a (m) <sup>9</sup>	CD (R <sup>2</sup> ) <sup>10</sup>	SD(%) <sup>11</sup>	CV <sup>12</sup>
SB (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>1</sup>	1.18	0.39	Exponential	0.02	0.14	170.9	0.92	13.50	0.980
(H+Al) (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>2</sup>	6.10	1.75	Exponential	0.05	1.52	77.4	0.93	3.39	1.121
Mg <sup>2+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	0.64	0.32	Spherical	3.8 x 10 <sup>-3</sup>	0.07	74.1	1.00	5.31	0.902

K <sup>+</sup> (cmol <sub>c</sub> dm <sup>-3</sup> )	0.07	0.04	Gaussian	8.7 x 10 <sup>-4</sup>	632 x 10 <sup>-5</sup>	630.2	0.82	13.76	0.765
CEC <sub>potential</sub> (cmol <sub>c</sub> dm <sup>-3</sup> ) <sup>3</sup>	7.28	1.74	Spherical	0.25	2.52	396.0	0.99	10.08	0.968
V (%) <sup>4</sup>	16.21	7.09	Spherical	0.90	48.72	60.2	0.96	1.84	1.092
m (%) <sup>5</sup>	54.44	11.85	Exponential	20.60	133.20	50.1	0.82	15.46	0.212

<sup>1</sup>Sum of bases; <sup>2</sup>Potential acidity; <sup>3</sup>Potential cation exchange capacity; <sup>4</sup>Base saturation; <sup>5</sup>Saturation by aluminum; <sup>6</sup>Standard deviation; <sup>7</sup>Nugget effect; <sup>8</sup>Sill; <sup>9</sup>Range; <sup>10</sup>Coefficient of determination; <sup>11</sup>Spatial dependence = (Nugget effect)/(Range) x 100. (SD≤25% is strong; 25<SD<75% is moderate; and SD≥75% is weak); <sup>12</sup>Cross validation.

**Table 4. Models and parameters of semivariograms of forest species in natural regeneration with higher absolute density in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil**

Forest	Mean	SDV <sup>1</sup>	Model	C <sub>0</sub> <sup>2</sup>	(C <sub>0</sub> + C <sub>1</sub> ) <sup>3</sup>	a (m) <sup>4</sup>	CD (R <sup>2</sup> ) <sup>5</sup>	SD (%) <sup>6</sup>	CV <sup>7</sup>
<i>B. rubescens</i>	37.5	82.17	Spherical	204.5	6389	119.2	0.98	3.20	0.320
<i>T. mangle</i>	14.0	21.45	Spherical	83	742.6	130.8	1.00	11.17	0.240
<i>A. dolichocarpa</i>	8.5	14.24	Spherical	24.36 x	162.8	107.8	0.97	14.96	1.041
<i>P. arachouchini</i>	7.0	11.36	Spherical	6.44 x 10 <sup>3</sup>	79.58	129.2	0.97	8.11	1.344
<i>C. densifolia</i>	7.0	18.14	Spherical	129	715.4	81.4	0.99	18.03	0.293
<i>T. retusa</i>	6.5	10.75	Spherical	5.62 x 10 <sup>3</sup>	74.56	117.0	0.91	7.53	0.202
<i>I. capitata</i>	6.2	10.29	Spherical	56.1	112.3	122.0	0.98	49.95	0.719
<i>P. heptaphyllum</i>	6.0	10.07	Linear	-	-	-	-	PNE <sup>8</sup>	-

<sup>1</sup>Standard deviation; <sup>2</sup>Nugget effect; <sup>3</sup>Sill; <sup>4</sup>Range; <sup>5</sup>Coefficient of determination; <sup>6</sup>Spatial dependence = (Nugget effect)/(Range) x 100. (SD≤25% is strong; 25<SD<75% is moderate; e SD≥75% is weak); <sup>7</sup>Cross validation; <sup>8</sup>Pure Nugget Effect.

The pure nugget effect occurred to *Protium heptaphyllum*, indicating absence of spatial dependence. Therefore, for this species the shortest distance between sampling points (25 m) was not enough to detect the spatial variability among the samples (Table 4).

For the correlation between soil attributes and geospatial distribution of the species in the fragment, *Thyrsodium spruceanum* and *Eschweilera ovata* were also disregarded, as they presented the lowest factor loads in the principal components analysis.

Spherical and exponential models were the ones that best fit the semivariograms of soil chemical attributes, except for the K<sup>+</sup> concentration that conformed to the Gaussian model (Table 3). For studied species, semivariograms conformed to spherical model (Table 4).

The analysis of the relationship of spatial dependence degree showed that the species presented a strong spatial dependence, except for *Inga capitata*, which presented moderate dependence (Table 4).

All soil chemical attributes and forest species presented a coefficient of determination (R<sup>2</sup>) higher than 0.80 (Tables 3 and 4), at least 80% of the variability in estimated semi variance values were explained by adjusted models.

Considering the range, soil chemical attribute that presented the highest value was K<sup>+</sup> concentration (630 m), with the lowest range observed for saturation by Al (50 m) (Table 3). For species, the highest



245 range value was obtained for *Tovomita mangle* (131 m) and the lowest for *Caraipa densifolia* (81 m)  
246 (Table 4).


247

### 248 **3.4 Soil-vegetation relationship**

249


250 In order to study correlations of soil chemical attributes of the forest fragment and the distribution of  
251 natural regeneration species, kriging maps were elaborated with adjusted semivariograms models  
252 parameters (Fig. 3 and 4), and a Pearson correlation was performed (Table 5). **Spatial** distribution of  
253 *Brosimum rubescens* occurred throughout the fragment area (Fig. 4). However, it concentrated the largest  
254 number of individuals around 300 ind. ha<sup>-1</sup>, in a small region where more elevated values of SB, (H+Al),  
255 CEC<sub>potential</sub>, and exchangeable Mg were found (Fig. 3 and 4).

256



257  
258  
259  
260

Fig. 3. Kriging maps of the spatial distribution of soil chemical attributes in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil.



261  
 262  
 263  
 264  
 265  
 266

**Fig. 4.** Kriging maps of the spatial distribution of forest species in natural regeneration with higher Absolute Density (AD) in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil.

267  
268  
269  
270

**Table 5. Correlation matrix (Pearson) between soil chemical attributes and spatial distribution of forest species in natural regeneration with higher Absolute Density (AD) in a fragment of Lowlands Dense Ombrophilous Forest, Pernambuco, Brazil**

Forest species	Soil chemical attributes						
	SB <sup>1</sup>	(H+Al) <sup>2</sup>	Mg <sup>2+</sup>	K <sup>+</sup>	CEC <sub>potential</sub> <sup>3</sup>	V <sup>4</sup>	m <sup>5</sup>
	cmol <sub>c</sub> dm <sup>-3</sup>				%		
<i>Brosimum rubescens</i>	ns	0.389*	0.431*	ns	0.446*	ns	ns
<i>Tovomita mangle</i>	ns	ns	ns	-0.403*	ns	ns	ns
<i>Anaxagorea dolichocarpa</i>	ns	-0.315*	ns	ns	-0.340*	ns	ns
<i>Protium arachouchini</i>	ns	ns	ns	ns	ns	ns	ns
<i>Caraipa densifolia</i>	ns	ns	ns	ns	ns	ns	-0.338*
<i>Talisia retusa</i>	-0,390*	ns	-0.405*	ns	ns	ns	0.470*
<i>Inga capitata</i>	-0,436*	0.509*	ns	-0.451*	0.412*	-0.568*	ns

271 <sup>1</sup>Sum of bases; <sup>2</sup>Potential acidity; <sup>3</sup>Potential cation exchange capacity; <sup>4</sup>Base saturation; <sup>5</sup>Saturation by aluminum. Sig: Significance (T test: ns, \*  
272 Not significant or significant at P = 0.05, respectively).

273  
274 The species *Brosimum rubescens* populated areas with higher CEC<sub>potential</sub>, probably because soils are  
275 more clayey and/or have higher organic matter concentrations, but mostly composed by acid cations  
276 (H + Al) and with a preference for higher levels of exchangeable Mg in soil.

277  
278 The species *Tovomita mangle* presented a negative correlation with exchangeable K concentration in soil  
279 of the fragment (Table 5), also verified by spatial distribution of species by the kriging maps (Fig. 3 and 4).  
280 This indicated that the higher exchangeable K concentration of the soil (Fig. 3) is associated with the  
281 smaller number of individuals of this species (Fig. 4). Additionally, a small relation of *Tovomita mangle*  
282 occurrence, between 60 and 80 ind. ha<sup>-1</sup>, was observed in regions of the forest fragment that presented  
283 higher (H+Al) concentration (Fig. 3 and 4).

284  
285 Unlike the two previous species, *Anaxagorea dolichocarpa* showed a correlation with the availability of  
286 exchangeable K of the soil by the kriging maps. A higher number of individuals per hectare was also  
287 reported in areas with higher exchangeable K concentrations (Fig. 3 and 4), but this behavior was not  
288 significant through Pearson's correlation (Table 5). However, it was correlated with low values of  
289 CEC<sub>potential</sub> and low levels of (H+Al) (Table 5).

290  
291 The correlations performed to evaluate the *Protium arachouchini* spatial distribution did not identify any  
292 soil chemical attributes that were related to this species (Table 5). However, the kriging maps allowed to  
293 infer that the highest individuals per hectare concentration occurred in areas with higher Mg concentration  
294 and higher CEC<sub>potential</sub>, and, in a less expressive way, it presented a correlation with (H+Al) similar to the  
295 behavior presented by *Brosimum rubescens* (Fig. 3 and 4).

296  
297 *Caraipa densifolia* was correlated negatively with saturation by Al (Table 5), also found in the kriging  
298 maps of species spatial distribution and chemical attributes in the fragment area (Fig. 3 and 4). The  
299 spatial distribution pattern of *Caraipa densifolia* was antagonistic to that presented by *Talisia retusa*,  
300 which correlated negatively with SB and exchangeable Mg and positively with saturation by Al (Table 5).  
301 Thus, the highest number of individuals per hectare of *Talisia retusa* was found in areas with higher  
302 percentages of saturation by Al. This behavior was also identified in the kriging maps for *Talisia retus* and  
303 the soil chemical attributes of the fragment.

304

305 For spatial distribution of *Inga capitata* in the fragment area, its occurrence was verified where there were  
306 the lowest levels of exchangeable K, lower values of SB, and base saturation (Table 5 and Fig. 3 and 4).  
307 In the kriging maps it was also possible to observe that in the lower area, the eastern portion of the  
308 fragment, there is a greater concentration of individuals in the higher potential acidity region (H+Al) and  
309 higher CEC<sub>potential</sub>, confirming the positive correlation of this species with these attributes.

310

## 311 4. DISCUSSION

312

### 313 4.1 Predominant species in the natural regeneration of the forest fragment

314

315 Lopes [36] studying the dynamics of natural regeneration found *Brosimum rubescens* as one of the most  
316 abundant species in a fragment of Lowlands Dense Ombrophilous Forest, as in this study. *Thyrsodium*  
317 *spruceanum* was also one of the most abundant species found in the study of the natural regeneration of  
318 tree species conducted by Silva [37] in a fragment of Dense Ombrophilous Forest, Brazil.

319

320 According to Corrêa [38] *Tovomita mangle* is typically found in the more advanced successional stage of  
321 a forest. This species was abundant in the fragment. Therefore, its presence is an indicative that the  
322 successional process is evolving in the fragment area. *Eschweilera ovata*, *Anaxagorea dolichocarpa*, and  
323 *Protium heptaphyllum* were also highlighted with regards to the number of individuals in a study on  
324 natural regeneration developed by Lima [39].

325

326 Understanding the factors that interfere or contribute to the establishment of natural regeneration species  
327 is critical to ensuring the balance and sustainability of forest ecosystems.

328

### 329 4.2 Soil chemical attributes of the forest fragment

330

331 The pH influences vegetation development by interfering on the soil nutrient availability [39]. For Mafra  
332 [41] the acidification of the soil in forest areas can be related to the leaching of bases or absorption of  
333 these bases by the plants. Some studies also found soils in similar conditions in fragments of Dense  
334 Ombrophilous Forest of Pernambuco, Brazil, with pH values between 4.04 and 4.75 [42, 43].

335

336 Espig [42] and Teixeira [43] also found similar results for the concentrations of Ca<sup>2+</sup>, Mg<sup>2+</sup>, P, K<sup>+</sup>, Al<sup>3+</sup>,  
337 (H+Al), SB, CEC<sub>effective</sub>, CEC<sub>potential</sub>, and V, as in this study.

338

339 Concentrations of the bases Ca<sup>2+</sup>, Mg<sup>2+</sup>, and K<sup>+</sup> were considered low. However, they were same to  
340 results obtained by Teixeira [43], which found Ca<sup>2+</sup> concentrations between 0.04 and 1.14 cmol<sub>c</sub> dm<sup>-3</sup>,  
341 Mg<sup>2+</sup> between 0.12 and 0.96 cmol<sub>c</sub> dm<sup>-3</sup>, and K<sup>+</sup> between 0.04 and 0.16 cmol<sub>c</sub> dm<sup>-3</sup>. Jandl [44] reported  
342 that low Ca<sup>2+</sup> levels in forest soil suggest that the species access this nutrient from other sources. Thus,  
343 litter may be one of these sources, because Espig [45] found that Ca was the nutrient with the highest  
344 litter concentration (15.73 g kg<sup>-1</sup>) and with the highest contribution (170.7 kg ha<sup>-1</sup> year<sup>-1</sup>), in a similar area  
345 of this study. This result was confirmed by Godinho [46] in a submontane seasonal semideciduous forest,  
346 an ecosystem associated with Atlantic Forest biome, where Ca was also found with higher concentration  
347 and content in the litter.

348

349 Barreto [47] found that the concentration of the bases Ca<sup>2+</sup>, Mg<sup>2+</sup>, and K<sup>+</sup> in forest areas was high in the  
350 superficial layer and decreased with the depth, favoring the concentration of potential acidity (H+Al).

351

352 Furtini Neto [7] affirmed that excess exchangeable Al in soil can limit the development of species,  
353 inhibiting the growth, and the acquisition and utilization of nutrients by plants. Beutler [48] studied the  
354 effect of exchangeable Al on the initial growth of two forest species, and identified that Al toxicity was  
355 characterized by reduction in plant height and dry matter production, as well as by decrease of N and P  
356 concentration in the aerial part of the two species.

357

358 Despite the high soil acidity of study fragment, exchangeable Al concentrations did not impede the  
359 development of the forest species, because there was no delay in tree growth. This proves the tolerance  
360 and the adaptability of **these** forest species in acidic soil, or the chelating effect of organic matter on the  
361  $Al^{3+}$ . In fact, the total organic C concentration (TOC) of the soil of fragment was high, mainly in  
362 subsurface, and it could have exerted a chelating effect on the exchangeable Al (Table 2).

363  
364 According to Jansen [49], Al seems to be a beneficial element for some individuals of native species and  
365 plants that are adapted to acid soils and high concentration of exchangeable Al do not show symptoms of  
366 toxicity. For Hartwig [50], the exudation of organic acids activated by Al presence at the root apex of  
367 tolerant species is the main mechanism of plants tolerance to  $Al^{3+}$  of the soil. However, due to the  
368 differences between the species, other mechanisms should be investigated.

369  
370 The highest P concentration was observed in the superficial layer, probably due to the decomposition of  
371 leaves and branches [51], decreased with depth. Similar behavior was observed in forest fragments  
372 studied by Espig [42] and Teixeira [43] in Pernambuco, Brazil.

### 373 **4.3 Geospatial variability of soil chemical attributes and forest species distribution**

374  
375 Kerry [52] affirmed that the type of result occurring in *Protium heptaphyllum* may also be associated with  
376 the small size of the data set, not necessarily meaning that they are spatially independent. Additionally,  
377 important spatial variation characteristics may be lost because of the large-scale sampling spacing.

378  
379 The occurrence of spatial dependence of forest species may represent a positive indicator of fragment  
380 conditions because, according to Amaral [19], disturbances in forests can disrupt the spatial dependence  
381 of this and other variables, since it allows the emergence of independent regions, eliminating the spatial  
382 influence of the variables.

383  
384 The spherical model adjusted for the studied species corroborates with the results of many studies that  
385 found the spherical and exponential mathematical models as the most adapted to describe the  
386 semivariograms behavior of attributes of plants and soils [11, 16, 17, 53, 54, 55].

387  
388 The greater attribute dependence degree, better the estimation of the kriging technique for non-sampled  
389 sites [16]. High range values demonstrated the effectiveness of sampling to portray the spatial variation of  
390 soil chemical attributes and forest species (Tables 3 and 4). Artur [55] affirmed that the knowledge of  
391 spatial dependence range allows one to define the sampling radius, guaranteeing the sampling points  
392 independence, the minimization of standard error of the mean, and also the number of samples to be  
393 collected, serving for further surveys in areas with similar characteristics.

### 394 **4.4 Soil-vegetation relationship**

395  
396 The pattern of the spatial distribution of trees in a forest has influences from biotic and abiotic processes,  
397 and these factors may fit into an intrinsic form to the species (reproductive, social) or extrinsic (wind,  
398 luminosity and edaphic conditions) [56].

399  
400 The wide spatial distribution of *Brosimum rubescens* may have occurred because of the dispersion of  
401 their seeds carried by animals, especially mammals. The fruits of this species are fleshy and attractive,  
402 having seeds with high self-regeneration capacity and with formation of abundant seedlings bank [57].  
403 Santo [58] found *Brosimum rubescens* occurred preferentially in areas with low exchangeable K and Mg  
404 concentrations in a study carried out in the Amazon region. These attributes, except SB, also correlated  
405 positively with *Brosimum rubescens* (Table 5).

406  
407 In programs for recovery of degraded areas, *Brosimum rubescens* can be used in soils with different  
408 chemical characteristics by their dispersion capacity, but it can be preferably used in high potential acidity  
409 soils, as long as it presents adequate levels of exchangeable Mg.

410  
411

412  
413 *Tovomita mangle* did not present a specific relation with any chemical attribute, and it can be used in any  
414 environment, provided that it presents low levels of exchangeable K. Unlike *Brosimum rubescens* and  
415 *Tovomita mangle*, *Anaxagorea dolichocarpa* was more demanding, and it can not be indicated for  
416 recovery of degraded areas in soils of low natural fertility, especially when the exchangeable K  
417 concentrations are restrictive.

418  
419 **Spatial** distribution of *Protium arachouchini*, resembling the behavior presented by *Brosimum rubescens*,  
420 corroborate with Santo [58], who found that *Protium arachouchini* occurs only in typologies with better  
421 natural soil fertility, especially with higher organic matter and exchangeable Mg and low Al exchangeable  
422 concentrations. At first, as there was no identification of significant correlations between soil attributes and  
423 spatial distribution of *Protium arachouchini*, it could be said that it would be a species to be used in any  
424 soil chemical condition. However, kriging maps restricted **this** species distribution to specific areas, with  
425 indicators of higher natural fertility (Fig. 3 and 4), which suggests that caution is required in  
426 recommending this species to generic areas, restricting its use for more fertile soils or at least with  
427 moderate chemical restriction.

428  
429 The lowest **density** of *Caraipa densifolia* **was** observed in areas with higher saturation **by** Al. This restricts  
430 the recommendation of this species to non-alic environments, and it should be used only in areas with  
431 high natural fertility. The pattern of spatial distribution of *Caraipa densifolia* was antagonistic to that  
432 presented by *Talisia retusa*. Therefore, in alic environments, where *Caraipa densifolia* can not be  
433 recommended, *Talisia retusa* can be alternatively recommended for its tolerance to high levels of Al<sup>3+</sup>.

434  
435 In this study, *Inga capitata* showed a positive correlation with potential acidity (H+Al) and CEC<sub>potential</sub>.  
436 Thus, it is a species that can be recommended for restrictive environments with low natural fertility.

## 437 438 5. CONCLUSION

439  
440 In conclusion, the results of this study showed that the soils of the studied area presented high acidity,  
441 high saturation by aluminum, and low base saturation, indicating low natural fertility. There was a  
442 correlation between spatial distribution of the species of natural regeneration and soil chemical attributes,  
443 suggesting that the soil contributes effectively to the density of the species. Some recommendations for  
444 species selection may be useful in programs for recovery of degraded areas in forest environments where  
445 soil chemical attributes are known. For example: *Tovomita mangle* may be indicated for soils with  
446 different chemical characteristics and diverse natural fertility; *Brosimum rubescens* and *Inga capitata* for  
447 acid soils with low natural fertility, but with moderate Al exchangeable concentrations; *Talisia retusa*  
448 should be recommended for alic soils, while *Caraipa densifolia* for non-alic soils; and *Anaxagorea*  
449 *dolichocarpa* and *Protium arachouchini* for soils of high natural fertility. Thus, there is no specific soil  
450 chemical attribute that has affected species. For example, where Al levels were high some species did  
451 not settle, but others were more tolerant and regenerated.

452  
453 In general, the attributes that most affected the distribution of the species were: SB, (H+Al), Mg<sup>2+</sup>, K<sup>+</sup>,  
454 CEC<sub>potencial</sub>, V e m.

455  
456 The knowledge of the spatial distribution of native species and soil attributes, through methods that  
457 consider the spatial dependence between samples, can contribute with accuracy in the interpretation of  
458 data behavior in forest fragments. It aims to define strategies for the recovery of areas with similar  
459 characteristics, as well as to predict the variations in the study environment in order to subsidize  
460 management techniques for their conservation.

## 461 462 REFERENCES

463  
464 1. Varjabedian, R. Atlantic Rainforest law: Environmental Regression. Estud. Av. 2010;24:147-160.  
465 <http://dx.doi.org/10.1590/S0103-40142010000100013>. Portuguese.

- 466 2. Mantovani MC, Scachetti AL, Garrido O, Mikami L, Spina L, Herrera A. et al. Atlantic rainforest  
467 remnants atlas. 2009. Accessed 13 July 2016. Available: [https://www.sosma.org.br/projeto/atlas-da-](https://www.sosma.org.br/projeto/atlas-da-mata-atlantica/dados-mais-recentes.html)  
468 [mata-atlantica/dados-mais-recentes.html](https://www.sosma.org.br/projeto/atlas-da-mata-atlantica/dados-mais-recentes.html) Portuguese.
- 469 3. Silva RKS, Feliciano ALP, Marangon LC, Lima RBA. Floristic and ecological succession of the arboreal  
470 vegetation in the spring area of an Atlantic forest fragment, Pernambuco, Brazil. *Rev. Bras. Ciênc.*  
471 *Agrár.* 2010;5:550-559. <http://dx.doi.org/10.5039/agraria.v5i4a829>. Portuguese.
- 472 4. Silva RKS, Feliciano ALP, Marangon LC, Lima RBA, Santos WB. Structure and dispersal syndromes of  
473 tree species in a stretch of riparian vegetation, Sirinhaém, Pernambuco State, Brazil. *Pesq. Flor. Bra.*  
474 2012;32:1-11. <http://dx.doi.org/10.4336/2012.pfb.32.69.01>. Portuguese.
- 475 5. Pessoa MML. Ecological succession in fragments of atlantic rainforest and in *Corymbia citriodora*  
476 (Hook.) K.D. Hill & L.A.S. Johnson, in Sirinhaém, Pernambuco [Dissertation]. Recife: Federal Rural  
477 University of Pernambuco, Brazil; 2012. Portuguese.
- 478 6. Brandão CFLS. Structure of tree component and natural regeneration in Atlantic Rainforest fragments  
479 of different sizes in Sirinhaém, Pernambuco [Thesis]. Recife: Federal Rural University of Pernambuco,  
480 Brazil; 2013. Portuguese.
- 481 7. Furtini Neto AE, Resende AV, Vale FR, Faquin V, Fernandes LA. Soil acidity, growth and mineral  
482 nutrition of some tree species at seedling phase. *Cerne.* 1999;5:1-12. Portuguese.
- 483 8. Silva RBM, Francelino MR, Moura RA, Moura TA, Pereira MG, Oliveira CP. Soil-vegetation relation in  
484 cerrado environment under influence of the group *Urucuia*. *Ci. Fl.* 2015;25:363-373.  
485 <http://dx.doi.org/10.5902/1980509818455>. Portuguese.
- 486 9. Silva Neto SP, Santos AC, Leite RLL, Dim VP, Neves Neto DN, Silva JEC. Spatial variation of the  
487 content of soil organic matter and production of pasture grass marandu. *Biosc. J.* 2012;28:41-53.  
488 Portuguese.
- 489 10. Kanegé Junior H, Mello JM, Scolforo JRS, Oliveira AD. Evaluation of spatial continuity of dendrometric  
490 characteristics of clonal stands of *Eucalyptus* sp. at different ages. *Rev. Árvore.* 2007;31:859-866.  
491 <http://dx.doi.org/10.1590/S0100-67622007000500010>. Portuguese.
- 492 11. Rosa Filho G, Carvalho MP, Montanari R, Silva JM, Siqueira GM, Zambiano EC. Spatial variability of  
493 dendrometric properties of eucalyptus and physical attributes of a *Rhodic Eutrudox*. *Bragantia.*  
494 2011;70:439-446. <http://dx.doi.org/10.1590/S0006-87052011000200027>. Portuguese.
- 495 12. Bognola IA, Ribeiro Junior PJ, Silva EAA, Lingnau C, Higa AR. Univariate and bivariate modeling of  
496 the spatial variability of yield of *Pinus taeda* L. *Ci. Fl.* 2008;38:373-385. Portuguese.
- 497 13. Assis AL, Mello JM, Guedes ICL, Scolforo JRS, Oliveira AD. Development of a sampling strategy for  
498 young stands of *Eucalyptus* sp. using geostatistics. *Cerne.* 2009;15:166-173. Portuguese.
- 499 14. Guedes ICL, Mello JM, Mello CR, Oliveira AD, Silva ST, Scolforo JRS. Geostatistical techniques and  
500 spatial interpolators in the stratification of *Eucalyptus* sp. *Stands. Ci. Fl.* 2012;22:541-550.  
501 <http://dx.doi.org/10.5902/198050986621>. Portuguese.
- 502 15. Wojciechowski JC, Schumacher MV, Kilca RV, Brun EJ, Pires CAF, Silva CRS. Et al. Geostatistics  
503 applied to the study of soil physiochemical characteristics in seasonal deciduous forest areas. *Ci. Fl.*  
504 2009;19:383-391. <http://dx.doi.org/10.5902/19805098894>. Portuguese.
- 505 16. Lima JSS, Souza GS, Silva SA. Sampling and spatial variability of chemical attributes of a soil under  
506 regenerating natural vegetation. *Rev. Árvore.* 2010;34:127-136. [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-67622010000100014)  
507 [67622010000100014](http://dx.doi.org/10.1590/S0100-67622010000100014). Portuguese.
- 508 17. Neves DA, Lemos F, Gonzalez AP, Vieira SR, Siqueira GM. Using geostatistics for assessing  
509 biodiversity of forest reserve areas. *Bragantia* 2010;69:131-140. Portuguese.
- 510 18. Skorupa ALA, Guilherme LRG, Curi N, Silva CPC, Scolforo JRS, Sá JJG. et al. Soil properties under  
511 native vegetation in Minas Gerais, Brazil: distribution by phytophysiology, hydrography and spatial  
512 variability. *R. Bras. Ci. Solo* 2012;36:11-22. <http://dx.doi.org/10.1590/S0100-06832012000100002>.  
513 Portuguese.
- 514 19. Amaral LP, Ferreira RA, Watzlawick LF, Longhi SJ, Sebem E. Influence of forest disturbance in the  
515 spatial distribution of the number of individuals of three species of *Araucaria* Forest assessed by  
516 geostatistical. *Rev. Árvore* 2013;37:491-501. <http://dx.doi.org/10.1590/S0100-67622013000300012>.  
517 Portuguese.



- 518 20.Higuchi P, Silva AC, Van Den Berg E, Pifano DS. Spatial association among individuals of different  
519 species of *Miconia* Ruiz & Pav spp. (Melastomataceae). Rev. Árvore 2011;35:381-389.  
520 <http://dx.doi.org/10.1590/S0100-67622011000300002>. Portuguese.
- 521 21.Martins L, Cavararo R. Technical Manual of the Brazilian vegetation. Phytogeographic system.  
522 Inventory of forest and rural formations. Techniques and management of botanical collections.  
523 Procedures for mappings. Rio de Janeiro: Instituto Brasileiro de Geografia e Estatística; 2012.  
524 Portuguese.
- 525 22.Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate classification  
526 map for Brazil. Meteorol. Z. 2013;22:711-728. <http://dx.doi.org/10.1127/0941-2948/2013/0507>.
- 527 23.Oliveira PRS, Guedes RVS, Silva CAWS, Santos EP, Oliveira FP, Silva-Júnior HD. et al. Climate  
528 Bulletin: Climate Synthesis. Recife: Agência Pernambucana de Águas e Clima; 2016. Portuguese.
- 529 24.Santos HG, Jacomine PKT, Anjos LHC, Oliveira VA, Lumberras JF, Coelho MR. et al. Brazilian  
530 System of Soil Classification. Brasília: Embrapa; 2013. Portuguese.
- 531 25.Donagema GK, Campos DVB, Calderano SB, Teixeira WG, Viana JHM. Manual of Soil Analysis  
532 Methods. Rio de Janeiro: Embrapa Solos; 2011. Portuguese.
- 533 26.Finol UH. Nuevos parámetros a considerarse en el analisis estructural de las selvas virgenes  
534 tropicales. Rev. For. Ve. 1971;14:29-42. Spanish.
- 535 27.Marangon LC. Floristic and phytosociology of area of semideciduous seasonal forest aiming to study  
536 dynamic of tree species in Viçosa, MG. [Thesis]. Federal University of São Carlos, Brazil; 1999.  
537 Portuguese.
- 538 28.APG. An update of the Angiosperm Phylogeny Group classification for the orders and families of  
539 flowering plants: APG III. Bot. J. Linn. Soc. 2009;16:105-121.
- 540 29.Mueller-Dombois D, Ellemberg H. Aims and methods for vegetation ecology. New York: John Wiley &  
541 Sons; 1974.
- 542 30.Fisher RA. Statistical methods, experimental design and scientific inference. New York: Oxford  
543 University Press; 1990.
- 544 31.Vieira SR, Hatfield JL, Nielsen DR, Biggar JW. Geostatistical theory and application to variability of  
545 some agronomical properties. Hilgardia. 1983;51:1-75. <http://dx.doi.org/10.3733/hilg.v51n03p075>
- 546 32.Trangmar BB, Yost RS, Ueharaa G. Application of geostatistics to spatial studies of soil properties.  
547 Adv. Agron. 1985;38:45-94. [https://doi.org/10.1016/S0065-2113\(08\)60673-2](https://doi.org/10.1016/S0065-2113(08)60673-2)
- 548 33.Robertson GP. GS+ geostatistics for the environmental sciences: GS+ user's guide. Michigan:  
549 Gamma Design Software; 1998.
- 550 34.Cambardella CA, Moorman TB, Novak JM, Parkin TB, Karlen DL, Turco RF. et al. Field-scale  
551 variability of soil properties in Central Iowa Soils. Soil Sci. Soc. Am. J. 1994;58:1501-1511.
- 552 35.Guimarães WD, Gripp Junior J, Marques EAG, Santos NT, Fernandes RBA. Spatial variability of the  
553 physical attributes of soil under pasture. Rev. Ciên. Agron. 2016;47:247-255.  
554 <http://dx.doi.org/10.5935/1806-6690.20160029>. Portuguese.
- 555 36.Lopes IS, Feliciano ALP, Marangon LC, Alencar AL. Dynamics of natural regeneration in the  
556 understory of *Pinus caribaea* Morelet. var. *caribaea* in Biological Reserve Saltinho, Tamandaré - PE.  
557 Ci. Fl. 2016;26:95-107. <http://dx.doi.org/10.5902/1980509821094>. Portuguese.
- 558 37.Silva WC, Marangon LC, Ferreira RLC, Feliciano ALP, Costa Junior RF. Natural regeneration's study  
559 of arboreal species in humid forest fragment in city of Catende, south zone of Pernambuco. Ci. Fl.  
560 2007;17:321-331. <http://dx.doi.org/10.5902/198050981964>. Portuguese.
- 561 38.Corrêa, MM. Cutting ants (Hymenoptera, Formicidae) as agents modifying the light availability and  
562 structure of the plant community in northeastern Atlantic forest [Thesis]. Federal University of  
563 Pernambuco, Brazil; 2006. Portuguese.
- 564 39.Lima AS, Feliciano ALP, Marangon LC, Oliveira LSB, Pessoa MML. Natural regeneration of one  
565 Dense Rainforest fragment the Watershed of River Capibaribe, PE-Brazil. Rev. Bras. Ci. Agr.  
566 2013;8:273-278. <http://dx.doi.org/10.5039/agraria.v8i2a2369>. Portuguese.
- 567 40.Alt F, Oelmann Y, Herold N, Schrumpt M, Wilcke W. Phosphorus partitioning in grassland and forest  
568 soils of Germany as related to land-use type, management intensity, and land use-related pH. J. Plant  
569 Nutr. Soil Sci. 2011;174:195-209. <http://dx.doi.org/10.1002/jpln.201000142>

- 570 41. Mafra AL, Guedes SFF, Klauberg Filho O, Santos JCP, Almeida JA, Rosa JD. Organic carbon and soil  
571 chemical attributes in forest areas. Rev. Árvore. 2008;32:217-224. [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-67622008000200004)  
572 67622008000200004. Portuguese.
- 573 42. Espig AS, Freire FJ, Marangon LC, Ferreira RLC, Freire MBGS, Espig DB. Nutrient distribution  
574 between the forest vegetation and the soil on remnants of Atlantic Forest. Rev. Bras. Ci. Agr.  
575 2008;3:132-137. <http://dx.doi.org/10.5039/agraria.v3i2a342>. Portuguese.
- 576 43. Teixeira LJ, Feliciano ALP, Galindo ICL, Martins CM, Alencar AL. Relationships between tree floristic  
577 and soil characteristics in an Atlantic Forest patch, Tamandaré, Pernambuco, Brazil. Rev. Floresta  
578 2010;40:625-634. <http://dx.doi.org/10.5380/rev.v40i3.18924>. Portuguese.
- 579 44. Jandl R, Alewell C, Prietz J. Calcium loss in Central European forest soils. Soil Sci. Soc. Am. J.  
580 2004;68:588-595.
- 581 45. Espig AS, Freire FJ, Marangon LC, Ferreira RLC, Freire MBGS, Espig DB. Litter seasonality,  
582 composition and nutrient input in remnant of Atlantic forest in the state of Pernambuco, Brazil. Rev.  
583 Árvore 2009;33:949-956 <http://dx.doi.org/10.1590/S0100-67622009000500017>. Portuguese.
- 584 46. Godinho TO, Caldeira MVW, Rocha JHT, Caliman JP, Trazzi PA. Quantification of biomass and  
585 nutrients in the accumulated litter in a section of Submontane Seasonal Semideciduous Forest, ES.  
586 Cerne 2014;20:11-20. <http://dx.doi.org/10.1590/S0104-77602014000100002>. Portuguese.
- 587 47. Barreto AC, Lima FHS, Freire MBGS, Quintino RA, Freire FJ. Chemical and physical characteristics in  
588 soil under forest, agroforest system and pasture. Rev. Caatinga 2006;19:415-425. Portuguese.
- 589 48. Beutler AN, Fernandes LA, Faquin V. Aluminum effect on the growth of two forest species. R. Bras. Ci.  
590 Solo. 2001;25:923-928. <http://dx.doi.org/10.1590/S0100-06832001000400015>. Portuguese.
- 591 49. Jansen S, Watanabe T, Smets E. Aluminum accumulation in leaves of 127 species in  
592 Melastomataceae, with comments on the Order Myrtales. Ann. Bot. 2002;90:53-  
593 64. <http://dx.doi.org/10.1093/aob/mcf142>.
- 594 50. Hartwig I, Oliveira AC, Carvalho FIF, Bertan I, Silva JIG, Schmidt D. et al. Associated mechanisms of  
595 aluminum tolerance in plants. Semin: Cien. Agrar. 2007;28:219-228. Portuguese.
- 596 51. Oliveira PC, Carvalho CJR, Sá TA. Trees of ecological services in the Brazilian Amazon. Univ. Sci.  
597 2010;15:265-277. <http://dx.doi.org/10.11144/javeriana.SC15-3.toes>. Portuguese.
- 598 52. Kerry R, Oliver MA. Determining nugget: sill ratios of standardized variograms from aerial photographs  
599 to krige sparse soil data. Precis. Agric. 2008;9:33-56. <http://dx.doi.org/10.1007/s11119-008-9058-0>
- 600 53. Corá JE, Araujo AV, Pereira GT, Beraldo JMG. Assessment of spatial variability of soil attributes as a  
601 basis for the adoption of precision agriculture in sugarcane plantations. R. Bras. Ci. Solo. 2004;  
602 28:1013-1021. <http://dx.doi.org/10.1590/S0100-06832004000600010>. Portuguese.
- 603 54. Chiba MK, Guedes Filho O, Vieira SR. Spatial and temporal variability of weed population in an Oxisol  
604 under no-till system. Acta Sci. Agron. 2010;32:735-742.  
605 <http://dx.doi.org/10.4025/actasciagron.v32i4.5445>. Portuguese.
- 606 55. Artur AG, Oliveira DP, Costa MC, Romero RE, Silva MVC, Ferreira TO. Spatial variability of soil  
607 chemical attributes associated with microrelief. Rev. Bras. Eng. Agríc. Ambient. 2014;18:141-149.  
608 <http://dx.doi.org/10.1590/S1415-43662014000200003>. Portuguese.
- 609 56. Silva KE, Martins SV, Santos NT, Ribeiro CAAS. Spatial patterns of tropical tree species. In: Martins,  
610 SV, editor. Ecology of tropical forests in Brazil. Viçosa: Federal University of Viçosa; 2009.  
611 Portuguese.
- 612 57. Marimon BS, Felfili JM. Seed rain in a monodominant *Brosimum rubescens* Taub. forest and an  
613 adjacent mixed forest in the Araguaia River Valley, Mato Grosso State, Brazil. Acta Bot. Brasílica.  
614 2006;20:423-432. <http://dx.doi.org/10.1590/S0102-33062006000200017>. Portuguese.
- 615 58. Santo NMC. Phytopathology of forest disjunctions (forest islands) in a savannah area of Boa Vista,  
616 Roraima [Dissertation]. Boa Vista: Federal University of Roraima, Brazil; 2010. Portuguese.