

Solubilization of Phosphorus in Phosphate Fertilizers After Treatment With Different Organic Residues

ABSTRACT

Phosphorus is an element of low mobility in soil. It is combined with iron compounds, aluminum and calcium, and organic matter. Phosphorus inorganic compounds found in the soil are conditioned by pH, type and quantity of minerals present in the clay fraction. This study aimed to evaluate the influence of three types of organic waste and the incubation time in the solubilization of three sources of P in the soil of the northeastern semi-arid region. The treatments were arranged in a 3 x 3 x 7 factorial arrangement, and a solo, 3 sources of phosphorus (P), 3 organic waste and 7 incubation times. The treatments were arranged in a randomized block design with three replicates. All soil samples were determined: pH, Ca and P concentrations, 10, 20, 30, 40, 50, 70 and 90 days of incubation. Data were submitted to ANOVA and Tukey test to compare the treatment averages. For the incubation time factor, regression equations were set at the level of 1% probability. The most efficient organic waste in reducing pH and solubilizing phosphate was Leucaena; The biofertilizer was the most efficient source of phosphate in making phosphorus available for soil.

Keywords: fertilizer, availability, soil, biofertilizers.

1. INTRODUCTION

The soils of the Brazilian semi-arid region contain low levels of phosphorus (P), one of the elements that most limit productivity in this region. Consequently, to obtain high yields, the addition of phosphate fertilizers is required. It is known that only a fraction of P applied to the soil of which 15 to 25% is used and the rest remains in the soil. Another part is adsorbed by colloids, another part is combined with the soil components, remaining in the insoluble form and also, another part is immobilized by microorganisms [1].

Practically, there are no detailed studies on the desorption of P in soils of the northeastern semi-arid, perhaps as a consequence of the lack of use of fertilizers, since, P is an element whose use is quite expensive in agriculture and presents great power of adsorption to soil colloids, which highlights a need to use alternative sources, such as fertilizers of organic origin, which present costs lower than conventional fertilizers and greater residual effect.

However, since this source of P has low solubility, one of the alternatives for this is to increase the activity of the microbiota that participates in the solubilization of P in the soil. Thus increasing the production of organic acids and other products of microbial metabolism [2,3], which can be increased with the addition of organic compounds to the soil [4].

Through this process, there is a slow but continuous release of phosphorus from phosphate sources (FF), in contrast to soluble fertilizers that provide high content of soluble phosphorus immediately after its application.

In this way, it was tried to evaluate the influence of three types of organic residues and the incubation time in the solubilization of three P sources in the Northeastern semi-arid soil.

2. MATERIAL AND METHODS

The experiment was conducted in the Laboratory of Physics and Soil Management of the Department of Agroecology, of the Federal Institute of Alagoas, Campus Piranhas-AL, in 2016.

The experiment consisted of a 3x3x7 factorial arrangement (phosphate, residue, time), with three replicates, being the treatments were arranged in a randomized block design.

The following materials were selected: a characteristic Luvisolic soil with greater predominance in the region; three sources of phosphorus: rock biofertilizer, natural phosphate (Fertipar), and simple superphosphate; and three organic residues: one legume (*Leucaena leucocephala*), earthworm humus, and tanned bovine manure.

Vegetable residues were ground in forage. Humus and manure were sieved in a 2-milimeter sieve. These materials were applied to soil samples, collected in the 0-20-centimeter layer. After collection, the soil was air-dried, broken down and sieved in a 2-milimeter mesh and submitted to chemical analysis (Table 1), according to [5]. The pH of water was determined in the ratio 1: 2.5 soil: solution.

Table 1: Chemical characteristics of the soil used in the experiment. Piranhas. IFAL, 2016.

pH	P	Ca ²⁺	Mg ²⁺	Al ³⁺	Na ⁺	K ⁺	H+Al
mg dm ⁻³	cmol _c dm ⁻³						
6.90	11.00	12.50	3.20	0.00	0.07	0.85	1.65

The available phosphorus was determined by colorimetry using ascorbic acid as a reducing agent after extraction with 0.05 mol L⁻¹ HCl solution and 0.025 mol L⁻¹ H₂SO₄ solution (Mehlich). Exchangeable potassium and sodium were determined by flame photometry, after extraction with Mehlich⁻¹, while calcium, magnesium and exchangeable aluminum were be extracted with 1mol L⁻¹ KCl and determined by titrimetry. The potential acidity (Al + H) was evaluated by extraction with 0.5 mol L⁻¹ calcium acetate at pH 7.0.

Table 2: Main characteristics of the organic residues used in the experiment. Piranhas. IFAL, 2016.

Residues organic ¹	Parameters						
	K	Mg	Ca	P ₂ O ₅	C	N	C/N
	g kg ⁻¹						
Leucena	3.60	18.10	38.50	4.50	500.00	26.00	19
Humus	12.80	3.88	5.80	8.70	190.00	15.00	12
Bovine manure	14.28	4.65	9.35	5.75	315.50	21.10	15

The dose of 600 kg ha⁻¹ of P₂O₅ was applied. Organic residues were submitted to nitric-perchloric digestion for subsequent determination of Ca and Mg contents by complexometry, P by colorimetry, and K by flame emission photometry [5]. Total organic carbon was

determined by the method of [6]. N content was determined in 100 mg of sample digested with sulfuric acid in the presence of a mixture of copper sulfate and potassium sulfate, by the Kjeldahl method [5] (Table 2). Fertilizers were characterized according to methodology described in [5] (Table 3).

Table 3. Principal characteristics of the phosphate fertilizers used in the experiment. Piranhas. IFAL, 2016.

Phosphate fertilizers ^{1/}	Parameters						
	P ₂ O ₅ total	P ₂ O ₅	P ₂ O ₅	P ₂ O ₅ CNA+ water	MgO	CaO	SiO ₂
	-----%						
SS	18.7	-	10.3	-	-	12.0	-
RBF	25.6	-	16.6	-	5.0	15.0	-
FNP	27.7	-	19.0	-	-	22.8	12.3

^{1/} SS = Simple Superphosphate; RBF = Rock Biofertilizer ; FNP = Fertipar Natural Phosphate

As an experimental unit, plastic bags with a capacity of 500 ml with 100 g of soil were used. After mixing the phosphates, residues and soils, these will be maintained in the field capacity, using distilled water, and will be incubated for 90 days. Up to 50 days of incubation, samples were taken every 10 days. After 50 days, the samples were taken at intervals of 20 days, completing 90 days of incubation. Immediately after the withdrawal of samples, they were submitted to pH and P analysis according to [5]. Available P was extracted with 0.5 mol L⁻¹ sodium bicarbonate solution, with pH adjusted to 8.5 at 1:10 ratio (soil: solution) [7]. The obtained extract was purified with activated charcoal, filtered and an aliquot of clarified extract was used for determination of P by colorimetry, using the phospho-molybdc complex method according to [8].

Data were submitted to analysis of variance and, afterwards, the Tukey test was applied to compare the means of the treatments. For the incubation time factor, regression equations were adjusted at a 1% probability level.

3. RESULTS AND DISCUSSION

The analysis of variance indicated a difference (P = 0.01) for the pH value and Ca and P contents as a function of the factors studied and for all interactions tested (Table 4).

Table 4. Test F in function of the effects of the following factors: organic residues (R), phosphorus sources (PS) and incubation times (T) for the variables pH, contents of P and Ca. Piranhas. IFAL, 2016.

FV	Parameters	
	pH	P
Organic Residues (R)	46.65**	5.58**
Phosphorus Sources (PS)	53.85**	19.43**
Incubation Time (T)	14.53**	145.54**
R x FPS	8.41**	14.32**
T x R	13.08**	70.64**
T x PS	3.78**	67.62**
R x PS x T	3.02**	148.29**
CV (%)	1.96	3.55

** significant at 1% probability level.

Of the added residues, those of leucine resulted in lower mean pH values relative to the phosphorus sources (PS) used (Table 5). This behavior may be a reflection of increased microbial metabolism with production of acidifying compounds [9]. Leucine residues present higher C/N ratio [10,11] compared to other residues and, therefore, greater resistance to decomposition [12].

Table 5. Mean of incubation times of pH values, P content for the type of residue applied and the source of P used. Piranhas. IFAL, 2016.

Residues	SS	BFR	FNP
pH			
Leucine	7.18 b	6.79 b	7.20 a
Humus	7.56 a	7.25 a	7.21 a
Bovine manure	7.62 a	7.28 a	7.33 a
DMS	0.14		
Standard error	0.04		
CV (%)	2.79		
General average	7.27		
P (mg dm ⁻³)			
Leucine	27.51 a	32.83 a	32.02 a
Humus	26.56 a	23.12 b	26.18 b
Bovine manure	27.15 a	31.30 a	19.56 c
DMS	3.93		
Standard error	1.17		
CV (%)	19.68		
General average	27.36		

Means followed by the same lowercase letter in the line do not differ from each other by the Tukey test at 5% probability level. SS = Simple Superphosphate; RBF = Rock Biofertilizer ; FNP = Fertipar Natural Phosphate. MSD = Minimum significant difference; CV = Coefficient of variation.


This higher resistance to decomposition, besides a higher content of N, resulted in a greater decrease of soil pH because of humus and bovine manure residues. A higher acidity observed in treatments with leucine residue had a higher available content of P only when associated to the biofertilizer (Table 5). However, the availability of P must be related to other parameters, because the higher acidity will not result in higher P, since Al³⁺, Fe³⁺ and Mn³⁺ were more available in acid soil. Having these minerals more positive charges, although there are negative and positive sites, the predominance of positive charges attracts H₂PO₄⁻ easily. Therefore, phosphate ions absorb the oxide surface of Al, Fe and Mn by changing with OH and OH²⁺. With respect to bovine manure, it is possible to observe that it provided, on average, lower release of P, showing differences of the other residues in the phosphates of simple superphosphate and natural phosphate.

The content of P made available by the other phosphates is dependent on the type of residue used. In addition, one of the conditions for the solubilization of phosphates is the existence of acidic soil conditions [13,14,15].

The greater decrease in soil pH and increase in P availability of the phosphates when in the presence of leguminous residues, may be due to their higher ammoniacal N content. Organic waste has a higher concentration of N [16], which produces acidification of the environment after oxidation due to the transformation of ammonium into nitrate. Acidification is one of the necessary conditions for increased solubilization of phosphates [13].

Solubilization of P may also result, of the production of CO₂ and organic acids from the mineralization of C-organic and the production of enzymes and chelating and complexing compounds by the microbiota [17]. These factors may exert direct solubilizing action on phosphates [18]. The reaction of CO₂ with water, in basic medium, releases H⁺ and HCO₃⁻ ions being able to solubilize Ca and Mg carbonates and phosphates [19,20].

As for P, there is a great fluctuation in the levels released, in relation to incubation time. For the three sources used, a slow release of P was initially observed in the first 10 days, occurring, from that point, a great increase in the available P content, reaching the maximum value between 30 and 50 days of incubation (Figure 1). Some researchers, studying the kinetics of P release by phosphate solubilizing microorganisms, observed variations in solubilized P contents giving rise to several peaks of maximum concentration [21,22,23,24]. These changes in P concentration may be a consequence of their precipitation with organic metabolites [21,24] and/or of the formation of compounds with the excreted organic acids, which are subsequently used as energy source or nutrient, which can occur several times in the medium [22]. Another alternative to explain this behavior is the difference between the release and the assimilation of P by microorganisms.



167


168 **Figure 1. Application of biofertilizer phosphorus sources, natural phosphate and**
 169 **simple superphosphate (A); and leucine residues, humus and bovine manure (B) on**
 170 **pH change as a function of incubation time.**
 171

172 The upward trend observed at the beginning of the incubation may reflect a microbial
 173 solubilization of the added phosphate, while the subsequent drop may reflect microbial
 174 immobilization in response to increased microbial metabolism. Similar results were observed
 175 by [4], using as carbon source soybean straw, corn and sugarcane bagasse, and [25]
 176 afterwards, addition of glucose. In addition, the decrease in P concentration in the solution
 177 may be associated with its attachment to the colloidal particles [13].
 178

179 Regression equations were adjusted using dependent variables, pH (Figure 1) and P (Figure
 180 2), and the incubation time as an independent variable. Quadratic equations were fitted for
 181 most treatments. In relation to soil pH, their values remained constant throughout the
 182 incubation time for all treatments.
 183

184 Leucena

185 Humus



Bovine manure




Figure 2. Application of legume residues, humus and bovine manure in the availability of P as a function of incubation time.

4. CONCLUSION

The most efficient organic residue in reducing pH and solubilizing phosphate was leucine.

However, other factors also alter the availability of P.

Natural phosphate and simple superphosphate showed very similar pH at all incubation times. All the materials used showed a certain pH instability during the incubation time. Nevertheless, studies with a greater time of incubation, with the exception of the biofertilizer of rock that stabilized around pH 6.7 at 70 days, are still necessary.

In the first 10 days, it was found that the release of P was slow for the three sources of P. From that point, we observed an increase in the P content available, having reached peaks of availability from 40 to 50 days for legume residues, humus and bovine manure.

209 **COMPETING INTERESTS**

210

211 Authors have declared that there are no competing interests.

212

213

214 **REFERENCES**

215

216 1. Tisdale SL, Nelson WL. Soil fertility and fertilizers. 2th ed. London: Macmillan; 1969.

217

218 2. Gadd G. Fungal production of citric and oxalic acid: importance of metal specification,
219 physiology and biochemical processes. *Adv. Mic. Phys.* 1999;41(1):47-92.

220

221 3. Whitelaw MA. Growth promotion of plants inoculated with phosphate-solubilizing fungi.
222 *Adv. Agron.* 2000;69(1):99-151.

223

224 4. Minhoni MTA, Cardoso EJBN, Eira AF. Effect of five types of organic matter on the
225 microbial solubilization of rock phosphate. *Rev. Braz. of Sci. of Soil.* 1991;15(1):29-35.

226

227 5. EMBRAPA. Brazilian company of agricultural research. Brazilian system of soil
228 classification. Brasília: Embrapa; 2009.

229

230 6. Yeomans JC, Bremner JM. A rapid and precise method for routine determination of
231 organic carbon in soil. *Comm. Soil Sci. Plant.* 1988;19(1):1467-1476.

232

233 7. Raij BV. Selection of laboratory methods to evaluate the availability of phosphorus in soils.
234 *Rev. Braz. of Sci. of Soil,* 1978;2(1):1-9.

235

236 8. Braga JM, Defelipo BV. Spectrophotometric determination of phosphorus in soil and plant
237 extracts. *Rev. Cer.* 1974;113(1):73-85.

238

239 9. Nahas E, Centurion JF, Assis LC. Phosphate solubilising microorganisms and producers
240 of phosphatases from various soils. *Rev. Braz. of Sci. of Soil.* 1994;18(1):43-48.

241

242 10. Fontes RLF, Jucksch I, Mendonça ES, Santos RHS, Dias LE. Environmental impacts of
243 agricultural land use. In: Fontes RLF. Fertility and soil management. Brasília: ABEAS; 1998.

244

245 11. Zucareli C. Seed production of four species of green summer fertilizer and its effect on
246 the fertility of an Eutrophic Purple Latosol. Londrina; Brazilian Congress on Soil
247 Science/Publishing company Midiograf; 2001.

248

249 12. Pelá A, Silva MS, Costa LA, Silva CJ, Zucareli C, Decarli LD, Matter UF. Evaluation of
250 the resistance to the decomposition of ten species of cover crops for no-till. *Rev. Plant. Dir.*
251 1999;53(1):26-33.

252

253 13. Novais RF, Smith TJ. Phosphorus in tropical conditions. Viçosa: UFV; 1999.

254

255 14. Gonçalves GK, Sousa RO, Vahl LC, Bortolon L. Solubilization of the natural phosphates
256 of ducks of mines and arad in two flooded soils. *Rev. Braz. of Sci. of Soil.* 2008;32(5):2157-
257 2164.

258

259 15. Chagas Junior AF, Oliveira LA, Oliveira AN, Willerding AL. Phosphate solubilization
260 capacity and symbiotic efficiency of rhizobia isolated from Amazonian soils. *Acta Sci. Agro.*
261 2010;32(2):359-366.

262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297

16. Freire VP. Study of the C:N ratio in several organic substrates grown with corn, *Zea mays* L. Londrina; Brazilian Congress on Soil science / Publishing company MIDIOGRAF; 2001.

17. Souza DI, Fagotti DL, Saturno DF, Cerezini P, Cervantes VNM, Nogueira MA. Green fertilization associated with basalt and natural phosphate in agroecological systems in southern Paraná and northern Santa Catarina. *Agroec. Not.* 2011;6(2):1-5.

18. Souza CES, Silva MO, Duda GP, Mendes AMS. Phosphorus solubilization of phosphate fertilizers after treatment with different organic residues. *Rev. of Biol. and Ear. Sci.* 2007;7(1):1-7.

19. Skujins J. Semiarid lands and deserts: soil resource and reclamation Logan. Utah: Marcel Dekker; 1991.

20. Meurer EJ. Fundamentals of soil chemistry. Porto Alegre: Genesis; 2006.

21. Babenko YS, Tyrygina G, Grigoryev EF, Dolgikh LM, Borisova TI. Biological activity and physiologo-biochemical properties of bacteria dissolving phosphates. *Mikr.* 1984;53(1):533-539.

22. Illmer P, Schinner F. Solubilization of inorganic phosphates by microorganisms isolated from forest soil. *Soil Biol. and Bioc.* 1992;24(1):389-395.

23. Rodríguez H, Rodríguez M, Goire I. Characterization of Phosphorus solubilizing *Pseudomonas* strains. *Rev Icidca.* 1996;30(1):47-54.

24. Khan JA, Bhatnagar RM. Studies on solubilization of insoluble phosphate rocks by *Aspergillus niger* and *Penicillium* sp. *Fert. Tec.* 1977;14(1):329-333.

25. Minhoni MTA, Eira AF, Bull LT. Microbial biomass, release of CO₂, available phosphorus and pH in soil receiving glucose and rock phosphate. *Rev. Braz. of Sci. of Soil.* 1996;20(2):387-392.