

Proteomic Analysis Reveals Differentially Accumulated Proteins in Banana Somaclonal Variants.

ABSTRACT

Aims. To compare the protein profile of leaves from normal and giant banana plants to identify differentially accumulated proteins specific to each type.

Place and Duration of Study. Central Molecular Biology Lab (LCBM), between April and November of 2017.

Methodology. Protein samples were analyzed in triplicate by two-dimensional gel electrophoresis. The gels were stained with Coomassie Blue G250 and evaluated by the ImageMaster 2D Platinum 5.0 software. Differentially abundant proteins were treated with trypsin enzyme, and aliquots of each hydrolysate sample were analyzed by matrix-assisted laser desorption/ionization tandem time-of-flight mass spectrometer. The molecular mass profile of each sample was subjected to comparative analysis using the MASCOT program and the NCBI database.

Results. Of the 36 differentially abundant proteins that were analyzed, 21 were identified as being related to root metabolism, photosynthesis, protein translation, and carbon and nitrogen fixation. Notably, glutamine synthetase was more abundant in the leaves of the giant banana plants.

Conclusion. This pioneering work used proteomic analysis to identify differentially abundant proteins in the leaves of normal and giant banana plants. Glutamine synthetase was revealed as a potential molecular marker that could be used to screen out plants of undesired variants. These results highlight some of the biochemical alterations that occur in banana trees with gigantism, providing a basis for future research on micropropagation programs and biofactories.

Keywords: banana; somaclonal variation; bidimensional electrophoresis; mass spectrometry, proteomic profile

1. INTRODUCTION

Banana (*Musa* spp.) is one of the world's most important economic and social crops, with cultivation occurring in more than 125 countries. Globally, banana ranks as one of the most popular fruits, with 106.5 million tons produced per year [1, 2]. Brazil cultivates more than 474 thousand hectares of banana plants and produces seven million tons of the fruit, placing it among the five largest producers in the world [3].

Conventional field propagation relies on transplanting banana suckers that have been severed from existing plants. Plant production of banana suckers is seasonally dependent, limiting the amount of available planting material [4]. Conventional propagation also risks spreading pathogenic agents from the field plant donors.

In Brazil, micropropagation has increasingly been used to meet demand. [5]. Micropropagation ensures the production of plants that are free of pests and disease, preventing dissemination of potential pathogens [6]. Nevertheless, *in vitro* propagation can lead to the occurrence of somaclonal variations, caused either by genetic or epigenetic modifications [7]. In Cavendish banana cultivars, for example, the occurrence of off-types in tissue-cultured plantlets can range from 6 to 38 % [4].

In the banana cultivar Prata Anã (AAB), the most frequent somaclonal variation is characterized by plant overgrowth and the production of bunches with low economical value, known as giant plants [6].

Since these morphological characteristics can only be identified in adult variant plants, molecular markers are needed to facilitate early detection.

Guimarães et al. [9] worked with somaclonal variants of gigantism in the Prata Anã cultivar and concluded that markers for simple sequence repeats (SSR) and random amplified polymorphic DNA (RAPD) were unable to detect variants. Chromosome counting was similarly ineffective. In other research, RAPD primers were used to identify a variegated plant in a micropropagated banana cultivar Grand Naine (AAA). However, other variations such as giant plants were grouped as normal plants [10].

Proteins play a central role in biological processes, and proteomic differential analysis can be used to determine which proteins are affected by genetic variation during plant growth and development [11]. The identification of up- and down-regulated proteins related to gigantism could provide useful information on the biochemical processes related to this type of somaclonal variation. These proteins could be used as molecular markers for the early identification of undesired somaclonal variants so that these off-type plants could be excluded from the field.

Recent research has used proteomic profiles to evaluate banana protein expression under various conditions. In the case of the banana cultivar Grand Naine, it was possible to correlate the induction of somatic embryogenesis with increases in a stress-related cationic peroxidase and plant growth hormone-related proteins such as indole-3-pyruvate monooxygenase and adenylate isopentenyl transferase [12].

Many researchers have used the proteomic approach to identify and track changes in proteins involved in growth mechanisms. Two-dimensional gel electrophoresis (2-DE) and mass spectrometry have been used to identify proteins at different ripening and climacteric stages in bananas and other fruits, such as papaya and apple [13, 14]. One study compared banana pulp at pre-climacteric and climacteric stages [15]. Chitinases were the most abundant type of protein in unripe banana while three heat shock proteins and isoflavone reductase were abundant at the climacteric stage [15].

This study identified differently abundant proteins on banana leaves with and without the giant plant somaclonal variation using 2-DE and matrix-assisted laser desorption/ionization time-of-flight/time-of-flight (MALDI-TOF/TOF) tandem mass spectrometry.

2. MATERIALS AND METHODS

2.1 Plant Material

Foliage leaves from both normal and giant plants of *Musa* spp. cultivar Prata Anã (AAB) were provided by a private property located in Lavras, MG, Brazil (Figure 1).



Figure 1. Normal and giant banana plants. The arrow highlights the giant plant.

All plants were 18 months old. The giant plants were selected according to morphological parameters, such as lanceolate and more upright leaves, larger size, and thinner pseudostems.

All samples were obtained from the same clone established through an *in vitro* propagation protocol conducted by BIOCELL Ltda (Sete Lagoas, MG, Brazil).

2.2 Banana leaf protein extraction

Total protein extracts were prepared from the leaves of both normal and giant plants. Triplicate preparations of 200 mg each were ground to a fine powder with a mortar and pestle in combination with liquid nitrogen, following the method described by Gallardo et al. [16]. Approximately 150 mg of powder was used for protein extraction at 4°C in 660 µL of a buffer containing: 7 M urea, 2 M thiourea, 4 % (w / v) CHAPS, 18 mM Tris-HCl, 14 mM Trizma base, a protease inhibitor cocktail (GE Healthcare), 53 units mL⁻¹ DNase I (Sigma), 4.9 Kunitz units mL⁻¹ RNase A (Sigma), and 0.2 % (v / v) Triton X-100. After ten minutes, 14 mM dithiothreitol (DTT) was added. The protein extracts were stirred for 20 minutes and then centrifuged (35,000g for ten minutes) at 4°C. The supernatant was submitted to a second centrifugation as above. The final supernatant, corresponding to the total protein extract, was stored at -20°C.

Protein concentrations were measured according to the Bradford procedure (1976) [17] by applying the GeneQuant unit (GE-Healthcare) and using bovine serum albumin as a standard. The extracted protein samples were loaded onto a sodium dodecyl sulfate (SDS)-polyacrylamide gel and dyed with Coomassie G-250 blue.

2.3 Protein separation by 2-DE

The Multiphor II (GE Healthcare) system was used for 2-DE following the manufacturer's instructions. Protein samples of 250 mg each were loaded onto 18 cm immobilized pH gradient (IPG) strips (linear pH gradient of 3–10; GE Healthcare). The Multiphor II (GE Healthcare) was used under the following conditions: a total of 55,000 Voltage-hours at 15°C; gradient 150 V for one minute; gradient 150 V for three hours; gradient 500 V for three hours; gradient 3500 V for five hours; gradient 3500 V for 12 hours; maximum current setting of 50 µA per strip. Next, for the first-dimension isoelectric focusing step, the IPG strips were equilibrated for 20 minutes on a buffer (100 mM Tris-HCl pH 8.8, 6 M urea, 2 % SDS, 30 % glycerol, and 0.2 mg / mL bromophenol blue) containing 5 mg / mL DTT (to reduce the proteins). Then, the strips were equilibrated for another 20 minutes in a similar buffer that contained 25 mg / mL of iodoacetamide (to alkylate proteins) instead of DTT. The second dimension SDS-polyacrylamide gel electrophoresis (PAGE) was performed at 15°C with 12 % resolving gels (GE Healthcare) using the Multiphor II apparatus (GE Healthcare) under the following conditions: Step-n-hold 600 V for 30 minutes and 600 V for 2 hours (programmed as step and hold) or until the bromophenol blue dye front had run off the gel. The low and high molecular weight protein markers (SDS Calibration Kit) for SDS electrophoresis (GE Healthcare) were used for the molecular mass determinations. At least three replicate gels were run for each sample.

Following electrophoresis, the gels were stained with Coomassie blue G-250 solution for 48 hours and then stored in an ammonium sulfate 20 % (p / v) solution, according to Neuheff, Arold, Taube & Ehrhardt [18].

2.4 Image analysis

The stained two-dimensional gels were digitalized at high resolution using a scanner (ImageScanner, GE-Healthcare) and processed with the Ulmax MagicScan 4.6 software. Image analysis was performed using the ImageMaster 2D Platinum 5.0 (GE-Healthcare) software. To evaluate the differences in protein abundance across different 2-DE gels, the normalized protein spot volume (spot area multiplied by its intensity, as measured by optical density) was applied as a parameter. The relative spot volume (% Vol) represents the ratio of a given spot volume (Vol_s) to the sum of all spot volumes detected in the gel with *n* spots (%Vol = $\frac{Vol}{\sum_{s=1}^n (Vol_s)} \times 100$).

Protein spots with significant and reproducible changes were considered as being differentially expressed proteins. The normalized volumes of the spots from replicate gels were subjected to a student's ANOVA test (*P* = .05) and only statistically significant data, with a difference of at least 1.2 X between treatments was considered.

2.5 In-gel protein digestion

Spots with differential expression were excised, and the gel fragments were washed in 25 mM ammonium bicarbonate and 50% acetonitrile until completely de-stained. After drying, gel fragments were placed in 50 μ L protease solution [20 ng/ μ L sequence grade modified trypsin (Promega Biosciences, CA, USA) in 25 mM ammonium bicarbonate] on ice for 30 min. Excess protease solution was then removed and replaced with 20 μ L 25 mM ammonium bicarbonate. Digestion was performed at 58°C for 30 min. Peptide extraction was performed twice for 15 min with 30 μ L 50% acetonitrile and 5% formic acid. Trypsin digests were then concentrated in a SpeedVac (Savant, USA) to a volume of about 10 μ L and desalted using a Zip-Tip (C18 resin, P10; Millipore Corporation, Bedford, MA, USA).

2.6 Protein identification by MALDI-TOF/ TOF

The spectrometric analyses were performed by matrix-assisted laser desorption-time of flight-tandem mass spectrometry (MALDI-TOF-TOF MS) (Autoflex III Smartbeam, Bruker Daltonics, Germany). The analysis was performed by applying 0.5 μ L of sample and 0.5 μ L of matrix α -cyano-4-hydroxycinnamic acid (CHCA) in a Maldi Target Plate (MTP) *AnchorChip* 800/384.

The spectra generated by mass spectrometry (MS) and tandem mass spectrometry (MS/MS) were obtained in the positive mode. The operation mode for acquisition was linear, with a laser repetition rate of 50 Hz, ion source voltage 1 at 20 kV, ion source voltage 2 at 18.3 kV, ion source lens voltage 6.75 kV and a shot number of 200. The collision-induced dissociation (CID) was used to fragment trypsin peptides in the peptide mass fingerprinting (PMF).


The mass spectrometry results were obtained as the ratio mass/charge for each sample using the software FlexControl 3.0. The analyses of the acquired datasets were performed with the aid of FlexAnalysis 3.0 and Biotools 3.0 software. The apparatus was calibrated by external standards (*Protein Calibration Standard* I and IV - Bruker Daltonics).

The molecular mass profiles of the samples obtained by MS and MS/MS were subject to comparative analysis using the software MASCOT to compare the masses of the observed peptides with a database and consequently identify the most probable proteins in the sample. The database used in the analysis was NCBI (<http://www.ncbi.nlm.nih.gov/>).

3. RESULTS AND DISCUSSION


Plant tissues contain several substances that can interfere with protein extraction [19] and impact proteomic analysis [20], such as phenolic compounds, proteolytic and oxidative enzymes, terpenes, pigments, organic acids, inhibitor ions, and carbohydrates.

In our study, the protein extraction process resulted in 18 mg / ml of proteins. Protein integrity was verified on SDS-PAGE gels that generated protein profiles with well-defined bands (Figure 2), demonstrating that our methods extracted the proteins similarly and efficiently.



202 **Figure 2. Protein profile of leaf proteins detected on SDS-PAGE gels. Protein samples**
 203 **were loaded as follows: 7 µg of protein in well 1; 10 µg in well 2; 15 µg in well 3, and 20**
 204 **µg in well 4.**

205
 206 To analyze the quality of the 2-DE gel results, protein extracts were first separated under
 207 isoelectric focalization at a 3–10 pH range. The 2-DE gel presented a large number of
 208 individualized proteins (spots) without any horizontal or vertical smearing (Figure 3), indicating
 209 the absence or limited number of potential interfering compounds, such as lipids and
 210 carbohydrates.



211
 212 **Figure 3. 2-DE gels for both normal banana leaf (A) and giant plant leaf (B) samples.**
 213

214 Using the ImageMaster 2D Platinum 5.0 software (ANOVA, $P = .05$), 224 spots were detected
 215 on the normal banana samples, and 267 spots were detected on the giant plant samples. Out of
 216 all these spots, 36 represented percent volume differences (% volume = volume occupied by
 217 each spot on the gel), with protein abundance varying from 1.2 X to 7.3 X. In this subgroup, 31
 218 spots were more abundant on the normal banana plant, while five were more abundant on the
 219 giant banana plant.

220
 221 MALDI-TOF/TOF tandem mass spectrometry enabled the identification of 21 proteins from the
 222 36 spots excised from the gel (Table 1).


223 **Table 1** - Significantly changed protein abundance in banana normal and 'giant Plant'. Proteins identified after analysis in MALDI-ToF/ToF, followed by
 224 analysis in the MASCOT program. Mean of protein abundance (% Vol) * in normal banana and "giant plant", with a quantitative increase from 1.2 to 7.3 times
 225 and a significant result in the *t-student test* ($P=0.05$).

226

Spot no.	Accession no.	(% Vol)*		Variation of normalised volume	Protein	Score	Peptide sequence
		Normal	Giant plant				
1556	XP_009393673.1	0,09 $\square$ 0,01	0,158 $\square$ 0,08	1.7X ²	Glutamine synthase	146	IIAEYIWIGGSGGLDLR HETADINTFLWGVANR
1231	NP_084803.1	0,64 $\square$ 0,30	0,21 $\square$ 0,19	3.0X ¹	ATP synthase CF1/alfa	281	ADEISNIIR LIESPAPGIISR ASSVAQVVTTFR IAQIPVSEGYLGR IVNTGTVLQVGDGIAR
1237	NP_084803.1	0,37 $\square$ 0,14	0,17 $\square$ 0,11	2.2X ¹	ATP synthase CF1/alfa	72	EAYPGDVFYLHSR IVNTGTVLQVGDGIAR
1323	NP_084803.1	0,28 $\square$ 0,16	0,17 $\square$ 0,08	1.6X ¹	ATP synthase CF1/alfa	140	IAQIPVSEGYLGR EAYPGDVFYLHSR IVNTGTVLQVGDGIAR
1649	XP_021609330.1	0,38 $\square$ 0,04	0,22 $\square$ 0,08	1.7X ¹	Fructose 1,6 bisphosphate aldolase	102	YAAVSQDNGLVPIVEPEILLDGEHGIDR LASIGLENTEANR
1664	AAV70659.1	0,36 $\square$ 0,05	0,30 $\square$ 0,04	1.2X ¹	Gliceraldehyde-3-phosphate dehydrogenase	181	VPTVDVSVVDLTVR GILGYVEEDLVSTDFVGDSR
1843	gi 22796842	1,27 $\square$ 0,15	0,21 $\square$ 0,08	6.0X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	91	VALEACVKAR DNGLLLHIHR
1847	gi 209417537	1,00 $\square$ 0,07	0,137 $\square$ 0,03	7.3X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase.	101	NHGMHFR IMTLGFVBLLR
1863	gi 4538610	2,82 $\square$ 0,33	0,94 $\square$ 0,12	3.0X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	123	VALEACVKAR DNGLLLHIHR EMTLGFVDLLR WSPELAAACEVWK
2005	gi 131974	0,35 $\square$ 0,09	0,12 $\square$ 0,01	2.9X ¹	Ribulose-1,5- bisphosphate	56	DNGLLLHIHR

					carboxylase/oxygenase		
2086	gi 30421728	0,36 □ 0,04	0,17 □ 0,02	2.1X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	71	VALEACVKAR DNGLLLHIHR
2237	gi 156143103	0,70 □ 0,12	0,16 □ 0,10	4.3X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	90	TFQGPPHGIQVER LIYYTPDYETKNTDILAAFR
2284	gi 17224634	8,72 □ 0,40	3,40 □ 0,30	2.6X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	157	LNYYTPDYEVKDTDILAAFR TFQGPPHGIQVER
2324	gi 304011	4,14 □ 0,50	1,64 □ 0,69	2.5X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	97	VALEACVKAR EMTLGFVDLLR
2327	gi 18026902	0,42 □ 0,21	0,18 □ 0,07	2.3X ¹	Ribulose-1,5- bisphosphate carboxylase/oxygenase	66	EMTLGFVDLLR
2489	gi 209417537	1,12 □ 0,22	2,00 □ 0,57	1.8X ²	Ribulose-1,5- bisphosphate carboxylase/oxygenase	146	EMTLGFVDLLR EIKFEFEPVDKIDK
2589	gi 172246334	1,72 □ 0,96	3,03 □ 0,20	1.8X ²	Ribulose-1,5- bisphosphate carboxylase/oxygenase	130	LNYYTPDYEVKDTDILAAFR
1987	XP_009418696.1	0,54 □ 0,12	0,35 □ 0,05	1.5X ¹	Triose phosphate isomerase	88	VIACIGEKLEER VATPQQAAQEVHLAVR
1800	ABF69972.1	0,38 □ 0,05	0,22 □ 0,08	1,7X ¹	T-protein	178	GGAVDWHIHDER VRDDHIYLVVNAGCR IYFGEFCTLDIHGSHCFLTR
3872	XP_010105219.1	0,21 □ 0,06	0,12 □ 0,02	1.7X ¹	Elongation factor Tu (EF-Tu)	71	QDQVDDEELLQLVELEVR GITINTATVEYETESR

228 These proteins were related to energy, photosynthesis, translation, and carbon and nitrogen
 229 fixation (Figure 4).



230 **Figure 4. The Blast2GO results are summarized into three main GO categories: biological**
 231 **process (BP), cellular component (CC), and molecular function (MF). Only the top ten GO**
 232 **terms for each main function category are shown (green: BP; blue: MF; yellow: CC).**
 233

234
 235 In the samples from the giant banana plants, a large amount of glutamine synthetase (GS)
 236 protein was observed. GS is responsible for catalyzing glutamine formation by condensing
 237 molecules of glutamate and ammonia [21] with the concomitant hydrolysis of adenosine
 238 triphosphate (ATP). GS plays a crucial role in the assimilation and re-assimilation of ammonia
 239 derived from a wide variety of metabolic processes during plant growth and development,
 240 including photorespiration and nitrogen transport [22]. The amide moiety of glutamine is used to
 241 synthesize essential metabolites, such as amino acids, nucleic acids, and amino sugars [23].
 242 Optimal functioning of the metabolic pathways for nitrogen and carbon are critical in plant
 243 growth and biomass accumulation [24].

244
 245 Biochemical studies have identified two GS isoenzymes: cytosolic GS (GS1) and chloroplastic
 246 GS (GS2). GS2 is abundantly expressed in the leaf mesophyll, while GS1 is less frequently
 247 expressed in the leaves and more concentrated in the phloem [25].

248
 249 Previous studies have evaluated GS1 expression in tobacco leaves. Overexpression of GS1 led
 250 to 70 % higher shoots and 100 % greater root dry weight as well as a 50 % increase in leaf area
 251 [26]. Other research presented a large-growth, light-dependent phenotype characterized by
 252 increases of fresh weight, dry weight, and soluble leaf proteins. These findings demonstrate the
 253 importance of this enzyme to the photosynthesis-photorespiration ratio (via ammonia
 254 assimilation), as noted by the lower levels of free ammonia, the increase in the ratio of serine to
 255 glycine (3.5 times), the higher intermediate photorespiratory levels, and the increases in the
 256 levels of glutamine and carbon dioxide [27].

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274


275

Recent research has correlated higher plant biomass and better growth phenotypes of transgenic *Arabidopsis thaliana* [28] and transgenic *Nicotiana tabacum* [29] with the improved efficiency of nitrogen assimilation due to overexpression of the GS gene.

This data suggests that a greater abundance of GS might be related to the giant banana plant's increased growth. Nevertheless, more studies should be conducted to confirm that GS can be used as a molecular marker for the early identification of giant banana plants.

In our study, several proteins were identified as having higher concentrations in the normal banana plants compared to the giant plants. The normal plants had higher protein concentrations of glyceraldehyde-3-phosphate dehydrogenase (GAPDH), fructose 1,6-biphosphate aldolase (ALDO), triose-phosphate isomerase (TPI), synthase ATP sub-unities, ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCo) sub-unities proteins, THF-dependent enzyme (T-protein), and Elongation factor Tu (EF-Tu).

The RuBisCo protein is an important Calvin cycle enzyme with a carboxylase that catalyzes the incorporation of carbon dioxide into organic forms. Other Calvin cycle enzymes were identified as being abundant in the normal banana plants, including GAPDH, ALDO, and TPI (Figure 5).



276

277

278

279

280

281

282

283

284

285

Figure 5. The Calvin cycle. Abundant proteins in normal banana plants are shown in bold letters. Enzymes encoded by non-differentially expressed genes are shown with normal letters. Abbreviations for enzymes: ALDO: aldolase (fructose-bisphosphate aldolase); FBPase: fructose-1,6-bisphosphatase; GAPDH: glyceraldehyde-3-phosphate dehydrogenase; PGK: phospho-glycerate kinase; PRUK: phosphoribulokinase; RPE, ribulosephosphate 3-epimerase; RPI: ribose 5-phosphate isomerase; Rubisco: ribulose-1,5-bisphosphate carboxylase/oxygenase; SBPase: sedoheptulose-1,7-bisphosphatase; TKT: transketolase; TPI: triose-phosphate isomerase.

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

The three latter (GAPDH, ALDO and TPI) proteins also act as glycolytic enzymes in the cytosol [30]. Therefore, these proteins participate both in the metabolic paths to obtain energy through the oxidation of sugars and in the conversion of carbon dioxide to triose-phosphate, which can be converted into glucose, starch, and sucrose. By playing an important role in the synthesis of triose-phosphate, these proteins participate in the creation of compounds that will either be used for the generation of energy or as construction blocks in growing plants.

Like the previously discussed enzymes, the ATP synthase CF1 alpha chain is a key enzyme in plant metabolism and performs an important role in multiple cellular processes [30, 31].

When kept at their ideal levels, these enzymes play a fundamental role in maintaining cellular homeostasis, preserving metabolism, and, consequently, promoting normal growth of the banana tree and its fruit. A decreased amount of these enzymes in giant banana plants might compromise metabolism.

Our results also showed that normal banana plants had higher concentrations of the T-protein. The T-protein and three other enzymes (pyridoxal phosphate-dependent enzyme or P-protein,

NAD⁺ dependent enzyme or L-protein, and lipoic acid-containing protein or H-protein) make up the glycine cleavage multi-enzyme system (GCS). This system is essential for photorespiration and for one-carbon metabolism [32] and directly affects the biosynthesis of nucleic acids, proteins, and lignin [33, 34].

GCS also combines with serine hydroxymethyltransferase (SHM), a mitochondrial enzyme, to convert glycine to serine. The combined GCS/SHM reaction represents the mitochondrial portion of the photorespiration cycle, which occurs in all C3 plant photosynthesis tissues, and converts the 2-phosphoglycolate generated by the RuBisCo enzyme to the 3-phosphoglycerate Calvin cycle metabolite [35, 36].

Other studies have demonstrated that GCS suppression can be lethal to plants [32, 37]. However, if the *Arabidopsis* T-protein is reduced by only five percent, there is no significant alteration to plant growth and photosynthesis, as well as no change in the photorespiratory carbon flux. An increase in *Arabidopsis* T-protein expression neither increased GCS activity nor altered any other metabolic process. Thus, it had no effect on plant growth and photosynthesis [32].

The present data provide no insight on the effect of decreased T-protein expression in giant banana plant leaves. We believe that this finding should be assessed in future studies, since the decreased T-protein levels converge with lower concentrations of RuBisCo, GAPDH, ALDO, and TPI.

In our results, one last protein, EF-Tu, was shown to have higher concentrations in normal plants. EF-Tu proteins belong to either the translational GTPase group or the G-Protein group. EF-Tu proteins are multifunctional and well preserved [38]. They play a fundamental role in the elongation phase of protein synthesis in bacteria and organelles, including mitochondria and plant plastids [39, 40]. EF-Tu genes are positively regulated during plant abiotic stress [41]. In plants under temperature stress, EF-Tu protein precursors protect synthase citrate, malate dehydrogenase, and RuBisCo proteins from thermal aggregation and inactivation, avoiding cellular respiration and photosynthesis restrictions [42, 43].

The current study found larger concentrations of EF-Tu proteins in normal banana plants, indicating their importance in preserving metabolic enzymes and ensuring balance of cell activities, such as protein synthesis and energy generation.

4. CONCLUSION

In conclusion, comparative proteomic analysis has allowed us to identify energy metabolite proteins and stress-responsive proteins from normal and giant banana plants.

Proteins related to photosynthesis, energy production, and translation were identified as differentially expressed in normal plants.

In giant plants, GS was the only abundant protein revealed by our analysis, and it could be a potential molecular marker for early identification of variant plants. Our results also highlight some of the biochemical alterations that occur in banana trees with gigantism, providing a basis for future research.

ACKNOWLEDGEMENTS

We wish to acknowledge financial support from CNPq, CAPES, FAPEMIG and FINEP.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1 FAO (Food and Agriculture Organization of the United Nation). Banana market reviews 2015 - 2016. Rome, 2016. Available: <http://www.fao.org/3/a-i7410e.pdf> Acesso em: 04 janeiro 2018.

2. Singh HP, Uma S, Selvarajan R, Karihaloo JL. Micropropagation for Production of Quality Banana Planting Material in Asia- Pacific. Asia-Pacific Consortium on Agricultural Biotechnology (APCoAB), 2011. New Delhi, India. p.92.

3. IBGE (Brazilian Institute of Geography And Statistics). Levantamento Sistema de Produção Agrícola. Rio de Janeiro, 2016, 29 (12):1-82. ISSN 0103-443X. Portuguese. Available: [ftp://ftp.ibge.gov.br/Producao_Agricola/Levantamento_Sistematico_da_Producao_Agricola_\[mensal\]/Fasciculo/2016/lspa_201612_20170222_133000.pdf](ftp://ftp.ibge.gov.br/Producao_Agricola/Levantamento_Sistematico_da_Producao_Agricola_[mensal]/Fasciculo/2016/lspa_201612_20170222_133000.pdf)

4. Sahijram L, Soneji J, Bollamma K. Analyzing somaclonal variation in micropropagated bananas (*Musa spp.*). In *Vitro Cell Dev Biol Plant*. 2003; 39:551–556.

5. Carvalho ACPP, Rodrigues AAJ, Santos, EO. Produção de Mudanças Micropropagadas de Bananeira. Circular técnica, 2012; 37. ISSN 1679-2254. Portuguese. Available: <https://www.infoteca.cnptia.embrapa.br/infoteca/bitstream/doc/942959/1/CIT12001.pdf>

6. Álvares MC, Caldas LS. Crescimento, produção e variação somaclonal em bananeiras micropropagadas. *Pesquisa Agropecuária Brasileira*. 2002; 37 (3): 415-420. Portuguese.

7. Krishna H, Alizadeh M, Singh D, Chauhan N, Eftekhari M, Sadh RK. Somaclonal variations and their applications in horticultural crops improvement. *Biotech*. 2016; 6:54. DOI 10.1007/s13205-016-0389-7.

8. Stover, R.H. Somaclonal variation in Grande Naine and Saba bananas in the nursery and field. In: PERSLEY, G.J.; DELANGHE, E.A. (Eds.). INTERNATIONAL WORKSHOP ON BANANA AND PLANTAIN BREEDING STRATEGIES, 1986, Cairns. **Proceedings...** Cairns, 1987. p.136-139.

9. Guimarães NCC, Torga PP, Resende EC, Chalfun-Júnior A, Paiva E, Paiva LV. Identificação de variantes somaclonais em bananeiras ‘Prata Anã’, utilizando técnicas moleculares e citogenéticas. *Ciência e Agrotecnologia*. 2009; 33(2): 448-454.

10. Abdellatif KF, Hegazy AE, Aboshama HM, Emara HA, El-Shahed AA. Morphological and molecular characterization of somaclonal variations in tissue culture-derived banana plants. *J Genet Eng Biotechnol*. 2012;10(1):47-53. <https://doi.org/10.1016/j.jgeb.2012.05.002>

11. Canovas FM, Gaudot ED, Recorbet G, Jorin J, Mock HP, Rossignol M. Plant proteome analysis. *Proteomics*. 2004;4:285–298

12. Kumaravel M, Uma S, Backiyarani S, Saraswathi MS, Vaganan MM, Muthusamy M, Sajith KP. Differential proteome analysis during early somatic embryogenesis in *Musa spp.* AAA cv. Grand Naine. *Plant Cell Rep*. 2016. DOI: 10.1007/s00299-016-2067-y

13. Nogueira SB, C. A. Labate CA, Gozzo FC, Pilau EJ, Lajolo FM, Oliveira-do-Nascimento JR. Proteomic analysis of papaya fruit ripening using 2DE-DIGE. *J Proteomics*. 2012; 75:1428-1439. PMID: 22134357. DOI: 10.1016/j.jprot.2011.11.015

14. Zheng Q, Song J, Campbell-Palmer L, Thompson K, Li L, Walker B et al. A proteomic investigation of apple fruit during ripening and in response to ethylene treatment. *J Proteomics* 2013; 93: 276–294. PMID: 23435059 DOI: 10.1016/j.jprot.2013.02.006

15. Toledo TT, Nogueira SB, Cordenunsi BR, Gozzoc FC, Pilau EJ, Lajolo FM, Nascimento JR. O Proteomic analysis of banana fruit reveals proteins that are differentially accumulated during ripening. *Postharvest Biol. Technol.* 2012; 70:51-58. <https://doi.org/10.1016/j.postharvbio.2012.04.005>

16 Gallardo K, Job C, Groot SPPC, Puype M, Demol H, Vandekerckhove J, Job D. Proteomics of Arabidopsis seed germination: A comparative study of wild- type and gibberellin-deficient seeds. *Plant Physiol.* 2002;129:823–837.

17. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principles of protein-dye binding. *Anal Biochem.*1976; 72: 248-254. DOI:10.1016/0003-2697(76)90527-3

18.. Neuhoﬀ V, Arold N, Taube D, Ehrhardt W. Improved staining of proteins in polyacrylamide gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis.* 1988; 9: 255-262. PMID: 2466658 DOI: 10.1002/elps.1150090603

19. Carpentier SC, Witters E, Laukens K, Deckers P, Swennen R, Panis B. Preparation of protein extracts from recalcitrant plant tissues: An evaluation of different methods for two-dimensional gel electrophoresis analysis. *Proteomics.* 2005; 5: 2497–2507. PMID: 15912556 DOI: 10.1002/pmic.200401222

20. Sapcariu SC, Kanashova T, Weindl D, Ghelfi J, Dittmar G, Hiller K. Simultaneous extraction of proteins and metabolites from cells in culture. *MethodsX.* 2014; 1:74–80. PMID: 26150938. DOI: 10.1016/j.mex.2014.07.002

21. Miflin BJ, Habash DZ. The role of glutamine synthetase and glutamate dehydrogenase in nitrogen assimilation and possibilities for improvement in the nitrogen utilization of crops. *J Exp Bot.* 2002; 53:979–987. <https://doi.org/10.1093/jexbot/53.370.979>. PMID: 11912240

22. Lea PJ, Forde BG. The use of mutants and transgenic plants to study amino acid metabolism. *Plant cell envir.* 1994; 179(5):541–556. DOI: 10.1111/j.1365-3040.1994.tb00148.x

23. Unno H, Uchida T, Sugawara H, Kurisu G, Sugiyama T, Yamaya T, Sakakibara H, Hase T, Kusunoki M. Atomic structure of plant glutamine synthetase: a key enzyme for plant productivity. *J Biol Chem.* 2006; 281:29287-29296. DOI: 10.1074/jbc.M601497200

24. Krapp A, Saliba-Colombani V, Daniel-Vedele F. Analysis of C and N metabolisms and of C/N interactions using quantitative genetics. *Photosynthesis Research* 2005. 83: 251–263. DOI: 10.1007/s11120-004-3196-7

25. Kamachi K, Yamaya T, Hayakawa T, Mae T, Ojima K. Vascular bundle-specific localization of cytosolic glutamine synthetase in rice leaves. *Plant Physiol.* 1992; 99:1481–1486. PMID: 16669062; PMCID: PMC1080651

26. Fuentes SI, Allen DJ, Ortiz-Lopez A, Hernández G. Over-expression of cytosolic glutamine synthetase increases photosynthesis and growth at low nitrogen concentrations. *J Exp Bot.* 2001;52(358):1071-81.

27. Oliveira IC, Brears T, Knight TJ, Clark A, Coruzzi GM. Overexpression of Cytosolic Glutamine Synthetase. Relation to Nitrogen, Light, and Photorespiration. *Plant Physiol.* 2002;129(3):1170-80. PMID: 12114571 DOI: <https://doi.org/10.1104/pp.020013>

28. Chenguang Z, Guimin Z, Shilin C, Wei W, Yuanping T, Rentao S. The better growth phenotype of DvGS1-transgenic arabidopsis thaliana is attributed to the improved efficiency of nitrogen assimilation. *Arch Biol Sci.* 2015;67(4):1107-1118. DOI:10.2298/ABS150414079Z

29. Seger M, Gebril S, Tabilona J, Peel A, Sengupta-Gopalan C. Impact of concurrent overexpression of cytosolic glutamine synthetase (GS1) and sucrose phosphate synthase (SPS) on growth and development in transgenic tobacco. *Planta.* 2015;241(1):69-81. PMID: 25213117. DOI: 10.1007/s00425-014-2165-4.

30. Nelson, David L.; COX, Michael M. *Princípios de bioquímica de Lehninger.* Porto Alegre: Artmed, 2011. 6. ed. Porto Alegre: Artmed, 2014. ISBN 978-85-8271-073-9

473 31. Shao H, Cao Q, Tao X, Gu Y, Chang M, Huang C, Zhang Y, Feng H. Cloning and
474 characterization of ATP synthase CF1 α gene from sweet potato. *African J Biotechnol.* 2011;
475 10(82):19035-19042. DOI: 10.5897/AJB11.2238

476 32. Timm S, Jonas Giese J, Nadja Engel N, Wittmi M, Florian A, Fernie AR, Bauwe H.
477 T α protein is present in large excess over the other proteins of the glycine cleavage system in
478 leaves of *Arabidopsis*. *Planta.* 2018;247:41-51. PMID: 28866761. DOI: 10.1007/s00425-017-
479 2767-8

480 33. Hanson AD, Roje S. One-carbon metabolism in higher plants. *Annu Rev Plant Physiol Plant*
481 *Mol Biol.* 2001;52:119-137. PMID:11337394. DOI:10.1146/annurev.arplant.52.1.119

482 34. Li R, Moore M, King J. Investigating the regulation of one- carbon metabolism in *Arabidopsis*
483 *thaliana*. *Plant Cell Physiol.* 2003;44:233-241. PMID: 12668769

484 35. Tolbert NE. The C2 oxidative photosynthetic carbon cycle. *Annu Rev Plant Physiol Plant*
485 *Mol Biol.* 1997;48:1–25. PMID: 15012254.
486 DOI: 10.1146/annurev.arplant.48.1.1

487 36. Douce R, Neuburger M. Biochemical dissection of photorespiration. *Curr Opin Plant Biol.*
488 1999;2:214–222. PMID:10375564. DOI:10.1016/S1369-5266(99)80038-7
489 sd

490 37. Engel N, Daele K, Kolukisaoglu U, Morgenthal K, Weckwerth W, Parnik T, Keerberg O,
491 Bauwe H. Deletion of Glycine Decarboxylase in *Arabidopsis* Is Lethal under
492 Nonphotorespiratory Conditions. *Plant Physiol.* 2007;144:1328-1335. DOI:
493 <https://doi.org/10.1104/pp.107.099317>

494 38. Caldas T, Laalami S, Richarme G. Chaperone properties of bacterial elongation factor EF-G
495 and initiation factor IF2. *J Biol Chem.* 2000;275(2):855-860. PMID: 10625618

496 39. Riis B, Rattan SIS, Clark BFC, Merrick WC. Eukaryotic protein elongation factors. **Trends**
497 *Biochem Sci.* 1990;15:420-424. PMID: 2278101

498 40. Fu J, Momcilovic I, Prasad PV. Roles of Protein Synthesis Elongation Factor EF-Tu in Heat
499 Tolerance in Plants. *J Bot.* 2012, Article ID 835836, 8 pages DOI:10.1155/2012/835836.

500 42. Ristic Z, Momcilovic I, Fu J, Callegari E, Deridder BP. Chloroplast protein synthesis
501 elongation factor, EF-Tu, reduces thermal aggregation of rubisco activase. *J Plant Physiol.*
502 2007;164:1564-1571. PMID: 17766005. DOI: 10.1016/j.jplph.2007.07.008

503 43. Lee GJ, Pokala N, Vierling E. Structure and in vitro molecular chaperone activity of cytosolic
504 small heat shock proteins from pea. *J Biol Chem.* 1995;270(18):10432-10438. PMID: 7737977.