

Effect of substrate treated with tannery sludge on growth and anatomy of conilon coffee cuttings

ABSTRACT

This work evaluated the development and anatomy of the conilon coffee (*Coffea canephora* Pierre - Rubiaceae) seedlings on substrates with varying doses of dehydrated bovine tannery sludge. The experiment was divided into two stages; The first one was carried out in the field in a nursery of seedlings in the city of Colatina and the second was carried out in a Laboratory of Cell and Tissue Biology of Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes. During the field phase, the experiment was performed in randomly selected sections of a propagating nursery area in which each section received five treatments with twelve replicates per treatment. Seventeen replicates of seedlings were included in each treatment for a total of 85 plants per section and 1,020 plants for the whole experiment. Biometric analyzes and gravimetric evaluation of the development of seedlings were carried out at 120 days post - planting at the house of propagation of seedlings. For anatomical and ultrastructural analysis, plant material was processed in accordance with standard techniques for light and electronic microscopy. Despite increasing chromium levels in leaves with increasing doses of sludge, there was no impairment to plant development associated with other components of the tannery sludge treated substrate, such as humus and soil. Structural analysis revealed reduction and disruption of the palisade parenchyma and alteration to the shape and internal structure of chloroplasts. The use of tanning sludge in the proportion of 20% showed positive results, with potential of use in the substrate of *Coffea canephora* seedlings.

Keywords: Sustainability, Tannery sludge, Chromium, Sodium, Residue, Structure, *Coffea canephora*

1. INTRODUCTION

As a country, Brazil harbors one of the largest herds of cattle in the world, with the production of leather being one of the outcomes. In just the first trimester of 2014, 9,164 million units of cattle hides were produced in the country. A great amount of residue is produced on a daily basis as a byproduct of leather processing in tanneries, which has potential agricultural applications. Although there are no federal laws regulating the use of tannery sludge in Brazil, some states have local regulations (e.g., São Paulo) while others, such as Espírito Santo, discard the sludge in leased warehouses at high costs for tanneries. Because the residue is rich in organic matter and essential elements for plants, it has been the subject of numerous investigations into its use and viability, or restriction, for agriculture [1-7].

Tannery sludge may become contaminated with chromium during leather processing, leaving the destination of the residue more problematic. Thus, when used in the substrate of perennial plants, such as conilon coffee, it is diluted and systematically redistributed in the soil in an attempt to prevent major changes to the original basic structure of the soil [8]. Other wastes from human activity, such as sewage sludge, have been considered for the purposes of recovering degraded areas or for the production of biomass for burning. These wastes have desirable characteristics for these purposes, particularly organic matter content and high concentrations of macronutrients needed by plants, such as nitrogen [9,10], similar to the characteristics of tannery sludge.

32 The chromium contained in tannery sludge can bioaccumulate in plant tissues, which may lead to
33 alterations of cellular structures and compromise the processes involved in cellular division, thus
34 leading to reduced biomass production [11]. Han et al. [12] showed that chromium accumulation in
35 leaves can delay the growth and development of this organ and reduce the number of cells of
36 parenchyma that compose leaf mesophyll.

37 Other effects attributed to chromium toxicity are directly related to its participation in the formation of
38 reactive oxygen species (ROS) [13,14]. These molecules act as agents that promote lipid peroxidation
39 and cause severe injury to lipid cellular membranes [15]. Damage to the membrane structure of
40 chloroplasts, in the form of perceptible alterations to chloroplast membranes and disorganization of
41 thylakoids membranes, is evident that when plants are treated with high doses of chromium [16].

42 In addition to chromium, tannery sludge contains high concentrations of sodium, which is used in the
43 cleansing of hides. The resulting high salinity of the residue may cause osmotic disequilibrium in
44 plants when it is used as a component of the substrate, thereby altering plant physiology and
45 anatomy.

46 Waste generated throughout the world should be appropriately used and allocated as close as
47 possible to the site of generation, since the cost of logistics for transporting these wastes often
48 precludes their use in agriculture. In the state of Espírito Santo, some producers of conilon coffee, the
49 main agricultural activity of the state, could benefit from the use of the tannery residues produced
50 locally. Thus, this work aimed to determine whether structural characteristics of conilon coffee
51 seedlings grown on substrate with increasing doses of dehydrated bovine tannery sludge are altered,
52 and if these alterations interfere with plant development.

53 54 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

55
56 The experiment was divided into two stages; the first was carried out in the field in a seedling nursery
57 in the city of Colatina (19°32'22" S, 40°37'50" W; 71 m; Espírito Santo, Brazil), and the second was
58 performed in a Laboratório de Biologia Celular e Tecidual. During the field phase, the experiment was
59 conducted in randomly selected sections of an area of the coffee seedling propagation nursery. Each
60 section received five treatments with twelve repetitions per treatment. Seventeen seedling repetitions
61 were included in each treatment for a total of 85 plants per section and 1,020 plants for the entire
62 experiment.

63 The four treatments consisted of dehydrated tannery sludge, soil and humus, with the concentration
64 of sludge differing among treatments with 10%, 20%, 30% and 40% of the total volume of the
65 substrate (T-10, T-20, T-30 and T-40). The subsoil – humus – chemical fertilizer mixture traditionally
66 used by producers of conilon coffee seedlings in the northern region of Espírito Santo, was used as a
67 control. The genotype used for this work was the clonal conilon coffee variety Vitoria Incaper 8142.
68 This variety has 13 clones; however, for the purposes of the experiment, only clone V8 was tested
69 since it is considered by the producers of the region to be the most vigorous. The composition of the
70 treatments T-C, T-10, T-20, T-30 and T-40 are detailed in Table 1.

71
72
73 **Table 1. Description of the treatments containing the conventional treatment (control T-C), and**
74 **the different proportions of the tannery sludge in the soil (T-10, T-20, T-30 and T-40).**

Treatment	Components of substrate
	136 dm ⁻³ DM soil (even soil used in treatments T-10 and T40) + 625
T-C	g simple superphosphate + 200 g limestone + 200 g KCl and 3 dm ⁻³ DM + Humus.
T-10	10% tannery sludge + 30% humus + 60% soil
T-20	20% tannery sludge + 30% humus + 50% soil
T-30	30% tannery sludge + 30% humus + 40% soil

75

76 The characteristics of the soil used in the treatment substrates are: pH 5; contents of the P; Cr; K; Fe;
77 Cu; Zn; Mn; S and B was respectively of the 5, 32; 48; 7; 0,6; 0,8; 7,9; 112; 0,4 $\mu\text{g.g}^{-1}\text{DW}$; contents of
78 the Ca; Mg; and Na was respectively of the 0,8; 1,3; 0,03 $\text{cmol}_c.\text{kg}^{-1}\text{DM}$; and base saturation of the
79 74.

80 Tannery sludge was provided by a company located in Baixo Guandu (Espírito Santo, Brazil). After
81 processing bovine rawhide, the sludge, released in a concentrated liquid form with 97% humidity (dry
82 base), was placed in evaporation tanks until it reached a humidity of approximately 14 % (dry base).
83 The characteristics of the dehydrated sludge used in this experiment are: pH 12,3; contents of the N;
84 P; K Ca; and Mg was respectively of the 37; 0,09; 0,8; 27 and 1 $\mu\text{g.g}^{-1}\text{DW}$; content of organic matter
85 was 29%; electric conductivity was 17,3 dS.m^{-1} ; contents of the de Cr; Na; Fe; Cu; Zn and Mn was
86 respectively of the 9952; 550; 57; 1; 1; 1 $\mu\text{g.g}^{-1}\text{DW}$.

87 The humus used in the treatment substrates was obtained from the warmer at IFES-Itapina and made
88 with bovine manure. The characteristics of the humus are: pH 6,8; contents of the Cr; remaining
89 Phosphorus; K; and Ca was respectively of the 0,01; 90; 201; 93 $\mu\text{g.g}^{-1}\text{DW}$; contents of the Mg; Fe;
90 Cu; Zn and Mn was respectively of the 146; 57; 1; 1 e 1; and base saturation of 82.

91 Mineral analyses of the soil and the humus used in the treatment substrates were performed by
92 atomic absorption spectrophotometry [17]. Chromium was analyzed according to EPA-SW-846 Test
93 Method 3051A [18].

94 Seedlings were produced from cuttings obtained from adult tissue of orthotropic branches collected
95 from plants with good phytosanitary and nutritional conditions. After removal, the branches were sent
96 to a greenhouse where 30 cm of the extremities of the branches were eliminated (branches
97 containing several cuttings). Standardization of cuttings was then carried out by setting them at a
98 height of 6 cm and leaves with 1/3 of the leaf blades. The cuttings were planted in 11x20 cm
99 polyethylene bags, which had been filled with treatment substrate 30 days prior to the implementation
100 of the experiment. Irrigation by micro sprinklers was done on a daily basis throughout the experiment,
101 always maintaining the field capacity of the substrates.

102 The experimental field stage lasted for 120 days beginning on the day when the cuttings were
103 planted. The number of leaves, seedling height and stem diameter were evaluated and gravimetric
104 analyses were performed with fresh and dry mass of the shoot, root and entire plant (circulation
105 furnace at 70 °C until reaching a constant weight for dry masses). Through obtaining of the dry mass
106 of the root and aerial part, the root dry mass / shoot dry mass ratio was calculated. For weighing
107 shoots (stem and leaves), only leaves that had emerged from the planted cuttings were considered,
108 excluding the original mass of the cutting. Only the new leaves were taken into account when
109 determining the number of leaves. After being dried in the furnace, tissue samples were ground in a
110 Willey knife grinder equipped with two opening sieves with 2 mm mesh. The analysis of leaf chromium
111 was performed by means of wet digestion with $\text{HNO}_3 + \text{HClO}_4$.

112 Quantitative data were subjected to analysis of variance with the F test, and means compared by the
113 Duncan's test ($p < 0.05$). The statistical analyses were performed using the Assistat program version
114 7.7 [19].

115 For anatomical and ultrastructural analyses, fragments of the middle third of the leaf blade were
116 removed and placed in Karnovsky solution [20]. The samples were then washed in sodium cacodylate
117 buffer at 0.05 M and post-fixed in 1% osmium tetroxide in the same buffer for hours at room
118 temperature. The samples were then dehydrated in increasing amounts of acetone, infiltrated and
119 embedded in epoxy resin (Polybed), and polymerized in a furnace at 60 °C. Semi-thin sections of
120 approximately 0.7 μm were then obtained with the aid of an ultra-microtome (Reichert Ultracuts Leica
121 Instruments®) using a diamond knife (Diatome®). The sections were then stained with 1% toluidine
122 blue and 1% borax buffer, and observed under a light microscope (Axioplan, ZEISS), using a
123 Hamamatsu C3077 digitizing camera or a Cannon Power Shot A640 camera and Analysis® software
124 - LINK/ISIS/ZEISS (Oxford, UK). For ultrastructural analyses of the chloroplasts, ultrafine sections
125 were collected in copper grids (300 mesh), which were then contrasted with 5% lead citrate and 1%
126 uranyl acetate, and observed under a transmission electron microscope (TEM 900, ZEISS) at a
127 voltage of 80 Kv.

128 The anatomical and ultrastructural analyses were performed at the Laboratório de Biologia Celular e
129 Tecidual da Universidade Estadual do Norte Fluminense Darcy Ribeiro, Campos dos Goytacazes, Rio
130 de Janeiro, Brazil.

131

3. RESULTS

There were significant differences among treatments for practically all characteristics evaluated, with the only exception being stem diameter and root dry mass/shoot dry mass ratio, which exhibited no statistical differences.

When evaluating the plants height, it was observed that the treatment with the dosage of 20% dehydrated sludge had the highest average plant height, differing from practically all the other treatments. This proportion of sludge/soil/humus was the only treatment that had a positive significant difference ($p < 0.05$); in other words, the increase in height of the coffee seedlings in this treatment was greater than that for the control substrate with conventional chemical fertilizer (Table 2).

Table 2. Mean plant height, number of leaves, stalk diameter and chromium level in leaves of conilon coffee seedlings grown with different concentrations of dehydrated tannery sludge at 120 days

Treatment	Plant height (cm)	Number of leaves	Stalk diameter (cm)	Chromium level in leaves ($\mu\text{g.g}^{-1}\text{DW}$)
T-10	8.75 b	4.45 a	2.82 a	1.65 c
T-20	9.76 a	4.16 ab	2.82 a	3.75 b
T-30	8.29 b	4.14 ab	2.64 a	4.05 b
T-40	8.87 ab	4.07 ab	2.79 a	5.21 a
T-C	8.23 b	3.92 b	2.67 a	1.20 c
Average	8.78	14.15	2.75	3.03

Means followed by different letters in a column differ statistically according to Duncan's test at the 5% level.

The number of leaves per plant at 120 days did not differ among tannery sludge treatments, and only the 10% treatment had a greater number of leaves than conventional substrate (Table 2). There was no significant increase in chromium level in leaves of the 10% sludge treatment (Table 2), with the level being equivalent to that found for plants developed on conventional substrate (without sludge). However, higher proportions of sludge produced increased levels of chromium in leaves, with the 40% treatment having statistically higher levels than the others. Chromium levels in treatments T-10; T-20; T-30 and T40 were 37, 212, 237 and 334%, respectively, which were all higher than the chromium content of leaves grown on conventional substrate (T-C, without sludge).

Practically all characteristics evaluated for the conilon coffee seedlings exhibited higher gain for the 20% sludge proportions of sludge when compared to the treatment with conventional fertilization (T-C, Table 3).

Table 3. Mean fresh mass of shoot (MFPA), root (MF-Root), and plant (MFP); and mean dry mass of shoot (MSPA), root (MS-Root), plant (MSP) and root dry mass/shoot dry mass ratio (RDM/SDM) for conilon coffee seedlings grown on conventional substrate and substrate with different concentrations of dehydrated tannery sludge at 120 days.

Treatment	MFPA	MFR	MFP	MSPA	MSR	MSP	RDM/SDM
-----------	------	-----	-----	------	-----	-----	---------

..... g							
T-10	3.26 ab	1.84 b	5.10 b	0.68 b	0.26 b	0.94 b	0.38 a
T-20	4.08 a	2.79 a	6.87 a	0.90 a	0.48 a	1.38 a	0.53 a
T-30	2.40 b	1.89 b	4.29 b	0.53 b	0.23 b	0.76 b	0.43 a
T-40	3.07 b	1.96 b	5.04 b	0.69 b	0.34 b	1.03 b	0.49 a
T-C	2.62 b	1.85 b	4.47 b	0.56 b	0.29 b	0.84 b	0.52 a
Average	3.09	2.06	5.15	0.67	0.32	0.99	0,47

Means followed by different letters in a column differ statistically according to Duncan's test at the 5% level.

Anatomical examination showed a decrease in palisade parenchyma and a consequent increase in spongy parenchyma, with the changes being most evident in the 30% and 40% sludge treatments. In addition to these changes, disorganization of the treatment (T-20), exceeding not only the treatments with other palisade cells resulting from the loss of cellular volume was observed mainly in the 10%, 30% and 40% sludge treatments (Fig. 1).

The chloroplasts of the conilon coffee plants grown on substrates enriched with tannery sludge exhibited significant structural differences from those of the control plants. The control plants (T-C) possessed chloroplasts with an ellipsoidal shape, internal membranes organized in thylakoids, and stroma with starch grains and plastoglobules (Fig. 2 A and B). In the 10% and 20% tannery sludge treatments there was an increase in the grana and in the stroma, and a decrease in the quantity of starch grains (Fig. 2 C – F). The chloroplasts of plants of the 30% and 40% treatments exhibited internal membranes organized in thylakoids and grana similar to those of the control plants. However, there was also disorganization of the membranes and an absence of starch grains in the stroma of the chloroplasts, with only plastoglobules being present (Fig. 2 H – J). The chloroplasts of plants from all the treatments exhibited changes in shape, compared to those of the control plants (Fig. 2).




Figure 1. Anatomical characteristics of the leaf blade of conilon coffee plants grown in substrates enriched with different quantities of tannery sludge; A, B and C – control plant; D, E and F – plants grown with 10% tannery sludge; G, H and I – plants grown with 20% tannery sludge; J, K and L – plants grown with 30% tannery sludge; and M, N and O – plants grown with 40% tannery sludge. Adaxial epidermis (ed); palisade parenchyma (pp); Spongy parenchyma (pl); abaxial and abaxial epidermis (eb). Bars: A, D, G, J and K = 50µm; B, C, E, F, H, I, K, L, N and O = 20 µm.




Figure 2. Chloroplasts of conilon coffee plants grown on substrates enriched with different quantities of tannery sludge. A and B – control plant; C and D – plants grown with 10% tannery sludge; E and F – plants grown with 20% tannery sludge; G and H – plants grown with 30%

tannery sludge; I and J – plants grown with 40% tannery sludge; granum (G); thylakoid (arrow); starch grains (star); plastoglobules (asterisk). Bars: A, C, E, G and I = 2 µm; B, D, F, H, J = 500 µm.

4. DISCUSSION

Seedling height is one of the main indicators of quality, and has been the subject of several studies involving conilon coffee seedlings. It is also one of the main developmental parameters taken into account by nurseries [21]. In this sense, the average height of plants developed on conventional substrate and substrates containing tannery sludge are within acceptable limits of previous studies. Berilli et al. [2] showed that the height of conilon coffee seedlings grown on a substrate containing soil from the subsoil, tannery sludge in different proportions and without humus reached heights of 5 cm at 120 days. For seedlings on conventional substrate, the heights were approximately 8 cm, similar to those observed in the present study. However, in contrast to what was observed by Berilli et al. [2], the height of the seedlings developed on substrates with tannery sludge with humus and soil exhibited heights that were similar to or higher than that of seedlings developed on conventional substrate. These results reveal that the presence of humus in the substrate, with sludge and soil, produced significant increases in seedling height, most notably in the proportion of 20% sludge, 30% humus and 50% soil.

Mean heights of the conilon coffee seedlings observed in this experiment were below those of coffee seedlings of other studies (21,22). An explanation is that the seedlings of the present study had different ages and were not provided supplemental fertilization, because the purpose of this study was to assess only the influence of the application of tannery sludge as a source of initial organic fertilization, in contrast to the conventional chemical fertilizer used by Berilli et al. [2].

Apparently, the presence of high doses of tannery sludge in the substrate tends to decrease the number of leaves per coffee seedling. This is probably due to the greater amount of factors inhibiting the development of plants, such as the greater presence of chromium and the salinization of the substrate caused by greater doses of sludge. Excessive application of tannery sludge has the potential to be inhibitory for several crops, mainly due to factors such as increased salinity and chromium concentration [1,3,4,23].

The root dry mass/shoot dry mass ratio showed a vegetative growth pattern among the treatments. The shoot dry mass presented higher values in relation to root dry mass, revealing a larger development of the shoot part in comparison to the roots. This pattern of development was also observed by Sales et al. [24] in seedlings of the *Schinus terebinthifolius* Raddi. (Anacardiaceae) treat with tannery sludge. Thus, seedlings with this high ratio may present problems at transplanting moment, because the root system may present difficulties in water supply to the shoot part under conditions of water stress, if the development of the shoot part is much above the development of the roots.

However, even with increased chromium accumulation in leaf tissues of plants with the increased doses of sludge, there was no apparent interference in mass gain or in seedling development, which is similar to the findings of Potasso et al. [1], who assessed the influence of adding tannery sludge to soil on eucalyptus harvest and the assimilation of chromium by this plant (Tables 2 and 3).

The presence of chromium in leaf tissue may alter the natural biochemistry of cells by causing oxidative stress and disturbances in the ultrastructure of the chloroplasts, compromising the photosynthetic process. Thus, the activity of the chromium in plants depends on sensitivity of the particular crop and/or its concentration in the soil or plant tissue [26]. Thus, despite there being an increased presence of chromium in leaf tissue, it seems that conilon coffee experienced no great compromise to plant development.

The significant mass gain for plants in the 20% tannery sludge treatment reveals that the specific mixtures (T-20) can provide a relative balance among the components of the substrate (soil, humus and dehydrated tannery sludge). In this way, the influence of problematic elements, such as sodium or chromium, are minimized by the activity of the humus, while at the same time the nutrients made available by the tannery sludge aid seedling development better than other treatments, being recommended for coffee seedlings, since it did not negatively affect the quality of seedlings.

Despite the fact that the dehydrated tannery sludge consisted of 29% dry mass organic matter and that its composition is rich in essential nutrients for plants, the elevated pH and the presence of sodium and chromium in high concentrations may preclude the direct use of tannery sludge as source

of organic matter or alternative fertilizer [2,8]. However, by associating the sludge with humus in the substrate, there was a positive interaction between the balance of the toxic elements and the availability of the essential nutrients present in the sludge.

Anatomical changes observed in the leaves, such as disorganization of the palisade parenchyma (Figure 1), may be directly related to the quantity of heavy metal, mainly chromium, present in the tannery sludge. Han et al. [12] showed that in the presence of trivalent chromium (Cr^{3+}), white mustard plants (*Sinapis alba* L. - Brassicaceae) exhibit a decrease in both the palisade and the spongy parenchyma, which differs from the results obtained for the present study, in which only a decrease in the palisade parenchyma was observed. Results of studies by Sridha et al. [25] with Zinc (Zn) and Cadmium (Cd), also with mustard plants, were equivalent to those obtained in the present study with respect to the palisade parenchyma, and in which cell disorganization was also evident.

The fact that tannery sludge presents a great amount of chromium may be directly related to the alterations observed in the chloroplasts of the studied plants. Studies with chromium have shown this metal to have phytotoxic effects, mainly by affecting the ultrastructural of chloroplasts and thus altering the organization of the chloroplast membranes and influencing thylakoid membrane arrangements [16,26].

The loss of the standard shape of chloroplasts (ellipsoidal) in plants treated with tannery sludge may also be related to the effect of heavy metals, since radish plants (*Raphanus sativus* L. - Brassicaceae) treated with cadmium also exhibited alterations to chloroplast shape [27]. The same author related these changes to increased stroma volume and the disorganization of thylakoid membranes, characteristics found in the coffee plant chloroplasts analyzed in the present study.

A decrease in the number of starch grains and an increase in the number of plastoglobules, mainly in the 30% and 40% treatments, may be related to stress and chromium toxicity. Increased numbers of plastoglobules in plants grown on substrates treated with sludge can be explained by the fact that these structures act in response to different types of stress [28]. Plastoglobules are lipid reserves that are linked to thylakoid membranes and can act as regions of synthesis and recycling of proteins, and thus have an important role in the response of plants to oxidative stress to the photosynthetic apparatus [29]. The increase in the number of plastoglobules in plants treated with tannery sludge can be directly related to this role of protecting these structures against the oxidative stress caused by the excess sodium and chromium present in the substrate.

5. CONCLUSION

The results indicate alterations to the structural characteristics of leaves of conilon coffee seedlings grown on substrates with varying doses of dehydrated bovine tannery sludge, such as disorganization of the mesophyll and alterations to the chloroplasts, suggesting that these plants were experiencing stress. However, these alterations did not interfere with seedling development, of the coffee seedlings and they present an exhibited optimal performance on substrate enriched with tannery sludge, suggesting that this is an adaptive strategy of these plants.

The treatment that obtained the highest plant dry mass was the one with 20% dehydrated sludge in the substrate, which showed potential for use as a substrate for coffee plants. Such use favors mutually close coffee producers and tanneries, by promoting proper disposal of this residue in the environment and facilitating coffee plant production.

REFERENCES

1. Potasso EL, Scaramuzza WLMP, Weber OLS, Nascentes R, Bressiani AL, Calegario N. Chemical attributes of a cambisol and growth of eucalyptus seedling after the addition of tannery sludge containing chromium. *Journal Tree*. 2014;38: 847-856.
2. Berilli SS, Quiquij JPC, Rembinski J, Salla PHH, Berilli APCG, Louzada JM. Use of sludge tannery substrate as alternative to prepare conilon coffee seedlings. *Coffee Science*. 2014;9: 472-479.
3. Patra APDD. Effect of tannery sludge amended soil on glutathione activity of four aromatic crops: *Tagetes minuta*, *Pelargonium graveolens*, *Ocimum basilicum* and *Mentha picata*. *Ecological Engineering*. 2015;81: 348-352.

4. Chand S, Singh S, Singh VK, Patra DD. Utilization of heavy metal-rich tannery sludge for sweet basil (*Ocimum basilicum* L.) cultivation. *Environmental Science and Pollution Research*. 2015;22: 7470-7475.
5. Sales RA, Ambrozim CS, Vitória YT, Sales RA, Berilli SS. Influência de diferentes fontes de matéria orgânica no substrato de mudas de *Passiflora Morifolia*. *Enciclopédia Biosfera, Centro Científico Conhecer*. 2016;13: 606-6015.
6. Berilli SS, Martineli L, Ferraz TM, Figueiredo FAMMA, Rodrigues WP, Berilli APCG, Sales RA, Freitas SJ. Substrate Stabilization Using Humus with Tannery Sludge in Conilon Coffee Seedlings. *Journal of Experimental Agriculture International*. 2018;21: 1-10.
7. Quarteazani WZ, Sales RA, Pletsch TA, Berilli SS, Nascimento AL, Hell LR, Mantoanelli E, Berilli APCG, Silva RTP, Toso R. Conilon plant growth response to sources of organic matter. *African Journal of Agricultural Research*. 2018;13(4):181-188.
8. Berilli SS, Zooca AAF, Rembinski J, Salla PHH, Almeida JD, Martineli L. Influência do acúmulo de cromo nos índices de compostos secundários em mudas de café conilon. *Coffee Science*. 2016;11: 512-520.
9. Antonkiewicz J, Kołodziej B, Bielińska E. The use of reed canary grass and giant miscanthus in the phytoremediation of municipal sewage sludge. *Environmental Science and Pollution Research*. 2016;23: 9505-9517.
10. Kołodziej B, Antonkiewicz J, Sugier D. Miscanthus × giganteus as a biomass feedstock grown on municipal sewage sludge. *Industrial Crops and Products*. 2016;81: 72-82.
11. Henriques FS. Changes in biomass and photosynthetic parameters of tomato plants exposed to trivalent and hexavalent chromium. *Biologia Plantarum*. 2010;54: 583-586.
12. Han FX, Sridhar BBM, Monts DL, Su Y. Phytoavailability and toxicity of trivalent and hexavalent chromium to Brassica juncea. *New Phytologist*. 2004;162: 489-499.
13. Ali S, Bai P, Zeng F, Cai S, Shamsi IH, Qiu B, Wu F, Zhang G. The ecotoxicological and interactive effects of chromium and aluminum on growth, oxidative damage and antioxidant enzymes on two barley genotypes differing in Al tolerance. *Environmental and Experimental Botany*. 2011;70: 185-191.
14. Kovačik J, Babula P, Klejdus B, Hedbavny J. Chromium uptake and consequences for metabolism and oxidative stress in chamomile plants. *Journal of Agricultural and Food Chemistry*. 2013;61: 7864-7873.
15. Ali S, Zeng F, Cai S, Qiu B, Zhang G. The interaction of salinity and chromium in the influence of barley growth and oxidative stress. *Plant Soil Environment*. 2011;57: 153-159.
16. Choudhury S, Panda SK. Toxic effects, oxidative stress and ultrastructural changes in moss *Taxithelium nepalense* (Schwaegr.) Broth. Under chromium and lead phytotoxicity. *Water, Air, & Soil Pollution*. 2005;167: 73-90.
17. Embrapa. Manual de métodos de análise de solo. Rio de Janeiro, Embrapa 2 Ed. - Centro Nacional de Pesquisa de Solos, 2007;212.
18. Usepa. A Method 3051. 1998. Disponível em: <<http://www.epa.gov/SW-846/3051a.pdf>>. Acesso em: Nov. 2016.
19. Silva FAS, Azevedo CAV. The Assistat Software Version 7.7 and its use in the analysis of experimental data. *African Journal of Agricultural Research*. 2016;11: 3733-3740.
20. Da Cunha M, Gomes VM, Xavier FJ, Attias M, Souza W, Miguens FC. Laticifer system of *Chamaecypariss thymifolia*. A closed host environment for Trypanosomatids. *Biocell*. 2000;24:123-132.

- 370
371
372 21. Filho ACV, Mauri AL, Volpi PS, Fonseca AFA, Ferrão RG, Ferrão MAG, Rodrigues WN, Júnior
373 AS. Growth and quality of clonal plantlets of conilon coffee (*Coffea canephora* Pierre ex A. Froehner)
374 influenced by types of cuttings. American Journal of Plant Sciences. 2014;5: 2148-2153.
375
376 22. Colodetti TV, Rodrigues WN, Martins LD, Tomaz MA. Differential tolerance between genotypes of
377 conilon coffee (*Coffea canephora*) to low availability of nitrogen in the soil. Australian Journal of Crop
378 Science. 2014;8: 1648-1657.
379
380 23. Gupta K. Modulation of antioxidant defense system for detoxification of oxidative stress caused by
381 tannery effluent in *Eichhornia crassipes*. International Journal of Environment. 2014;3: 101-110.
382
383 24. Sales RA, Sales RA, Nascimento TA, Silva TA, Berilli SS, Santos RA. Influence of different
384 sources of organic matter in the spread of schinus terebinthifolius raddi. Scientia Agraria. 2017;18:
385 99-106.
386
387 25. Sridhar MBB, Diehl SV, Han FX, Monts DL, Su Y. Anatomical changes due to uptake and
388 accumulation of Zn and Cd in Indian mustard (*Brassica juncea*). Environmental and Experimental
389 Botany. 2005;54: 131-141.
390
391 26. Panda SK, Choudhury S. Chromium stress in plant. Brazilian Journal of Plant Physiology.
392 2005;17: 95-102.
393
394 27. Vitoria AP, Da Cunha M, Azevedo RA. Ultrastructural changes of radish leaf exposed to cadmium.
395 Environmental and Experimental Botany. 2006;58: 7-52.
396
397 28. Ytterberg AJ, Peltier JB, Wijkk JV. Protein profiling of plastoglobules in chloroplast and
398 chromoplasts. A surprising site for differential accumulation of metabolic enzyme. Plant Physiology.
399 2006;140: 984-997.
400
401 29. Austin JR, Frost E, Vidi PA, Kessler F, Staehelin LA. Plastoglobules are lipoprotein
402 subcompartments of the chloroplast that are permanently coupled to thylakoid membranes and
403 contains biosynthetic enzymes. The Plant Cell. 2006;18: 1693-1703.