

Nursery water supply and biomass accumulation relationships in super sweet maize (*sh2*)

A. Riverti¹, E. Giardina¹ and A. Di Benedetto^{1,2}

¹Faculty of Agronomy, University of Buenos Aires, Avenue San Martín 4453 (C1417DSE), Buenos Aires, Argentina.

²Faculty of Agricultural Sciences, National University of Mar del Plata, Route 226, km. 73.5 (B7620ZAA), Balcarce, Province of Buenos Aires, Argentina.

Authors' contributions

This work was carried out in collaboration between all authors.
All authors read and approved the final manuscript.

E-mail address: dibenede@agro.uba.ar (A. Di Benedetto)

ABSTRACT

An agricultural intensification with maize transplanted crops for increasing food production is needed. However, super sweet maize water management during nursery has been designed from old genotypes in the 90's decade, which indicate that a water stress during nursery did not change plant growth. The aim of this work was to determine the effects of the watering regime during nursery on biomass accumulation for three different super sweet *sh2* maize hybrids to test the hypothesis that a different nursery water supply driven pre- and post-transplant super sweet maize biomass accumulation. Our results from seedlings watered with 100%, 75% or 50% daily evaporation are not in agreement with this previous report and showed that a water stress during nursery would be considered as a limiting source to decrease maize early biomass accumulation. A very early water stress during nursery decrease early post-transplant biomass accumulation, leaf area expansion and change photo assimilate partitioning. These results would force to a change in the commonly accepted water management when a transplant routine from plug trays was used for improving agricultural intensification.

Keywords: growth, transplant, water management

1. Introduction

Sweet corn (*Zea mays saccharata*), considered a vegetable, is a special type of corn destined exclusively for human consumption, in fresh form or in processed foods. As the demands for food increase in coming decades, society will be pressed to increase agricultural production by increasing yields on already cultivated lands, by cultivating currently natural areas or to change current crop consumption patterns. For the future, a strong increase in food demand is expected, which may be fulfilled by further agricultural intensification rather than expansion of agricultural area [1,2,3], especially because climate change can affect both crop yield and the land area suitable for agriculture [4].

Because transplanting provides optimal environmental conditions for seed germination, sweet corn has been transplanted experimentally in an attempt to improve stands and hasten maturity [5,6]. The growing period was significantly shortened with transplantation of sweet corn plants compared to direct seeded [7], although growth and development at early growth stages could become of critical importance for final yield [8]. On the other hand, Orgaz et al. [9] indicate that canopy development and management of some greenhouse horticultural crops is quite different from that of outdoors.

Drought stress is a major cause of yield instability in crops across diverse eco-geographic regions worldwide. In the scenario of global warming, drought is indeed considered to be one of the most important abiotic stresses that adversely affect food security [10]. Limited water supply to roots and/or higher transpiration rate due to elevated atmospheric temperature induces drought stress [11], which severely impairs growth and yield [12].

Maize has been reported in the literature as having high irrigation requirements; a frequent and uniform supply of water is extremely important for maize yield to meet the water requirements of plants [13]. In this way, Oktem et al. [14,15] showed that water applied as 100, 90, 80 and 70% of evaporation, decreased yield. Their results indicate that a 2-day irrigation frequency, with 100% evapotranspiration water application by a drip system, would be optimal for sweet corn grown in semi-arid regions. Seasonal dependence of biomass production on transpiration and linear relationships between relative yield and relative evapotranspiration has been previously found as well [16].

When plants do not receive sufficient water, plant growth is negatively affected [17]. Previous modeling studies indicated that changes in root architecture and water acquisition capacity directly affect biomass accumulation and yield increases [18]. The spatial configuration of root systems is an important factor determining belowground resource acquisition. Campos et al. [19] attributed the differences in water uptake between old and modern maize hybrids to differences in root depth and root intensity. Under water-limited situations, they observed that the old hybrid took up more water from shallow soil depth, whereas the new hybrid appeared to be more effective in acquiring water from the deeper soil layers. Plants growing in soil with less water, can survive the condition either by escape or by resistance. The resistance can be by two means, drought avoidance (maintenance of tissue water potential) and drought tolerance [20]. Recently, Rattin et al. [21] showed a close coordination between roots and shoot growth.

Moisture management, or watering, accounts for the greatest loss in plug vegetable production. Watering at the wrong time or in the wrong way adversely affects plug crops. Because water is essential in plants for many processes, plugs must never be allowed to dry completely. The smaller the cell size (or higher plant relative growth rate), the more vulnerable the plug is to fluctuations in moisture. According to Styer and Koransky's suggestions [22] for the best both root and shoot growth, in Stage 1 (sowing to radicle emergence), most crops require uniformly high moisture levels. Once radicle emerges occurs, the moisture level would be reduced for many crops (Stage 2); by allowing the soil surface to dry out slightly, the roots will tend to go down into the media. Once crops enter Stage 3 (full leaves expansion), leaves and roots are actively growing. The alternating wet/dry pattern increases in depth with advancing growth stages. However, Water et al. [23] using olds shrunken-2 (*sh2*) and sugary (*su*) sweet corn hybrids mutants showed no impact on plant growth and development related to seed moisture during nursery.

The aim of this work was to determine the effects of the watering regime during nursery on biomass accumulation for three different super sweet *sh2* maize hybrids to test the hypothesis that a different nursery water supply driven pre- and post-transplant super sweet maize biomass accumulation.

2. MATERIALS AND METHODS

2.1. Plant Material and Experiments

Two experiments were performed:

2.1.1. Experiment 1: Solute leakage

Three maize super sweet hybrids (*sh2*) ('Butter', '1441' and '3475') with yields ranging between 15 to 17 ton ha⁻¹ were provided by Semillera Basso (Argentina). For the experiment, four replications of 100 seeds from each maize hybrid were immersed in 1.000 cm³-distilled water for 96 hours and water volume was kept during all the time.

2.1.2. Experiment 2: Plug and pot responses

A greenhouse experiment was carried out at the Faculty of Agronomy, University of Buenos Aires, Argentina (34° 35' 59"S, 58° 22' 23"W) between October and November 2017.

Three super sweet maize hybrids (*sh2*) ('Butter', '1441' and '3475') seeds were sown in plastic plug trays (128 cells tray⁻¹; 17.37 cm³ cell⁻¹) using a commercial growing media (Klasmann 411® medium, Klasmann-Deilmann, GmbH, Germany). Daily water irrigation makes up for 100%, 75% or 50% evapotranspiration. Seed emergence (%) was recorded five days from sowing.

At the transplant stage (15 days from sowing), seedlings were kept in 5-litre pots filled with a *Sphagnum maguellanicum*-river waste-perlite (40-40-20, v/v/v) medium [24]. The plants were arranged at a density of 4 plants m⁻², which avoided mutual shading. A weekly ferti-irrigation of 1.0: 0.5: 1.0: 0.5 (v/v/v/v) N: P: K: Ca (nitric acid, phosphorus acid, potassium nitrate, and calcium nitrate) (150 mg l⁻¹ N) was included through to the overhead irrigation water.

Half hourly averages of the air temperature were measured using a HOBO H08-001-02 data logger (Onset Computer Corporation, MA, USA) protected from direct radiation by aluminum foil shades. Minimum temperature, maximum temperature and mean global solar radiation during the experiment were 25.11°C, 28.31°C and 25.60 MJ m⁻² day⁻¹ respectively.

2.3. Growth Evaluations

For experiment 1, solute leakage was estimated through electrical conductivity (dS m⁻¹) change during first 96 hours.

Seedling emergence rates (SER) is calculating by dividing the number of normal seedlings per each 128-plug cell try⁻¹ obtained by each counting in the seedling emergence by the number of days until true leaf expansion. The values obtained at each count are then summed at the end of emergence period to obtain SER according to the method adapted from Maguire [25].

$$SER = \left(\frac{\text{number of normal seedlings}}{\text{days to first count}} + \dots + \frac{\text{number of normal seedlings}}{\text{days to final counts}} \right)$$

For experiment 2, plants for destructive measurements were harvested (five per treatment and block) at emergence and at 7-day intervals during the experiment. Roots were washed and root, stem, leaf and petioles fresh weights (FW) were recorded. Dry weights (DW) were recorded after drying roots, stems, leaves and petioles to constant weight at 80°C for 96 hours. The number of leaves was recorded and each leaf area was determined using a leaf area meter LI-COR FL16 (LI-COR Inc., Lincoln, NE, USA)

The rate of leaf area expansion (RLAE) was calculated as the slope of the regression of the ln of total leaf area versus time (in days). The rate of leaf appearance (RLA) was calculated as the slope of the number of fully expanded leaves versus time (in weeks). The relative growth rate (RGR) was calculated as the slope of the regression of the natural logarithm of the whole plant on a DW basis versus time (in days).

The mean net assimilation rate (NAR), and the leaf area ratio (LAR) were calculated as follows:

149

$$NAR = \frac{k_w W_0 e^{k_w t}}{A_0 e^{k_a t}}$$

150

$$LAR = k_a / \frac{A_a e^{k_a t}}{k_w W_0 e^{k_w t}}$$

151

152 where: k_w : RGR (days^{-1}); W_0 : extrapolated value of total dry weight at time zero (g); A_0 : extrapolated
153 value of leaf area at time zero (cm^2); k_a : RLAE (days^{-1}); t : time (in days) at the midpoint of the
154 experimental period and e : base of natural logarithms.

155

156 The specific leaf area on a FW basis (SLA) was calculated as the ratio between the area of the new
157 individual leaf and leaf FW.

158

159 The allometric coefficients between root and shoot fraction were calculated as the slope (β) of the
160 straight-line regression of the natural logarithm of the root DW versus the natural logarithm of the
161 shoot DW. On the other hand, the β allometric coefficients between leaf blades and the petiole + stem
162 were calculated as the natural logarithm of the leaf blade DW versus the natural logarithm of the
163 petiole + stem DW.

164

165

166 2.4. Statistical Analysis

167

168 We used a complete aleatory design for both experiments. Data were subjected to one-way analysis
169 of variance for experiment 1 and to two-way analysis of variance for experiment 2. Means were
170 separated by Tukey's test ($P < 0.05$) and STATISTICA 8 (StatSoft) software was used. Slopes from
171 straight-line regressions of RLA, RLAE, RGR, NAR, LAR and allometric values were tested using the
172 SMATR package [26].

173

174

175 3. Results

176

177 Solute leakage (experiment 1), estimated through the change in electrical conductivity increase during
178 the first 96 hours of imbibition in distilled water with significant differences between the three sweet
179 maize hybrids tested (Fig. 1).

180

181 Seedling emergence (%) was higher in plants fully watered with no significant differences between
182 75% and 50% evapotranspiration rate for the three maize hybrids tested (Fig. 2). For 'Batter' and
183 '1441' hybrids, the seedling emergence rate (SER) (seeds day^{-1}) was significantly higher between
184 100% water supply and the rest of the watering treatments. However, '3475' maize hybrid showed no
185 SER differences for the two higher water supply (Figure 2). Both seedling emergence and SER
186 differences between maize hybrids tested were found as well.




Fig. 1. Changes in solute leakage during the first 96 hours of the imbibition period in three maize F₁ hybrids ('Batter', '1441' and '3475') seeds.





Fig. 2. Seedling emergence (%) for three super sweet maize mutants under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4). Lower case letters indicate statistically significant differences ($P < .05$) among water supply level for each sweet maize hybrid tested. SER (seeds day⁻¹) is indicated as well.


At the transplant stage, the higher water supply the higher fresh weight for 'Batter' and '3475' maize mutant hybrids. The accumulated '1441' mutant fresh weight showed no significant differences between plants watered with 75% and 50% of daily evapotranspiration (Fig. 3).

A decrease in root: shoot ratio on a dry weight base from transplant and four weeks later with no significant differences between water supply regimes or sweet maize hybrid tested were found. Anyway, at the transplant stage, seedlings watered with 100% evapotranspiration gave the higher root:shoot ratios (Fig. 4).




206

207 **Fig. 3. Changes in fresh weight at the transplant stage for three super sweet maize mutants**
 208 **grown under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4).**
 209 **Lower case letters indicate statistically significant differences ($P < .05$) among water supply**
 210 **level for each sweet maize hybrid tested.**



211



212

213 **Fig. 4. Changes in root: shoot dry weight ratio during the experiment for three super sweet**
 214 **maize mutants (A: Butter; B: 1441 and C: 3475) grown under three water supply levels (100%,**
 215 **75% and 50% evapotranspiration rate) (n = 4).**

216

217 At the transplant stage, there are no significant differences in shoot dry weight in plants watered to
 218 compensate evapotranspiration in different degree but the higher water supply the higher root dry
 219 weight (Fig. 5).




Fig. 5. Changes in both shoots and roots dry weight ratio at the transplant stage for three super sweet maize mutants (A: Butter; B: 1441 and C: 3475) grown under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4).

A linear increase between LMA during the experiment was found for the three maize hybrid mutants tested with no significant differences neither between them nor straight-line regression scopes of the watering routines (Fig. 6).

There is no significant differences in total leaf area at the end of experiment 2; this is when plants 100% watered have expanded near five leaves (V_5 stage). However, a decrease in water supply to compensate 75% daily evapotranspiration give a decrease in '1441' total leaf area but no changes in both 'Butter' and '3475' maize hybrid mutants. A strong decrease in water supply of 50% daily evapotranspiration significantly decreased total leaf area in 'Butter' and '1441' but no in '3475' maize hybrids (Fig. 7).




Fig. 6. Changes in leaf dry mass per unit leaf area (LMA) during the experiment for three super sweet maize mutants (A: Butter; B: 1441 and C: 3475) grown under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4).




Fig. 7. Changes in leaf area during the experiment for three super sweet maize mutants (A: Butter; B: 1441 and C: 3475) grown under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4). Vertical lines indicate least significant differences (LSD).

A decrease in water supply during nursery decrease RLA and RLAE and increase SLA in 'Butter' and '1441' maize mutant hybrids. However, there is no significant changes these growth parameters values in '3475' hybrid (Table 1).

Table 1: Changes in the rate of leaf appearance (RLA), the relative leaf area expansion rate (RLAE) and the specific leaf area (SLA) for three super sweet maize grown under three water supply levels (100%, 75% and 50% evapotranspiration rate, ET) (n = 4). Lower case letters indicate statistically significant differences ($P < .05$) among water supply level for each sweet maize hybrid tested.

Water supply (% ET)	RLA (leaves week ⁻¹)	RLAE (cm ² cm ⁻² day ⁻¹)	SLA (cm ² g ⁻¹)
'Butter'			
100	0.126 ^a	0.140 ^a	35.66 ^b
75	0.097 ^b	0.129 ^b	36.65 ^b
50	0.094 ^b	0.124 ^b	44.55 ^a
'1441'			
100	0.120 ^a	0.146 ^a	37.18 ^b
75	0.097 ^b	0.125 ^b	35.15 ^b

	50	0.094 ^b	0.129 ^b	41.41 ^a
'3475'				
	100	0.099 ^a	0.143 ^a	35.29 ^a
	75	0.097 ^a	0.147 ^a	34.97 ^a
	50	0.094 ^a	0.147 ^a	35.01 ^a

Both 'Butter' and '1441' maize hybrids decreased RGR and NAR when water supply decreased from 100% daily evapotranspiration, without changes in '3475'. These results would be explained by increases in NAR and minor changes in LAR (Table 2).

Table 2: Changes in the relative growth rate (RGR), the net assimilation rate (NAR) and the leaf area ratio (LAR) for three super sweet maize grown under three water supply levels (100%, 75% and 50% evapotranspiration rate, ET) (n = 4). Lower case letters indicate statistically significant differences ($P < .05$) among water supply level for each sweet maize hybrid tested.

Water supply (% ET)	RGR (g g ⁻¹ day ⁻¹)	NAR (g cm ⁻² day ⁻¹) (x 10 ⁻⁵)	LAR (cm ² g ⁻¹)
'Butter'			
100	0.142 ^a	13.29 ^a	106.57 ^b
75	0.134 ^b	11.50 ^b	116.44 ^a
50	0.111 ^b	10.69 ^b	104.08 ^b
'1441'			
100	0.141 ^a	13.17 ^a	106.72 ^b
75	0.128 ^b	11.32 ^b	113.79 ^a
50	0.126 ^b	11.48 ^b	109.52 ^b
'3475'			
100	0.133 ^a	12.52 ^a	107.33 ^a
75	0.135 ^a	11.56 ^b	115.29 ^a
50	0.137 ^a	11.89 ^b	115.22 ^a

When plotting the data from all treatments, we found a close direct relationship ($r^2 = 0.672$) between RGR and NAR and a weak direct relationship between RGR and LAR ($r^2 = 0.479$) (Fig. 8). Nevertheless, plants watered to compensate 100% evapotranspiration showed higher both NAR and LAR values.





Fig. 8. The net assimilation rate (NAR) (empty symbols) and the leaf area ratio (LAR) (full symbols) related to the relative growth rate (RGR). The straight-line regressions were $\text{NAR} = 63.20 \text{ RGR} + 3.46$ ($r^2 = 0.672$ $P < .001$) and $\text{LAR} = 270.71 \text{ RGR} + 74.28$ ($r^2 = 0.479$ $P < .01$). ($n = 4$). Water supply: 100%: Δ , 75%: $\circ$ and 50%: $\square$.

The lower water supply the higher root: shoot and stem: leaves allometries for all maize mutant hybrids tested (Table 3).

Table 3: Changes in allometric relationships between roots and shoots and between stem and leaves for three super sweet maize grown under three water supply levels (100%, 75% and 50% evapotranspiration rate, ET) ($n = 4$). Lower case letters indicate statistically significant differences ($P < .05$) among water supply level for each sweet maize hybrid tested.

Water supply (% ET)	Root: shoot	Stems: leaves
	β	β
'Butter'		
100	0.529 ^b	0.875 ^b
75	0.587 ^a	0.889 ^a
50	0.591 ^a	0.885 ^a
'1441'		
100	0.566 ^c	0.841 ^c
75	0.594 ^b	0.867 ^b
50	0.648 ^a	0.957 ^a
'3475'		
100	0.521 ^c	0.918 ^c
75	0.582 ^b	0.971 ^b
50	0.653 ^a	1.004 ^a

Positive relationships between RLAE ($r^2 = 0.672$ $P < 0.001$) (Fig. 9A), RGR ($r^2 = 0.611$ $P < .001$) (Fig. 9B) and NAR ($r^2 = 0.608$ $P < .001$) (Fig. 9C) and root DW were found. The higher values were those from plants grown with the higher water supply during nursery.






Fig. 9. Relationship between RLAE (A), RGR (B), NAR (C) and root dry weight for three super sweet maize grown under three water supply levels (100%, 75% and 50% evapotranspiration rate) (n = 4). The straight-line regressions were RLAE = 0.0084 root DW + 0.119 ($r^2 = 0.672$ $P < .001$), RGR = 0.011 root DW + 0.104 ($r^2 = 0.611$ $P < .001$), NAR = 0.92 root DW + 9.40 ($r^2 = 0.608$ $P < .001$) (n = 4). Water supply: 100%: Δ , 75%: $\circ$ and 50%: $\square$.

4. Discussion

Low field emergence and early field vigor are major problems in *sh2* maize. The reduced field emergence is due to lower levels of storage carbohydrate and pathogenic infection that thrive on electrolytes leaching out through fractures in the pericarp [27,28]. Sweet maize hybrids with the highest levels of percent emergence and marketable yield tended to have lower levels of soluble solids [29]. Seed vigor is defined as those properties that determine the potential for rapid, uniform emergence, and the development of normal seedlings under a wide range of field conditions [30]. The anion leakage of isolated embryos correlated closely to seed vigor, but sugar leakage was not closely related to total soluble sugar content of seeds. Electrical conductivity measurements of seed leachates are routinely used to determine seed vigor [31]. Results from Fig. 1 showed significant solute leakage differences between the three sweet maize hybrids tested, which are in agreement with results from Cardoso et al. [32] who showed that there is genetic variability in traits such as pericarp thickness. On the other hand, Fig. 2 did not show both low seedling emergence and a clear leachate losses and seedling emergence relationships.

Although a major goal of *sh2* maize-breeding programs is an increase in field emergence [33], transplant routine has shown an alternative to reach better field emergence [34]. When a dry seed is water imbibed, the uptake of water occurs in three stages. Stage I give a rapid initial water uptake due to the seed's low water potential. During this phase, proteins are synthesized using existing mRNA and DNA and mitochondria are repaired. In stage II, there is a slow increase in seed water content, but physiological activities associated with germination are initiated, including synthesis of proteins by translation of new mRNAs and synthesis of new mitochondria. There is a rapid uptake of water in stage III where the process of germination is completed culminating in radicle emergence [35]. In our experiments, the effect of a decrease in water supply on seed vigor status of three super sweet corn hybrids was examined by standard seedling emergence percentage and seedling emergence rate. The lower water supply during nursery, the lower both seedling emergence percentage and SER; differences between sweet maize hybrids were found as well (Fig. 2).

Working with old *sh2* and *su* sweet corn hybrids, Waters et al. [23] showed that moistened seed had no impact on plant growth and development. Our results are not in agreement with this previous report. Data from Fig. 3 showed the higher water supply, the higher the fresh weight in super sweet *sh2* maize plants sowed and grown in plug cell tray under the best greenhouse environmental facilities. In field adult plants, generally, when water availability is limited, the root: shoot ratio of plants increases because roots are less sensitive than shoots to growth inhibition by low water potentials

[36], however, an inverse response was found in seedlings grown in cell plug trays at the transplant stage. When dry weight data were plotted against time (Fig. 4), we found significant differences in the root: shoot ratio at the transplant stage, which decreased during the first four post-transplant weeks to pots. Significant differences between water supply treatments disappeared after transplant as well. A more detailed analysis of the dry weight differences at the transplant stage showed that they were the result of higher root dry weight accumulation in those seedlings 100% watered with no significant differences in shoot dry weight (Fig. 5). In agreement with Ahmed et al. [37], the growth of maize root system is regulated by its endogenous genetic program, but also by interactions with the soil and changing environmental conditions.

Leaf dry mass per unit leaf area (LMA) was introduced as a concept in the analysis of plant growth and defined at the plant level as the total leaf dry mass divided by the total leaf area. LMA is used in crop simulation models to predict leaf area from leaf dry mass. In this way, Drouet et al. [38] showed that LMA increased about two-fold during the first 37 days after sowing in maize plants; similar increments for the sweet *sh2* maize hybrids tested in our experiments were found (Fig. 6) and let to explain the decrease in the root: shoot ratios previously discussed (Fig. 4). Although LMA has been considered either as a constant for the whole growth cycle, as a function of the plant development stage and it has been correlated with variables as diverse as net photosynthesis, relative growth rate, leaf structure and yield [38], our results are not in agreement because LMA increased during our experiment (Fig. 6).

The susceptibility of plants to drought stress varies in dependence of stress degree, different accompanying stress factors, plant species, and their developmental stages [39]. Water deficit stress is more vulnerable at the germination and early seedling growth stages of plants [12]. Identifying corn hybrids with faster rates of seedling emergence and leaf appearance may allow development of a corn crop that achieves earlier canopy closure and better seasonal light interception. Water deficits reduce the number of leaves per plant and individual leaf size by decreasing the soil's water potential. Leaf area expansion depends on leaf turgor, temperature, and assimilating supply for growth. Our data showed that during nursery and four weeks post-transplant, the higher water supply the higher total leaf area (Fig. 7) but with significant differences in response to the three sweet maize hybrids tested. Results can be explained by higher as RLA as RLAE growth parameters (Table 1).

A common adverse effect of water stress on crop plants is the decrease in fresh and dry biomass production. Kamara et al. [40] revealed that water deficit imposed at various developmental stages of maize reduced total biomass accumulation at silking by 37%, at grain-filling period by 34% and at maturity by 21%. In the same way, Anjum et al. [10] indicated that drought stress in maize led to considerable decline in net photosynthesis (33.22%), transpiration rate (37.84%), stomatal conductance (25.54%), water use efficiency (50.87%), intrinsic water use efficiency (11.58%) and intercellular CO₂ (5.86%) as compared to well water control. In agreement, our results showed a fresh and dry weight decrease of 19-36% and 13-28% respectively at the transplant stage when water supply decrease from 100% to 50% daily evaporation (from Fig. 3 and 5). During the last four weeks, post-transplant dry weight differences were maintained and a decrease in RGR was found (Table 2). In previous reports [41,42,43] changes in RGR were associated exclusively with NAR changes. However, Fig. 8 showed RGR as the result of additive positive effects of NAR and LAR.

Water deficit stress has a profound adverse effect on photosynthesis, photosynthetic apparatus, and pigment content in adult field maize crops [44]. As a result, a considerable decline in net photosynthesis has been reported in drought-stressed maize in contrast to well-watered control plants [10,12]. The same response during nursery and early post-transplant in our experiment because NAR (the "photosynthetic" component of RGR) was higher in full watered plants (Table 2). On the other hand, the decrease in SLA in plants fully watered is an advice of higher leaf thickness (a morphological trait related to higher photosynthetic rate).

A water stress **alters** biomass allocation to shoots and roots or by changing the metabolic activities in these organs. According to the theory of functional balance, plants increase the allocation of biomass to shoots if limited resources aboveground affect carbon gain, such as light and CO₂. Similarly, plants increase biomass allocation to roots in the presence of low levels of belowground resources, such as water and nutrients [45]. Allometries from Table 3 showed that a **full** water supply during nursery, redirect photo assimilate partitioning from shoots and particularly to stems during at least four post-transplant weeks.

Maize yield was positively correlated in field adult plants with total root number and the ratio of root mass below 20 cm to total root mass [18], while Rattin et al. [21] showed a close coordination between roots and shoot growth as well. Long-distance signaling between the root and shoot is intrinsic to the control of plant growth and development, and its response to environmental perturbations. With respect to plant water use and control of plant water relations, Vandeleur et al. [46] currently know more about root-to-shoot signaling because of the important role of root-derived signals in informing the shoot of changes in soil water content. Although emerging evidence suggests that there may be a requirement for shoot water demand to be communicated to roots, to allow root water uptake to fulfil the shoots requirements [47], shoot-to-root signals have been investigated only rarely in respect to plant water relations [48,49]. Our results showed positive relationships between RLAE (Fig. 9A), RGR (Fig. 9B), NAR (Fig. 9C) and root dry weight during the early post-transplant growth with the higher values from plants grown with the higher water supply during nursery.

Cramer et al. [50] defined abiotic stress as an environmental condition that reduces growth and yield below optimal levels. Puig et al. [51] and Chen et al. [52] concluded that plants can sense the volume of the rooting space available, and a limited number of studies on individual roots have shown that plant roots may sense the identity of neighboring roots and respond accordingly [53,54]. Our results showed that a water stress during nursery would be considered as a limiting source to decrease maize early biomass accumulation and, presumably, plant yield. However, the validation of this hypothesis goes beyond the present experimental design and is the matter of further experiments.

5. CONCLUSIONS

Contrary to previous reports, our results showed that a water stress during nursery decrease seedling emergence and may be avoided during nursery for plants sown in plug trays. On the other hand, a very early water stress during nursery decrease early post-transplant biomass accumulation, leaf area expansion and change photo assimilate partitioning. These results would force to a change in the commonly accepted water management when a transplant routine from plug trays was used for improving agricultural intensification.

REFERENCES

1. Neumann K, Verburg PH, Stehfest E, Müller C. The yield gap of global grain production: A spatial analysis. *Agr. Syst.* 2010; 103(5): 316-326.
2. Licker R, Johnston M, Foley JA, Barford C, Kucharik CJ, Monfreda C, Ramankutty N. Mind the gap: how do climate and agricultural management explain the 'yield gap' of croplands around the world?. *Global Ecol. Biogeogr.* 2010; 19(6): 769-782.
3. Dietrich JP, Schmitz C, Müller C, Fader M, Lotze-Campen H, Popp A. Measuring agricultural land-use intensity—A global analysis using a model-assisted approach. *Ecol. Model.* 2012; 232: 109-118.
4. Zhang X, Cai X. Climate change impacts on global agricultural land availability. *Environ. Res. Lett.* 2011; 6(1): 014014.
5. Di Benedetto A, Rattin J. Transplant in Sweet Maize: A Tool for Improving Productivity. *The Americas Journal of Plant Sci. Biotech.* 2008; 2(2): 96-108.
6. El-Hamed K, Elwan M, Shaban W. Enhanced sweet corn propagation: studies on transplanting feasibility and seed priming. *Vegetable Crops Res. Bull.* 2011; 75, 31-50.
7. Orosz F. Morphological plant properties of sweet corn cultivated with different technologies. In: Sixth International Scientific Agricultural Symposium "Agrosym 2015", Jahorina, Bosnia and Herzegovina, 2015; 279-284.
8. García A, Guerra LC, Hoogenboom G. Impact of planting date and hybrid on early growth of sweet corn. *Agron. J.* 2009; 101(1): 193-200.
9. Orgaz F, Fernández MD, Bonachela S, Gallardo M, Fereres E. Evapotranspiration of horticultural crops in an unheated plastic greenhouse. *Agr. Water Manage.* 2005. 72(2): 81-96.
10. Anjum SA, Xie XY, Wang LC, Saleem MF, Man C, Lei W. Morphological, physiological and biochemical responses of plants to drought stress. *Afr. J. Agric. Res.* 2011; 6(9): 2026-2032.
11. Huber AE, Bauerle TL. Long-distance plant signaling pathways in response to multiple

- stressors: the gap in knowledge. *J. Exp. Bot.* 2016; 67(7): 2063-2079.
12. Lata C, Muthamilarasan M, Prasad M. Drought stress responses and signal transduction in plants. In: *Elucidation of Abiotic Stress Signaling in Plants*. 2015; 195-225. Springer New York.
 13. El-Hendawy S.E, Hokam EM, Schmidhalter U. Drip irrigation frequency: the effects and their interaction with nitrogen fertilization on sandy soil water distribution, maize yield and water use efficiency under Egyptian conditions. *J. Agron. Crop Sci.* 2008; 194(3): 180-192.
 14. Oktem A, Simsek M, Oktem AG. Deficit irrigation effects on sweet corn (*Zea mays saccharata* Sturt) with drip irrigation system in a semi-arid region: I. Water-yield relationship. *Agr. Water Manage.* 2003; 61(1): 63-74.
 15. Oktem A. Effects of deficit irrigation on some yield characteristics of sweet corn. *Bangladesh J. Bot.* 2008; 37(2): 127-131.
 16. Ben-Gal A, Karlberg L, Jansson PE, Shani U. Temporal robustness of linear relationships between production and transpiration. *Plant Soil.* 2003; 251(2): 211-218.
 17. Bray EA. Plant responses to water deficit. *Trends Plant Sci.* 1997; 2(2): 48-54.
 18. Wang Z, Ma BL, Gao J, Sun J. Effects of different management systems on root distribution of maize. *Canadian J. Plant Sci.* 2014; 95(1): 21-28.
 19. Campos H, Cooper M, Habben JE, Edmeades GO, Schussler JR. Improving drought tolerance in maize: a view from industry. *Field Crops Res.* 2004; 90(1): 19-34.
 20. Sarwat M, Tuteja N. Hormonal signaling to control stomatal movement during drought stress. *Plant Gene.* 2017; 11: 143-153.
 21. Rattin J, Wagner P, Ferreyri D, Riverti A, Giardina E, Di Benedetto A. Roots partially drive super sweet maize yield. *J. Exp. Agr. Int.* 2017; 16(6): 1-17.
 22. Styer RC, Koransky DS (Eds). *Plug and transplant production. A grower's Guide*, Ball Publishing, Batavia, Illinois, USA. 1997.
 23. Waters L, Burrows RL, Bennett MA, Schoenecker J. Seed moisture and transplant management techniques influence sweet corn stand establishment, growth, development, and yield. *J. Am. Soc. Hortic. Sci.* 1990; 115(6): 888-892.
 24. Di Benedetto A. Alternative substrates for potted ornamental plants based on Argentinean peat and Argentinean river waste: a review. *Floriculture Plant Ornamental Biotech.* 2007; 1(2): 90-101.
 25. Maguire JD. Speed of germination- Aid in selection and evaluation for seedling emergence and vigor. *Crop Sci.* 1962; 2(2): 176-177.
 26. Warton DI, Duursma RA, Falster DS, Taskinen S. SMATR 3-an R package for estimation and inference about allometric lines. *Methods Ecol. Evol.* 2012; 3(2): 257-259.
 27. Adetimirin VO. Stand establishment and early field vigour variation in a tropicalised shrunken-2 maize population. *Field Crops Res.* 2008; 108(2): 143-149.
 28. Kumari J, Gadag RN, Jha GK, Joshi HC, Singh RD. Combining ability for field emergence, kernel quality traits, and certain yield components in sweet corn (*Zea mays* L.). *J. crop Improve.* 2008; 22(1): 66-81.
 29. Kleinhenz MD. Sweet corn variety trials in Ohio: recent top performers and suggestions for future evaluations. *HortTechnology.* 2003; 13(4): 711-718.
 30. Hassell, R. L., Dufault, R. J., & Phillips, T. L. (2003). Low-temperature germination response of su, se, and sh2 sweet corn cultivars. *HortTechnology*, 13(1), 136-141.
 31. Ouyang X, van Voorthuysen T, Toorop PE, Hilhorst HW. Seed vigor, aging, and osmopriming affect anion and sugar leakage during imbibition of maize (*Zea mays* L.) caryopses. *Int. J. Plant Sci.* 2002; 163(1): 107-112.
 32. Cardoso ET, de Melo Sereno MC, Neto JB. Genetic progress for adaptative traits in sweet corn populations. *Maydica.* 2002; 47(2): 121-126.
 33. Zhao G, Zhong T, Zheng D. Improving the field emergence performance of super sweet corn by sand priming. *Plant Prod. Sci.* 2009; 12(3): 359-364.
 34. Rattin J, Valinote JP, Gonzalo R, Di Benedetto A. Transplant and a change in plant density improve sweet maize (*Zea mays* L.) yield. *Amer. J. Exp. Agric.* 2015; 5(4): 336-351.
 35. Varier A, Vari AK, Dadlani M. The subcellular basis of seed priming. *Current Sci. India.* 2010; 99(4): 450-456.
 36. Wu Y, Cosgrove DJ. Adaptation of roots to low water potentials by changes in cell wall extensibility and cell wall proteins. *J. Exp. Bot.* 2000; 51(350): 1543-1553.
 37. Ahmed MA, Zarebanadkouki M, Kaestner A, Carminati A. Measurements of water uptake of maize roots: the key function of lateral roots. *Plant Soil.* 2016; 398(1-2): 59-77.
 38. Drouet JL, Pagès L, Serra V. Dynamics of leaf mass per unit leaf area and root mass per unit

- root volume of young maize plants: implications for growth models. Eur. J. Agron. 2005; 22(2): 185-193.
39. Demirevska K, Zasheva D, Dimitrov R, Simova-Stoilova L, Stamenova M, Feller U. Drought stress effects on Rubisco in wheat: changes in the Rubisco large subunit. Acta Physiol. Plant. 2009; 31(6): 1129.
40. Kamara AY, Menkir A, Badu-Apraku B, Ibikunle O. The influence of drought stress on growth, yield and yield components of selected maize genotypes. J. Agr. Sci. 2003; 141(1): 43-50.
41. Gandolfo E, De Lojo J, Gómez D, Pagani A, Molinari J, Di Benedetto A. Anatomical changes involved in the response of *Impatiens wallerana* to different pre-transplant plug cell volumes and BAP sprays. Eur. J. Hortic. Sci. 2014; 79(4): 226-232.
42. Di Benedetto A, Galmarini C, Tognetti J. Exogenous cytokinin promotes *Epipremun aureum* L. growth through enhanced dry weight assimilation rather than through changes in partitioning. Amer. J. Exp. Agric. 2015; 5(5): 419-434.
43. Di Matteo J, Rattin J, Di Benedetto A. Increase of spinach growth through the use of larger plug cell volume and an exogenous BAP spray. American J. Exp. Agric. 2015; 6(6): 372-383.
44. Wahid A, Rasul E, Rao RA, Iqbal RM. Photosynthesis in leaf, stem, flower and fruit. Handbook of Photosynthesis. 2005; 2: 479-497.
45. Lavinsky AO, Magalhães PC, Ávila RG, Diniz MM, de Souza TC. Partitioning between primary and secondary metabolism of carbon allocated to roots in four maize genotypes under water deficit and its effects on productivity. Crop J. 2015; 3(5): 379-386.
46. Vandeleur RK, Sullivan W, Athman A, Jordans C, Gilliam M, Kaiser BN, Tyerman SD. Rapid shoot-to-root signalling regulates root hydraulic conductance via aquaporins. Plant Cell Environ. 2014; 37(2): 520-538.
47. Franks PJ, Drake PL, Froend RH. Anisohydric but isohydrodynamic: seasonally constant plant water potential gradient explained by a stomatal control mechanism incorporating variable plant hydraulic conductance. Plant Cell Environ. 2007; 30(1): 19-30.
48. Levin M, Resnick N, Rosianskey Y, Kolotilin I, Wininger S, Lemcoff JH, Cohen S, Galili G, Koltai H, Kapulnik Y. Transcriptional profiling of Arabidopsis thaliana plants' response to low relative humidity suggests a shoot-root communication. Plant Sci. 2009; 177(5): 450-459.
49. Sakurai-Ishikawa JUNKO, Murai-Hatano, MARI, Hayashi H, Ahamed A, Fukushi K, Matsumoto T, Kitagawa Y. Transpiration from shoots triggers diurnal changes in root aquaporin expression. Plant Cell Environ. 2011; 34(7): 1150-1163.
50. Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants: a systems biology perspective. BMC Plant Biol. 2011; 11(1): 163.
51. Puig J, Pauluzzi G, Guiderdoni E, Gantet P. Regulation of shoot and root development through mutual signaling. Molec. Plant. 2012; 5(5): 974-983.
52. Chen BJ, During HJ, Vermeulen PJ, Kroon H, Poorter H, Anten NP. Corrections for rooting volume and plant size reveal negative effects of neighbour presence on root allocation in pea. Funct. Ecol. 2015; 29(11): 1383-1391.
53. Hellmann E, Gruhn N, Heyl A. The more, the merrier: cytokinin signaling beyond Arabidopsis. Plant Signaling Behavior. 2010; 5(11): 1384-1390.
54. Bartrina I, Otto E, Strnad M, Werner T, Schmulling T. Cytokinin regulates the activity of reproductive meristems, flower organ size, ovule formation, and, thus, seed yield in *Arabidopsis thaliana*. Plant Cell. 2011; 23(1): 69-80.