

Original Research Article

Dispersion Syndrome, Ecological Sucession and Structure of the Arbor Component in an Atlantic Forest Fragment, Pernambuco

ABSTRACT

The Atlantic Forest is considered one of the richest areas of biodiversity on the planet. However, it has been continuously threatened and must receive strategies aiming for its restoration and conservation. Thus, the objectives of this work were to evaluate the arboreal component, to classify the ecological succession and the dispersion syndrome and to analyze the structure of a fragment located in the city of Goiana, Pernambuco. For this, 10 plots of 10m x 25m were allocated, where individuals with height greater than or equal to 5.0 m and chest circumference (CC) equal to or greater than 15 cm were sampled. As for dispersion syndrome, species were classified in autochoric, zoochoric and anemochoric, and ecological succession were classified in pioneers, early secondary and late secondary. For structural analysis, vegetation was divided into three classes, representing 0-50%, 50-80% and 80-100% of the maximum total height, and the diametrical structure was distributed in 7 classes with intervals of 5 cm between them. A total of 422 individuals were found, distributed in 47 species and 28 families. In relation to the dispersion syndrome, species were predominantly zoochoric. As for the ecological succession, most species of the area were early and late secondaries. Considering the diametric and hypsometric analysis, it can be seen that individuals were mainly concentrated in the first classes of diameter and height, indicating that the fragment has a high regeneration potential. Thus, with all analysis performed, it can be suggested that the studied fragment is in intermediate stage of succession.

Keywords: Phytosociology; biodiversity; zoochory.

1. INTRODUCTION

The tropical rainforest (Atlantic Forest) is considered one of the world's hotspots [1] considering that it has one of the richest biodiversity on Earth. However, it is under severe and progressive threatening process. Great part of its forest cover disappeared, with only approximately 11.73% left, which are very fragmented, with areas under 50 hectares that are considered insufficient for protection and maintenance of the existing biodiversity [2]. For this reason, it is necessary to develop studies that address ecological and biological characteristics of fragmented environments [3].

Among the ecological studies applied in environments under these conditions, phytosociology verifies the relationships between elements of plant communities and floristic that seeks to understand the distribution of these elements and their geographical occurrence. These factors are fundamental for a satisfactory understanding of forest environments, making it possible to construct a representation of its functioning [4, 5, 6].

Others analyzes, such as the dispersion of seeds and the characterization of successional groups enrich information related to floristic and phytosociological data. Dispersion is the action by which plants remove their seeds from their vicinity to minimize effects of competition and predation [7, 8]. This mechanism is efficient to provide propagules, subsiding the natural regeneration process [9], also

contributing to the planning and development of recovery actions in environments with intense loss of biodiversity as the present study.

The classification of successional groups enables the comprehension of adaptative qualities of each specie by the understanding of its distribution in the forest environments [10]. It helps to verify the successional phase and identify the factors that may be interfering in the ecosystems evolution for more complex successional levels and thus, contribute to the success in the ecosystems recovery, assisting in the selection of species which may vegetate degraded environments.

Finally, structural evaluations of vegetation, both horizontal and vertical, stand out. These are fundamental to understand the stage of forest development [11], serving as indicators of biodiversity and ecosystem services [12]. It is performed using parameters of density, dominance, frequency, coverage and importance values among others, providing information to assist in management and conservation of forest ecosystems [13, 12]. Moreover, it contributes to the understanding of species aggregation in the forest communities and in the habit of each population, besides of its ecological behavior [14].

Many studies have addressed only one of the analysis described above [15, 16, 17]. However, in order to clearly and significantly understand the dynamics and composition of a forest area, it is necessary to use as many analyzes as possible in order to acquire sufficient data that allow the establishment of efficient management and conservation strategies.

Phytosociological and floristic composition studies of vegetation are the basis for the development of strategies and public policies aimed at the conservation and restoration of forests remnants, providing information about species distribution, abundance, structure and dynamics in different locations. It is also important, for example, to provide a consistent basis for the creation of protected areas, especially, in fragmented ecosystems with relevant anthropic impact activities as it offers fundamental subsidies for the recovery management of fragments [18].

In this context, the objectives of this work were to characterize the arboreal component, to classify the dispersion syndromes and the ecological succession, and to analyse the diametric and hypsometric structures of a fragment of tropical rain forest located in the city of Goiana, Pernambuco.

2. MATERIAL AND METHODS

The present study was developed in a fragment of tropical rainforest with approximately 51.32 hectares located in the city of Goiana, PE (latitude 7 ° 37'30 "S, longitude 34 ° 57'30"W). The climate of the region is As' type, tropical humid with autumn-winter rain [19], with annual average temperatures of 24.8°C and annual precipitation of 1.990 mm [20].

To enable identification and classification of species in the area, 10 plots of 10m x 25m were randomly installed. Robustness of data was verified by sampling adequacy in which a probability level of 5% was used, with a maximum permissible error of 20%. Within plots, individuals with height greater than or equal to 5.0 m and chest circumference (CC) equal to or greater than 15 cm were sampled. Measurement for height occurred visually, taking a 6-m rod as reference, and for the CC, a metric tape was used.

Sampled species were botanically classified using APG IV [21]. Collected botanical material was herborized and deposited in the Herbarium Sérgio Tavares (HST) of the Federal Rural University of Pernambuco for identification purposes.

For the phytosociological analysis, the following parameters were determined: absolute density (AD) (ind./ha); relative density (RD) (%); absolute frequency (AF); relative frequency (RF); absolute dominance (ADo) (m² / ha); relative dominance (RDo); value of importance (VI) and the Shannon diversity index using Mata Nativa software version 4 [22].

Subsequently, individuals identified at species level were classified by their dispersion syndrome and ecological group. For dispersion, the following categories were adopted: Anemochory (Ane), Autochory (Aut) and Zoochory (Zoo), following methodology established by [23]. Regarding the ecological group, species were classified as pioneers, early secondary and late secondary, following [24].

Finally, structural evaluation of the fragment was given. The hypsometric structure followed methodology established by [25] dividing the vegetation into three classes representing 0-50%, 50-80% and 80-100% of the total maximum height, respectively. The diametric structure was distributed in 7 classes with intervals of 5 cm between them.

3. RESULTS AND DISCUSSION

Phytosociology

A total of 422 individuals were registered and identified, in which the floristic richness was 47 species distributed in 28 botanical families (Table 1). Among sampled families, 25 were identified and three were indeterminate. Regarding species, 35 were identified at species level, nine at gender level and one at family level. Families with the greatest species richness were Fabaceae (17.78) and Myrtaceae (8.89). Six families presented two species (4.44) each: Lecythidaceae, Anacardiaceae, Moraceae, Clusiaceae, Annonaceae and Chrysobalanaceae. The others presented one species (2.22) each.

The representativeness of Fabaceae family is justified by the fact that this group stands out among the most frequent families in the Atlantic Forest [26]. The results found corroborate with the study by [6], who evaluated a tree community of a Atlantic forest fragment in Pernambuco, and with the study of [27] in areas of Ombrophylous Dense Forest in Rio de Janeiro, where families that presented the largest number of species were also Fabaceae and Myrtaceae. [28] highlight the Fabaceae family in their study and justify that it is common for this family to appear in floristic and ecological surveys in Brazil.

Shannon-Wiener index (H') for the study area was 3.39 nats / individual, higher than 3.26 nats / individual found by [6] and less than 3.91 nats / individuals found by [29] in a forest fragment in Cabo de Santo Agostinho, and [30], which found 3.68 nats / individual in a Atlantic forest fragment in Igarassu, both in the State of Pernambuco. Differences in succession stages, different sampling methodologies, inclusion levels, efforts for taxonomical identification, as well as floristic variations in different communities can explain variations occurring in diversity values in areas of the same phytogeographic region [31].

Table 1. Phytosociological parameters, dispersion syndrome and ecological group of species sampled in an area of Ombrophylous Dense Forest, Pernambuco-Brazil. Values are in descending order of importance value (IV). Where: AD - absolute density (ind./ha); RD - relative density (%); AF - absolute frequency; RF - relative frequency; ADo - absolute dominance (m² / ha); RDo - relative dominance.

Species	Family	Disp. Syn.	Ecol. Group	AF	RF	AD	RD	ADo	RDo	IV
<i>Pogonophora schomburgkiana</i> Miers ex Benth	Peraceae	Aut	ST	80	5,1	180	10,66	0,91	3,49	19,25
<i>Eschweilera ovata</i> Mart. ex Miers	Lecythidaceae	Zoo	SI	70	4,46	96	5,69	2,04	7,78	17,93
<i>Tapirira guianensis</i> Aubl.	Anacardiaceae	Zoo	PI	70	4,46	60	3,55	2,44	9,31	17,33
<i>Manilkara salzmannii</i> (A.DC.) H.J.Lam	Sapotaceae	Zoo	SC	90	5,73	108	6,4	1,08	4,14	16,27
<i>Lecythis pisonis</i> Cambess.	Lecythidaceae	Aut	ST	40	2,55	120	7,11	1,42	5,41	15,06
<i>Coccoloba mollis</i> Casar.	Polygonaceae	Zoo	PI	70	4,46	76	4,5	1,54	5,88	14,84
<i>Thyrsodium spruceanum</i> Benth.	Anacardiaceae	Zoo	SI	90	5,73	84	4,98	0,64	2,45	13,16
Indetermined 3	SC	SC	SC	90	5,73	68	4,03	0,82	3,12	12,88
<i>Myrcia</i> sp 1	Myrtaceae	Zoo	SI	70	4,46	92	5,45	0,64	2,46	12,37
<i>Clusia nemorosa</i> G.Mey.	Clusiaceae	Zoo	ST	40	2,55	64	3,79	1,28	4,9	11,24
<i>Dialium guianensis</i> (Aubl.) Sandwith	Fabaceae	SC	ST	20	1,27	20	1,18	2,01	7,68	10,14
<i>Apuleia leiocarpa</i> (Vogel) J.F.Macbr.	Fabaceae	Ane/Aut	SI	30	1,91	36	2,13	1,37	5,22	9,26
Indetermined 2	SC	SC	SC	50	3,18	52	3,08	0,71	2,7	8,96

<i>Eriothea crenulicalyx</i> A.Robyns	Malvaceae	SC	SC	30	1,91	64	3,79	0,77	2,95	8,65
<i>Protium heptaphyllum</i> (Aubl.) Marchand	Burseraceae	Zoo	SI	50	3,18	52	3,08	0,49	1,86	8,12
<i>Symphonia globulifera</i> L. f.	Clusiaceae	Zoo	SC	10	0,64	48	2,84	1,19	4,53	8,01
<i>Brosimum rubescens</i> Taub.	Moraceae	Zoo	SI	50	3,18	28	1,66	0,6	2,29	7,13
<i>Himanthantus bracteatus</i> (A. DC.) Woodson	Apocynaceae	Ane	SI	60	3,82	32	1,9	0,25	0,96	6,68
<i>Maytenus distichophylla</i> Mart. ex Reissek	Celastraceae	Zoo	ST	40	2,55	48	2,84	0,22	0,83	6,22
<i>Byrsonima sericea</i> DC.	Malpighiaceae	Zoo	SI	30	1,91	16	0,95	0,61	2,31	5,17
<i>Hirtella hebeclada</i> Moric. ex DC.	Chrysobalanaceae	Zoo	ST	20	1,27	28	1,66	0,55	2,09	5,02
<i>Bowdichia</i> sp.	Fabaceae	SC	PI	20	1,27	12	0,71	0,78	2,99	4,98
<i>Guatteria pogonopus</i> Mart.	Annonaceae	Zoo	ST	30	1,91	36	2,13	0,12	0,46	4,5
<i>Samanea</i> sp.	Fabaceae	Zoo	PI	20	1,27	8	0,47	0,7	2,69	4,44
<i>Psidium guineense</i> Sw.	Myrtaceae	Zoo	SC	30	1,91	28	1,66	0,19	0,73	4,3
<i>Miconia</i> sp.	Melastomataceae	Zoo	SC	30	1,91	24	1,42	0,17	0,64	3,97
<i>Myrcia</i> sp 2	Myrtaceae	Zoo	SI	30	1,91	20	1,18	0,18	0,69	3,79
<i>Ocotea glomerata</i> (Nees) Mez	Lauraceae	Zoo	SI	30	1,91	20	1,18	0,18	0,69	3,78
<i>Couepia impressa</i> Prance	Chrysobalanaceae	SC	SC	20	1,27	20	1,18	0,3	1,13	3,59
<i>Inga</i> sp.	Myrtaceae	Zoo	PI	20	1,27	12	0,71	0,37	1,4	3,39
<i>Annona</i> sp.	Annonaceae	Zoo	SI	30	1,91	12	0,71	0,2	0,76	3,38
<i>Schefflera morototoni</i> (Aubl.) Maguire et al.	Araliaceae	Zoo	SI	20	1,27	8	0,47	0,41	1,57	3,31
<i>Albizia pedicellaris</i> (DC.) L.Rico	Fabaceae	SC	PI	20	1,27	8	0,47	0,39	1,5	3,25
<i>Myrtaceae</i> sp.	Myrtaceae	SC	SC	30	1,91	12	0,71	0,04	0,16	2,78
<i>Zolernia</i> sp.	Fabaceae	SC	SC	20	1,27	12	0,71	0,09	0,36	2,35
<i>Allophylus edulis</i> (A.St.-Hil. et al.) Hieron. ex Niederl.	Sapindaceae	Zoo	PI	20	1,27	12	0,71	0,08	0,32	2,3
<i>Manilkara salzmannii</i> (A. DC.) H.J.Lam	Sapotaceae	Zoo	PI	20	1,27	12	0,71	0,05	0,18	2,17
<i>Handroanthus impetiginosus</i> (Mart. ex DC.) Mattos	Bignoniaceae	Aut	ST	10	0,64	16	0,95	0,07	0,27	1,86
<i>Erythroxylum squamatum</i> Sw.	Erythroxylaceae	Zoo	ST	10	0,64	12	0,71	0,07	0,28	1,63
<i>Albarena cochliacarpus</i> (Gomes) Barneby & J.W. Grimes	Fabaceae	SC	ST	10	0,64	8	0,47	0,12	0,46	1,57
Indetermined 1	SC	SC	SC	10	0,64	8	0,47	0,03	0,1	1,21
<i>Simarouba amara</i> Aubl.	Simaroubaceae	Zoo	SI	10	0,64	4	0,24	0,03	0,1	0,98
<i>Senna phlebadenia</i> H.S. Irwin & Barneby	Fabaceae	Aut	SI	10	0,64	4	0,24	0,02	0,09	0,96
<i>Luehea ochrophylla</i> Mart.	Malvaceae	Ane	SI	10	0,64	4	0,24	0,01	0,04	0,91
<i>Artocarpus heterophyllus</i> Lam.	Moraceae	Ane	SI	10	0,64	4	0,24	0,01	0,04	0,91
Total				1570	100	1688	100	26,19	100	300

Species with the highest relative frequencies, in decreasing order, were: *Manilkara salzmannii*, *Thyrsodium spruceanum*, Undetermined 3, *Pogonophora schomburgkiana*. Species that presented the highest values for relative density, in descending order, were: *Pogonophora schomburgkiana*, *Lecythis pisonis*, *Manilkara salzmannii*, *Eschweilera ovata* and *Myrcia* sp 1, totalizing 35.31%.

In terms of relative dominance, the species *Tapirira guianensis* presented the highest value, followed by *Eschweilera ovata*, *Dialium guianensis*, *Coccoloba mollis* and *Lecythis pisonis*, representing 36.06% of the total. According to [6], it can be inferred that these species are better exploiting the natural resources of the environment.

The five species that presented the highest values of importance (VI) in decreasing order were: *Pogonophora schomburgkiana*, *Eschweilera ovata*, *Tapirira guianensis*, *Manilkara salzmannii*, and *Lecythis pisonis*.

Among these *Manilkara salzmannii*, *Pogonophora schomburgkiana*, *Lecythis pisonis* and *Eschweilera ovata* stand out for showing relevance in three parameters analyzed. These species are often found in studies performed in the region. The ecological domain of few species is frequently observed in tropical forests ecosystems, as in the present research [32], evidencing the importance of them to subsidize conservation strategies in this fragmented environment, especially because of urban growth and alcohol and sugar industries [33].

Dispersion syndrome

It was observed that the fragment presented in its composition 23 zoochoric (72%), six anemochoric (19%) and three autochoric (9%) species. This pattern of zoochoric dispersion domain is often observed in tropical forests [34, 35, 36]. The proportions of species among these variables for zoochoric were 65.1%, 74% and 58%, respectively, 22.5%, 6% and 13% for anemochoric and 22%, 20% and 29% for autochoric.

The strong interdependence of the Atlantic forest ecosystem with the faunal population to maintain its dispersion evidences the important role of this syndrome in the conservation of species of advanced ecological successional stages [37]. The highest seed richness with zoochoric dispersion in the bank is important for maintenance of the propagating fauna of propagules, mainly at the beginning of the regeneration of the forest after disturbance, influencing in its special distribution and frequency of species in the forest [38].

The reduced proportion of anemochoric species may be related to spatial restriction of this syndrome, as the seeds tend to remain in the vicinity of the mother plant, generating a greater aggregation effect [39]. Regarding the anemochoric, the low indices occurred because it is a limited characteristic of few families, confirming a successional stage varying between intermediary and advanced in the fragment of this study, as the greater incidence of anemochoric species is commonly observed in initial successional areas [40].

Ecological succession

Regarding ecological groups, a relevant aspect of this research is the number of species both initial secondary (S_i) and late secondary (S_l) stages, which together represent 57.44% of species and 67.1% of individuals inventoried. On the other hand, initial secondary species represented 28.20% of individuals grouped in 31.91% of species, and late secondary species represented 38.86% of individuals occurring in 25.53% of species. Initial pioneers represented 14.9% of species and 16.59% of individuals of the fragment. No data were obtained from 13 species (27.66%).

Low representativeness of pioneer species may suggest forest maturity [41]. According to [42], occurrence of late secondary species typical of the Atlantic forest is a good indicator of resilience of this community.

Regarding ecological groups, most of the species found are late secondary (S_l). The species *Pogonophora schomburgkiana* (10.7%), *Lecythis pisonis* (7.1%) and *Pisonia ambiguous* (6.4%) were more representative in this group. *Eschweilera ovata* (5.7%), *Myrcia* sp (5.4%) and *Thyrsodium spruceanum* (5%) stood out among initial secondary species (S_i) and the species *Coccoloba mollis* (4.5%) and *Tapirira guianensis* (3.55%) were the most representative among pioneers.

Considering the low floristic representativeness of pioneer species and the number of late secondary (S_l), the expression of an intermediate stage in successional development of the forest was found in this relation [10].

Classification of species in their respective ecological groups has encountered two primary factors. The first one is that criteria used differ between authors, which leads some species to be classified into distinct groups. The second one refers to the fact that the same species, depending on its genetic characteristics, may respond differently to environmental conditions in regions with different soil and

climate, since these responses do not occur for a single environmental factor alone [15]. However, classification in ecological groups combined with structure studies may provide better information on species behavior within a given area.

Structural analysis

Understanding the structure is an essential factor for proper management of a forest, both for economic and environmental purposes, and is also a relevant indicator of the biodiversity of the area. [43, 44]. In addition, it allows identifying how species are distributed, also providing information on successional dynamics [45].

In terms of diametric structure, seven classes with 5 cm intervals were found between each of them in this study, in which the first class comprised individuals from 3 to 8 cm (center 5.5 cm) and the last class ≥ 35 , 5 cm. As can be seen in figure 1, the curve followed an “inverted J” distribution, which is a pattern for native forests, with the majority of individuals concentrated in the first class, following a progressive downward trend. Results similar to this were found by [6] in a fragment of Atlantic forest in the municipality of Moreno, in Pernambuco, and by [46], in a fragment of Atlantic forest in Recife, located in the same state.




Fig. 1. Diametric distribution of arboreal individuals sampled in an Atlantic forest fragment in Goiana, Pernambuco, 2017.

These results indicate that the fragment is composed mainly of individuals who have reached maturity in recent time. In addition, this suggests that the forest presents potential for regeneration, also indicating that there is recruitment in the area [47, 48]. This type of forest distribution is the most desirable, in a way that allows individuals to continue to participate in natural dynamics [49].

In order to better understand the dynamics of tropical forests in relation to their structure, it is necessary to know the community that compose them, as well as to identify the behavior of each species [50]. Thus, ten species with the highest VI were analyzed (Figure 2).




Fig. 2. Diametric distribution of ten species with the highest VI in a fragment of Atlantic forest in Goiana, Pernambuco, 2017.

Of the ten species selected, most showed similar results to the general arrangement, only *T. guianensis*, *L. pisonis* and *C. nemorosa* did not follow the “inverted J” distribution. *T. guianensis* presented the majority of individuals distributed in class 4, different from *L. pisonis* in which the majority of individuals appeared in class 2, and of *C. nemorosa* that presented the same number of individuals equally distributed in classes 2, 3 and 4. These variations may be related to the population ecology of each species, where there are certain discontinuities in the distributions, and there may be a total absence of young individuals in some species [51]. This was the case of Undetermined 3, which did not present any individuals in the first class of diameter.

Vertical structure refers to characteristics of the vegetation as a function of vertical height of the trees, being an essential component that helps to describe the ecosystem [52]. Formation of a vertical stratification occurs commonly in tropical forests, where these stands are composed of plant species that have different modes of occupation in response to different light requirements, which may justify the enormous diversity in structurally more complex ecosystems [53].

Therefore, for hypsometric structure, distribution was carried out in three classes, in which the first class included individuals with height ≤ 11.5 cm; in the second, individuals 11.5 to 18.4 cm; and in the last class, individuals with height between 18.4 and 23.1 cm. It was observed that the majority of individuals (69.62%) are concentrated in the first class, followed by the second one with 27.75% and the last class comprising 2.63% of the individuals in the area (figure 3). These results coincide with those found in tropical forests, which present the highest number of individuals in lower stages, characterizing a negative exponential distribution [54]. The same results were found by [6] in his work.




Fig. 3. Hypsometric distribution of arboreal individuals sampled in Atlantic forest fragment in Goiana, Pernambuco, 2017.

As for diametric distribution, hypsometric analysis was also performed for the ten species of higher VI (figure 4). It can be seen that most species followed the pattern, with the largest number of individuals concentrating on the first classes, except for *T. guianensis* and *C. nemorosa* which had the majority of their individuals concentrated in class two. This may suggest that most individuals of these species are in a more advanced stage of succession when compared to others. Only *Eschweilera ovata*, *L. pisonis* ~~Cambess~~ and *C. mollis* presented individuals in the highest height class, demonstrating that these species have more individuals that reached maturity. Low representativeness of individuals of other species in the highest height class may indicate that they require higher nutritional requirements or luminosity, for example [17].




Fig. 4. Hypsometric distribution of the ten species with higher VI sampled in an Atlantic forest fragment in Goiana, Pernambuco, 2017.

The structural study of the forest, combining diametric and hypsometric analysis, allows to assess the stage of forest development through distribution of individuals in different strata [50]. Thus, it can be inferred that the studied fragment presents individuals with high potential of regeneration that fit, mainly, in the first classes of diameter and height, suggesting that the area is at intermediate stage of succession.

4. CONCLUSION

Regarding phytosociological analysis, it can be concluded that the family Fabaceae and Myrtaceae were the most representative for the study area. Most of the species that make up the area have zoochory as dispersion syndrome, with few anemochoric and autochoric species. As for ecological succession, the fragment showed to be predominantly composed of late secondary and early secondary species that recently reached maturity. Regarding structure, individuals were distributed mainly in the first classes of diameter and height, which indicates that the fragment has high regenerative potential. Thus, the analysis allows to infer that the studied fragment is at intermediate stage of succession.

REFERENCES

1. Mittermeier RA, Gil PR, Hoffmann M, Pilgrim J, Brooks T, Mittermeier CG, Lamoreux J, Fonseca GAB. 2005. Hotspots revisited: Earth's biologically richest and most endangered terrestrial ecoregions. Conservation International. 2005; 392.
2. Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM. The Brazilian Atlantic Forest: How much is left, and how is the remaining forest distributed? Implications for conservation. Biological Conservation. 2009; 142; 1141-1153.

3. Alves SL, Holanda AC, Wanderley JAC, Souza JS, Almeida PG. Regeneração natural em uma área de caatinga situada no município de Pombal-PB. *Revista Verde de Agroecologia e Desenvolvimento Sustentável*. 2010; 5; 152-168.
Available: <http://www.gvaa.com.br/revista/index.php/RVADS/article/view/286>
4. Marangon LC, Soares, JJ, Feliciano, ALP. Florística arbórea da mata da pedreira, município de Viçosa, Minas Gerais. *Revista Árvore*. 2003; 27; 2; 207-215.
Available: <http://www.scielo.br/pdf/%0D/rarv/v27n2/15939.pdf>
5. Silva WC, Marangon LC, Ferreira RLC, Feliciano ALP, Aparício PS, Costa Junior RF. Estrutura horizontal e vertical do componente arbóreo em fase de regeneração natural na mata de Santa Luzia, no município de Catende-PE. *Revista Árvore*. 2010; 34; 5; 863-869.
Available: <http://www.redalyc.org/html/488/48815860011/>
6. Batista APB, Marangon LC, Lima RBL, Santos RC, Baracho Júnior, E. Estrutura fitossociológica, diamétrica e hipsométrica da comunidade arbórea de um fragmento de floresta atlântica no Município de Moreno, Pernambuco, Brasil. *Revista Verde de Agroecologia e Desenvolvimento Sustentável*. 2012; 7; 5; 114-120.
Available: <http://www.gvaa.com.br/revista/index.php/RVADS/article/view/1756>
7. Cordeiro N, Howe HF. Forest fragmentation severs mutualism between seed dispersers and an endemic African tree. *Proceedings of the National Academy of Sciences of the United States of America*. 2003; 100; 24; 14052-14056.
Available: <https://doi.org/10.1073/pnas.2331023100>
8. Howe HF, Miriti MN. When seed dispersal matters. *Bioscience*. 2004; 54; 7; 651-660.
Available: [https://doi.org/10.1641/0006-3568\(2004\)054\[0651:WSDM\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2004)054[0651:WSDM]2.0.CO;2)
9. Dias CR, Umetsu F, Breier TB. Contribuição dos poleiros artificiais na dispersão de sementes e sua aplicação na restauração florestal. *Revista Ciência Florestal*. 2014; 24; 2; 501-507.
Available: <http://www.redalyc.org/articulo.oa?id=53431140023>
10. Paula A, Silva AF, Marco Júnior P, Santos FAM, Souza AL. Sucessão ecológica da vegetação arbórea em uma Floresta Estacional Semidecidual, Viçosa, MG, Brasil. *Revista Acta botanica brasiliica*. 2004; 18; 3; 407-423.
11. Rode R, Figueiredo Filho A, Galvão F, Machado SA. Estrutura horizontal da comunidade arbórea sob um povoamento com *Araucaria angustifolia* e uma Floresta Ombrófila Mista. *Pesquisa Florestal Brasileira*. 2010; 30; 64; 347-361.
Available: <https://pfb.cnpf.embrapa.br/pfb/index.php/pfb/article/view/104>
12. Rutten G, Ensslin A, Hemp A, Fischer M. Vertical and horizontal vegetation structure across natural and modified habitat types at Mount Kilimanjaro. *PloS one*. 2015; 10; 9; e0138822.
Available: <https://doi.org/10.1371/journal.pone.0138822>
13. Souza DR, Souza AL, Gama JRV, Leite HG. Emprego de análise multivariada para estratificação vertical de florestas inequidâneas. *Revista Árvore*. 2003; 27; 1; 59-63.
Available: <http://www.scielo.br/pdf/%0D/rarv/v27n1/15922.pdf>
14. Santos WB. Estrutura do componente arbóreo da borda e interior do fragment de Floresta ombrófila, Mata do Curumim, em São Lourenço da Mata – PE, Brasil: UFRPE, Universidade Federal Rural de Pernambuco, 2014. 90 p. Dissertação de Mestrado em Ciências Florestais.
Available: http://www.ppgcf.ufrpe.br/sites/www.ppgcf.ufrpe.br/files/documentos/wedson_batista_dos_santos_0.pdf
15. Silva AF, Oliveira RV, Santos NRL, Paula A. Composição florística e grupos ecológicos das espécies de um trecho de floresta semidecídua submontana da fazenda São Geraldo, Viçosa-MG. *Revista Árvore*. 2003; 27; 3; 311-319.

Available: <http://www.scielo.br/pdf/rarv/v27n3/a06v27n3>

16. Domingues CAJ, Gomes VGN, Quirino ZGM. Síndrome de dispersão na maior área de proteção da Mata Atlântica paraibana. *Revista Biotemas*. 2013; 3; 26; 99–108.
Available: <https://periodicos.ufsc.br/index.php/biotemas/article/view/26008>
17. Lambrecht FR, Dallabrida JP, Trautenmuller JW, Carli L, Burgin MRB, Fortes FO. Florística e estrutura em área de floresta estacional decidual na região do Alto Uruguais, RS. *Brazilian Journal of Biosystems Engineering*. 2016; 10; 2; 198-209.
18. Chaves ADCG, Santos RMS, Santos JO, Fernandes AA, Macarajá PB. A importância dos levantamentos florísticos e fitossociológicos para a conservação e preservação das florestas. *ACSA*. 2013; 9, 2, 43-48.
Available: <http://revistas.ufcg.edu.br/acsa/index.php/ACSA/article/view/449/pdf>
19. Köppen W. Tradução: Corrêa ACB. *Sistema Geográfico dos Climas*. 1996. 494 Notas e Comunicado de Geografia – Série B: Textos Didáticos nº13. Ed. 495 Universitária – UFPE, Departamento de Ciências Geográficas, UFPE, p.31.
20. APAC - Agência Pernambucana de Águas e Clima. 2016.
Available: <<http://www.apac.pe.gov.br/meteorologia/monitoramento-pluvio.php>>. Acesso em: 20 de April. de 2018.
21. Byng JW. et al. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society*. 2016; 181; 1–20.
Available: <http://www.bio-programmer.com/Uploads/files/2017-09-18/20170918212547j2TQ.pdf>
22. Cientec - Consultoria e Desenvolvimento de Sistemas. *Mata Nativa 4: Software para inventário de florestas nativas*. Viçosa, MG, 2016.
23. PIJL, L. van der. 1982. *Principles of dispersal in higher plants*, v. 2, 1982.
24. Gandolfi S, Leitao Filho H, Bezerra F, Carlos LF. Levantamento florístico e caráter sucessional das espécies arbustivo-arbóreas de uma floresta mesófila semidecídua no município de Guarulhos, SP. *Revista Brasileira de biologia*. 1995; 55; 4; 753-767.
Available: https://www.researchgate.net/profile/Sergius_Gandolfi/publication/259471542
25. Hasenauer H, et al. (Ed.). *Sustainable forest management: growth models for Europe*. (No. 634.92 S964). Berlin: Springer. 2006.
26. Rizzini CT. *Árvores e madeiras úteis do Brasil: manual de dendrologia brasileira*. São Paulo: Edgard Blücher; 1979.
27. Cruz AR, Silva KC, Freitas AFN. Estrutura e florística de comunidade arbórea em duas áreas de Floresta Ombrófila Densa em Macaé, RJ. *Rodriguésia*. 2013; 64; 4; 791-805.
Available: <http://dx.doi.org/10.1590/S2175-78602013000400009>
28. Silva ACC, Prata APN, Mello AA, Santos ACAS. Síndromes de dispersão de Angiospermas em uma Unidade de Conservação na Caatinga, SE, Brasil. *Hoehnea*. 2013; 40; 4; 601–609.
Available: <http://ri.ufs.br/jspui/handle/riufs/1517>
29. Silva Júnior JF, Marangon LC, Ferreira RLC, Feliciano ALP, Brandão CFL, Alves Junior FT. Fitossociologia do componente arbóreo em um remanescente de Floresta Atlântica no município do Cabo de Santo Agostinho, PE. *Revista Brasileira de Ciências Agrárias*. 2008; 3; 3; 276-282.
30. Brandão CFLS, Marangon LC, Ferreira RLC, Lins e Silva, A. C. B. Estrutura fitossociológica e classificação sucessional do componente arbóreo em um fragmento de floresta atlântica em Igarassu –Pernambuco. *Revista Brasileira de Ciências Agrárias*. 2009; 4; 1; 55–61.
Available: <http://www.redalyc.org/html/1190/119018227009/>

31. Marangon LC. Florística e fitossociologia de área de floresta estacional semidecidual visando dinâmica de espécies florestais arbóreas no município de Viçosa, São Carlos: UFscar, Universidade Federal de São Carlos 1999. 135 p. Tese doutorado.
32. Botrel RT, Oliveira Filho AD, Rodrigues LA, Curi N. Influência do solo e topografia sobre as variações da composição florística e estrutura da comunidade arbóreo-arbustiva de uma floresta estacional semidecidual em Ingaí, MG. *Revista Brasileira de Botânica*. 2002, 25, 2, 195-213.
33. Da Silva, JM. Revisão histórica da diversidade vegetal da Zona da Mata Norte de Pernambuco com ênfase no município de goiana. *Revista Espaço Acadêmico*. 2017, 16, 191, 12-26.
34. Liebsch D, Acra LA. Síndrome de dispersão de diásporos de um fragmento de floresta ombrófila mista em Tijucas do Sul, Paraná. *Revista Acadêmica*. 2007; 5; 2; 167-175.
Available: <http://dx.doi.org/10.7213/cienciaanimal.v5i2.9750>
35. Oliveira LSB, Marangon LC, Feliciano ALP, Lima AS, Cardoso MO, Silva VF. Florística, classificação sucessiona e síndromes de dispersão em um remanescente de Floresta Atlântica, Moreno-PE. *Revista Brasileira de Ciências Agrárias*. 2011; 6; 3; 502-507.
Available: <http://www.redalyc.org/html/1190/119021236019/>
36. Domingues CÂJ, Gomes, VN, Quirino ZGM. Síndromes de dispersão na maior área de proteção da Mata Atlântica paraibana. *Biotemas*. 2013; 26; 3; 99-108.
Available: <https://doi.org/10.5007/2175-7925.2013v26n3p99>
37. Tabarelli M, Peres CA. Abiotic and vertebrate seed dispersal in brazilian atlantic forest: implications for forest regeneration. *Biological Conservation*. 2002; 106; 2; 165-176.
Available: [https://doi.org/10.1016/S0006-3207\(01\)00243-9](https://doi.org/10.1016/S0006-3207(01)00243-9)
38. Franco BKS, Martins SV, Faria CL, Miranda Neto A. Estrato de regeneração natural de um trecho de floresta estacional semidecidual, Viçosa, MG. *Revista Árvore*. 2014; 38; 1; 31-40.
Available: <http://www.redalyc.org/html/488/48830661003/>
39. Negrini M, Aguiar MD, Vieira CT, Da Silva AC, Higuchi P. Dispersão, distribuição espacial e estratificação vertical da comunidade arbórea em um fragmento florestal no Planalto Catarinense. *Revista Árvore*. 2012, 36, 919-930.
Available: <http://dx.doi.org/10.1590/S0100-67622012000500014>.
40. Duarte E, Machado FD, Dos Santos GN, Farias KJ, Fernandes C, Higuchi P, Silva AC, Vieira F, Amaral RS, Aguiar V, Walter FF, Moraes B, Reis MA. Síndromes de dispersão de espécies arbóreas regenerantes em uma floresta ombrófila mista, Santa Catarina. *Revista Latinoamericana Ambiente e Saúde-rLAS*. 2016, 1, 1. Available: <http://revista.uniplac.net/ojs/index.php/rLAS/article/view/1920/1147>
41. Prado Júnior JA, de Faria LS, Vale VS, Oliveira AP, Gusson AE, Neto OCD, Schiavini I. Estrutura e caracterização sucessiona da comunidade arbórea de um remanescente de floresta estacional semidecidual, Uberlândia, MG. **Caminhos de Geografia**, 2011; 12; 39; 81-93.
Available: <http://www.seer.ufu.br/index.php/caminhosdegeografia/article/view/16329>
42. Medeiros AS, Pereira MG, Braz DM. Estrutura e Conservação de um Trecho de Floresta Estacional em Piraí, RJ. *Revista Floram*. 2016; 23; 3; 330-339.
Available: <http://www.scielo.br/pdf/floram/2016nahead/2179-8087-floram-2179-8087106214.pdf>
43. Pommerening, Arne. Evaluating structural indices by reversing forest structural analysis. *Forest Ecology and Management*. 2006; 224; 3; 266-277.
Available: doi.org/10.1016/j.foreco.2005.12.039

44. Pastorella F, Paletto A. Stand structure indices as tools to support forest management: an application in Trentino forests (Italy). *Journal of Forest Science*. 2013; 59; 4; 159-168.
Available: <https://www.agriculturejournals.cz/publicFiles/90901.pdf>
45. Callerago RM, Solon JL, Araujo ACB, Kanieski MR, Floss PA, Gracioli CR. Estrutura do componente arbóreo de uma floresta estacional decidual ripária em Jaguari, RS. *Ciência Rural*. 2012; 42; 2; 305-311.
46. Alves Júnior FT, Brandão CFLS, Rocha KD, Silva JT, Marangon LC, Ferreira RLC. Estrutura diamétrica e hipsométrica do componente arbóreo em um fragmento de mata atlântica, Recife-PE. *Cerne*. 2007; 3; 1; 83-95.
47. Almeida Junior EB, Santos Filho FS, Araújo EL, Zickel CS. Structural characterization of the woody plants in restinga of Brazil. *Journal of Ecology and the Natural Environment*. 2011. 3; 3; 95-103.
Available: http://www.academicjournals.org/article/article1379762293_Almeida%20et%20al.pdf
48. Christo AG, Guedes BRR, Sobrinho FAP, Silva AG, Peixoto AL. Structure of the shrub-arboreal componente of an Atlantic Forest fragment on a hillock in the central lowland of Rio de Janeiro, Brazil. *Interciencia*. 2009; 34; 4; 232-239.
Available: <http://www.redalyc.org/html/339/33911575002/>
49. EISFELD, R. L.; SOCHER, L. G.; ZACHOW, R. PORTES, M. C. G. O.; GARCIA, T. Caracterização florística e estrutura fitossociológica das espécies arbóreas da Floresta Nacional do Açungui, Campo Largo – PR. *Ambiência*. 2014; 10; 2; 429-448.
Available: <http://200.201.10.18/index.php/ambiencia/article/view/2391>
50. Marangon LC, Feliciano ALP, Brandão CFLS, Alves Júnior FT. Relações florísticas, estrutura diamétrica e hipsométrica de um fragmento de floresta estacional semidecidual em Viçosa (MG). *Floresta*. 2008; 38; 4; 699-709.
Available: <http://dx.doi.org/10.5380/ufv.v38i4.13166>
51. Felfili, JM. Diameter and height distributions in a gallery forest tree community and some of its main species in central Brazil over a six-year period (1985-1991). *Brazilian Journal of Botany*. 1997; 20; 2; 155-162.
Available: <http://dx.doi.org/10.1590/S0100-84041997000200006>
52. Treuhaft R, Siqueira PR. Vertical structure of vegetated land surfaces from interferometric and polarimetric radar. *Radio Science*. 2000; 35; 1; 141-177.
Available: <https://doi.org/10.1029/1999RS900108>
53. Guilherme FAG, Morellato PC, Assis MA. Horizontal and vertical tree community structure in a lowland Atlantic Rain Forest, Southeastern Brazil. *Revista Brasileira de Botânica*. 2004; 27; 4; 725-737.
Available: <http://dx.doi.org/10.1590/S0100-84042004000400012>
54. Turchetto F, Callegaro RM, Conte B, Pertuzzatti A, Griebeler AM. Estrutura de um fragmento de Floresta Estacional Decidual na região do Alto-Uruguaí, RS. *Revista Brasileira de Ciências Agrárias*. 2015; 10; 2; 280-285.
Available: <http://www.redalyc.org/pdf/1190/119039562018.pdf>