

Study of of nitrogen efficiency using three corn cultivars of green ears production

Cassiana Felipe de Sousa¹, Leilson Costa Grangeiro^{1*}, Valdivia de Fátima Lima de Sousa¹, Jader Vieira Cordeiro², Jorge Luis Alves Silva¹ and Lucas Pereira Gomes¹

¹Federal Rural University of the Semi-arid Region (UFERSA), Mossoró, RN, Brazil

²Federal University Ceará (UFC), Fortaleza, CE, Brazil.

ABSTRACT

The objective of this study was to evaluate the efficiency of nitrogen utilization by corn cultivars for the production of green ears. The experiment was conducted at garden of the Department of Plant Science of Federal Rural University of the Semi-arid, Mossoró, Rio Grande Norte state, Brazil, from February to May 2015. Treatments were constituted by the combination of two nitrogen doses (0 and 90 kg ha⁻¹ N) and three corn cultivars (Cruzeta, Truck and AG 1051). The experimental design was the randomized block, in a 2 x 3 factor schem, with five replications. Cultivar AG 1051, based on the values of the nutritional efficiency indices, was more effective in using nitrogen to produce of green ears. The dose of 90 kg ha⁻¹ N provided higher yield of green corn.

Keywords: Zea mays; productivity; efficiency indices; mineral nutrition.

1. INTRODUCTION

Corn (*Zea mays* L.) is the most cultivated cereal in the world, with the largest producers being the United States, China and Brazil. It has great economic importance, due to its high production potential and the various forms of use in human and animal food, *in natura* and in the high-tech industry. It also contributes to social development, due to its capacity to generate employment and income [1]. It is commercialized mainly in the form of grains; however, in the northeastern region of Brazil, due to higher financial returns, the production of green ears has become a promising activity, being cultivated throughout the year [2].

Nitrogen fertilization plays a very important role; it is one of the most influent factors on corn crop productivity. On the other hand, for the production of green ears, N has been responsible for a yield increase; its effect was also influenced by other factors, such as genotypic and environmental ones [3].

The identification of genotypes that are able to absorb and use nitrogen effectively is a way to increase the efficiency of using nitrogen fertilizer in corn cultivations, to increase production, minimize losses and reduce environmental contamination [4]. In this context, the importance of a genetic improvement is evident, aiming not only at genotypes that are responsive to nitrogen fertilization, but also at the study of effective genotypes in the use of nitrogen [5].

The objective of this study is to evaluate the efficiency of nitrogen utilization by corn cultivars for the production of green ears.

2. MATERIAL AND METHODS

2.1 Site location and characterization

The experiment was conducted at a garden in the Plant Science department of Federal Rural University Semi-arid, Mossoró, Rio Grande do Norte State, in Brazil, (5° 11" southern latitude and 37° 20" western longitude, with 18 m altitude), from February to May 2015. The soil of the experimental area is classified as Eutroferic Yellow - Red Argisol [6]. The chemical analysis conducted on soil samples from the experimental area, at a depth between 0 and 20 cm, presented the following results: pH in H₂O (1:2.5) = 6.5; P_(mehlich⁻¹) = 210.83 mg dm⁻³; K = 5.26 mmol_c dm⁻³; Ca = 30.90 mmol_c dm⁻³; Mg = 12.60 mmol_c dm⁻³; Al = 1.0 mmol_cdm⁻³; H + Al = 10.0 mmol_c dm⁻³ and Organic matter = 12.8 g dm⁻³.

2.2 Treatments and experimental design

Treatments were constituted by the combination of two nitrogen doses (0 and 90 kg ha⁻¹) and three corn cultivars (Cruzeta, Truck and AG 1051). The used experimental design was the randomized block, in a 2 x 3 factor scheme, with five replications. Each experimental plot consisted of three rows measuring 3 m in length, spaced 0.8 m apart, containing 10 plants in each row; the central row was considered as the usable area, discarding one plant at each end.

Nitrogen doses were used to evaluate the efficiency of using N in corn cultivars, where the dose "0" corresponded to the N content in the soil (D0) and the 90 kg ha⁻¹ N (D90), recommended dose for corn production [7]. The corn cultivars the ones that are commonly used in the region, Cruzeta (Open pollination), Truck (modified triple hybrid) and AG 1051 (double hybrid).

2.3 Field establishment

Soil preparation was performed by ploughing and harrowing, followed by opening furrows with an approximate 0.20 m depth for planting fertilization, based on soil analysis and recommendation for corn [7], using: 30 kg ha⁻¹ of N (treatment with N); 40 kg ha⁻¹ of P₂O₅, 40 kg ha⁻¹ of K₂O; 1.0 kg ha⁻¹ of B and 3.0 kg ha⁻¹ of Zn. In covering, in the N treatment, 30 kg ha⁻¹ of N were applied on 15 day after sowing (DAS) and 30 kg ha⁻¹ of N were applied when the plant presented eight definitive leaves. The used sources were urea, triple superphosphate, potassium chloride, boric acid and zinc sulfate.

Sowing was performed with three seeds per hole, at a spacing of 0.8 x 0.3 m and, when the plants had four definitive leaves (9 DAS), thinning was performed, leaving one plant per hole. The irrigation was by dripping, with tips spaced of 0.3 m apart and a flow of 1.4 L h⁻¹. Weed control, when necessary, was performed by manual weeding. Four sprayings were performed using the insecticides Decis® (deltamethrin) and Premio® (chlorantraniliprole) at doses of 200 and 100 mL ha⁻¹, respectively, for control of carpiaceous caterpillars (*Spodoptera frugiperda*).

2.4 Harvesting and parameters evaluated

Before harvesting (68 DAS), on all plants from the usable area of the plot, the height of plants, the insertion of the first ear and the stalk diameter were measured.

94 The harvesting of green ears was performed manually at the phenological stage R3 (milky
 95 grains), on day 70 DAS; on the occasion, the number and the productivity of marketable
 96 green ears that were not husked (damage-free ears, well wrapped, ≥ 22 cm long) and
 97 husked (ears with good grains and ≥ 18 cm long). Two plants per plot were cut on the stem
 98 region, separated in leaves, stalk and ears, washed and dried in a forced air circulation oven
 99 (65 °C) until reaching constant mass, in order to obtain the dry matter of the plant.
 100 Subsequently, content of N in said parts was then determined. The analysis of N was carried
 101 out in the extract obtained by sulfuric digestion (N) and quantified by the semi-micro method
 102 of Kjeldahl [8].

103
 104 Nitrogen accumulation values were obtained through the product between N content and dry
 105 matter. With the data of dry matter and N accumulation, the following indices were
 106 calculated [9]: Agronomic efficiency (AE); Physiological efficiency (PE); Ear production
 107 efficiency (CPE); Recovery efficiency (RE) and Use efficiency (UE), using the following
 108 equations:

109
 110
$$AE \text{ (kg kg}^{-1}\text{)} = (PV_{cN} - PV_{sN}) / (QN_a) \quad (1)$$

111
$$PE \text{ (kg kg}^{-1}\text{)} = (MST_{cN} - MST_{sN}) / (AN_{cN} - AN_{sN}) \quad (2)$$

112
$$CPE \text{ (kg kg}^{-1}\text{)} = (PV_{cN} - PV_{sN}) / (AN_{cN} - AN_{sN}) \quad (3)$$

113
$$RE \text{ (kg kg}^{-1}\text{)} = (AN_{cN} - AN_{sN}) / (QN_a) \quad (4)$$

114
$$UEN \text{ (kg kg}^{-1}\text{)} = PE \times RE \quad (5)$$

115 where: PV_{cN} is the production of green ears with nitrogen fertilization (kg); PV_{sN} is the
 116 production of green ears without nitrogen fertilization (kg); QN_a is the quantity of applied N
 117 (kg); MST_{cN} is the total production of dry matter with nitrogen fertilization (kg); MST_{sN} is the
 118 total production of dry matter without nitrogen fertilization (kg); AN_{cN} is the total N
 119 accumulation with nitrogen fertilization (kg) and AN_{sN} is the total N accumulation without
 120 nitrogen fertilization (kg).

121

122 2.5 Statistical Analysis

123

124 The obtained data were submitted to analysis of variance and the averages compared by
 125 Tukey's test at 5% probability, using the statistic software SISVAR [10].

126

127

128 3. RESULTS AND DISCUSSION

129

130 There was a significant effect in the cultivar and N dose interaction as for plant height and
 131 insertion of the first ear. Stem diameter, plant dry matter, total ear number, number of
 132 marketable-husked ears and productivity of marketable-husked ears were significant to
 133 isolated factors (cultivar and dose). As for prolificacy, the factor cultivar was the only
 134 significant one, and for the total ear productivity and the marketable non-husked ears, the N
 135 dose factor was significant.

136

137 3.1 Plant height, insertion of the first ear, stalk diameter and dry matter

138

139 Except for the cultivar Truck, the application of nitrogen increased height plant and insertion
 140 of the first ear. In the absence of N, cultivars did not differ significantly as for these
 141 characteristics. With the application of N (D_{90}), 'Cruzeta' had the highest plant and ear
 142 insertion, even if it did not differ from AG 1051 as for plant height (Table 1). In this cultivar,

143 increases in plant height and ear insertion with the application of N were 18 and 24%
 144 respectively, compared to the treatment with no application.

145

146 **Table 1. Plant height and insertion of the first ear, in corn cultivars, without the**
 147 **application of N (D₀) and with the application of 90 kg ha⁻¹ of N (D₉₀).**

148

Cultivars	Plant height (cm)		Insertion of ear (cm)	
	D ₀	D ₉₀	D ₀	D ₉₀
Cruzeta	209.0 aB	247.0 aA	110.0 aB	137.0 aA
Truck	205.0 aA	212.0 bA	99.0 aA	103.0 cA
AG 1051	204.0 aB	226.0 aA	106.0 aB	122.0 bA
CV (%)	5.86		7.91	

149

Means followed by the same lowercase letter in the column and upper case in the line, within each variable do not differ by Tukey test, 5% probability.

150

151

152 The increase in plant height and ear insertion, caused by providing N, is mainly due to the
 153 role that N plays in the vegetative development, influencing the height of plants and the
 154 accumulation of phytomass, since higher plants tend to provide a higher ear insertion [11].
 155 This characteristic of higher ear insertion is desirable from the operational point of view,
 156 since it provides significant advantages during harvesting.

157

158 Cultivars differed as for stem diameter, plant dry matter and prolificacy (Table 2). As for stem
 159 diameter, AG 1051 exceeded the others, which did not differ from one another. On the other
 160 hand, Cruzeta and Truck presented greater plant dry matter and prolificacy. The increase in
 161 the stem diameter is an important physiological factor, since the stem does not only have a
 162 leaf and inflorescence support function, but it mainly works as a structure for the storage of
 163 soluble solids that are later used during grain formation [12]. The application of N provided
 164 greater stem diameter and plant dry matter. As for prolificacy, there was no significant
 165 difference.

166

167 **Table 2. Stem diameter, plant dry matter and prolificacy according to corn cultivar and**
 168 **N dose.**

169

Cultivars	Stem diameter (mm)	Dry matter (g planta ⁻¹)	Prolificacy
Cruzeta	21.85 b	159.86 a	1.05 ab
Truck	20.98 b	140.80 ab	1.20 a
AG 1051	24.38 a	134.41 b	1.03 b
Dose N (Kg ha ⁻¹)			
0	20.67 b	126.25 b	1.07 a
90	24.21 a	163.38 a	1.12 a
CV(%)	7.69	12.66	13.35

170

Means followed by the same lowercase letter in the column, within each variable do not differ by Tukey test, 5% probability.

171

172

173 3.2 Number and the productivity of green ears

174 The cultivar Truck exceeded the others in terms of ear number (husked, non-husked and
 175 total). However, as for productivity, the cultivars differed significantly only in the production of

marketable husked ears (PMHE); Truck was superior but it did not differed from AG 1051 (Table 3). The percentage of non-husked marketable ears in relation to the total ear number was high in all cultivars; it was 94, 95 and 97% respectively for Cruzeta, Truck and AG 1051.

In spite of the superiority of Truck as for number of marketable non-husked ears (NME), it did not differ from the others in the production of marketable non-husked ears (PME), demonstrating that, even if greater in number, the ear matter was lower. The PMHE was lower in relation to the PME, in all cultivars. The reduction was 60, 42 and 43% respectively for Cruzeta, Truck and AG 1051. This difference occurs since not all non-husked ears have the pattern required by the market for husked ears (ears with good grains and ≥ 18 cm long).

Table 3. Total ear number (TEN), number of marketable non-husked ears (NME) and number of marketable husked ears (NMHE), total ear productivity (TEP), productivity of marketable non-husked ears (PME) and productivity of marketable husked ears (PMHE), according to corn cultivar and N dose.

Cultivars	TEN -----ear ha ⁻¹ -----	NME -----ear ha ⁻¹ -----	NMHE -----ear ha ⁻¹ -----	TEP -----kg ha ⁻¹ -----	PME -----kg ha ⁻¹ -----	PMHE -----kg ha ⁻¹ -----
Cruzeta	33984,65 b	2031,51 b	18652,49 c	10031,72 a	10008,61 a	4026,07 b
Truck	43055,90 a	1146,17 a	39062,81 a	10707,90 a	10209,20 a	5958,38 a
AG 1051	33333,60 b	2291,93 b	27778,00 b	9798,52 a	9521,95 a	5457,16 ab
Dose N (kg ha ⁻¹)						
0	34635,70 b	32465,54 b	24356,39 b	8791,30 b	8500,55 b	3999,22 b
90	38947,07 a	37847,53 a	32639,15 a	11567,45 a	11325,96 a	6295,19 a
CV(%)	12,69	15,26	20,71	17,78	18,25	25,74

Means followed by the same lowercase letter in the column, within each variable do not differ by Tukey test, 5% probability.

In green ears production system, ears are transported for benefit or consumption while non-husked, in order to avoid the degradation of sugars, the denaturing of the product and in order to keep the sweet taste. Thus, productivity with husk is an important characteristic for the commercialization of corn green [13].

The application of N, as expected, helped obtaining a greater number of ears and greater productivity (Table 3). This increase is due to the fact that N causes greater effects in the characteristics about plant growth and development, which affect productivity directly or indirectly [14].

3.3 Nutritional efficiency indices

Among the evaluated efficiency indices, there was a significant difference among the cultivars, as for agronomic efficiency and recovery efficiency (Table 4). AG 1051 was more effective in producing ears (AE) and in the quantity of accumulated N (RE), per unit of applied N, not differing significantly from the cultivar Truck as for RE.

Table 4. Agronomic efficiency (AE), physiological efficiency (PE), ear production efficiency (EPE), recovery efficiency (RE), N use efficiency (NUE), of corn cultivars for the production of green ears.

Cultivars	AE	PE	EPE	RE	NUE
	-----kg kg ⁻¹ -----				

* Tel.: +55 84 3312 6222.

E-mail address: leilson@ufersa.edu.br.

Cruzeta	27,08 b	54,18 a	134,06 a	0,17 b	7,98 a
Truck	19,42 b	42,38 a	56,97 a	0,32 ab	13,74 a
AG 1051	57,67 a	41,97 a	171,12 a	0,36 a	14,04 a
CV(%)	38,84	43,28	64,96	35,61	56,45

Means followed by the same lowercase letter in the column, within each variable do not differ by Tukey test, 5% probability.

The non-observation of a significant difference in the indices of physiological efficiency (PE), ear production efficiency (EPE) and N usage efficiency (NUE) is probably due to the high variation coefficient. However, AG 1051 was numerically superior in CPE than the others. This indicates that it was also more effective in the production of ears per unit of accumulated N, with a consequent higher NUE. On the other hand, Cruzeta was more effective in the biological production, that is, it produced a higher quantity of dry matter per applied N unit.

4. CONCLUSIONS

Based on the values of the nutritional efficiency indices, cultivar AG 1051 was more effective in using nitrogen to produce green ears corn. The dose of 90 kg ha⁻¹ N provided higher yield of green corn.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration between all authors. This work was carried out in collaboration between all authors. Author CFS participated in the idea and management of the experiment, besides writing the article. Author LCG participated in the elaboration of the research project, conduction of experiments and writing of the article. Author VFLS was responsible for collecting, tabulating and analyzing the data. Authors JVC, JLAS and LPG participated in the management of the experiment from the implantation to the data collection. All authors read and approved the final manuscript.

REFERENCES

1. Silva GR, Oliveira FHT, Pereira RG, Silva PSL, Diógenes TBA, Silva ARC. Doses de nitrogênio e fósforo para produção econômica de milho na Chapada do Apodi, RN. Rev Bras Eng Agríc Ambient. 2014; 18(12):1247-1254.
2. Rocha DR, Fornasier Filho D, Barbosa JC. Efeitos da densidade de plantas no rendimento comercial de espigas verdes de cultivares de milho. Hortic Bras 2011; 29(3): 392-397.
3. Freire FM, Viana MCM, Mascarenhas MHT, Pedrosa MW, Coelho AM, Andrade CLT. Produtividade econômica e componentes da produção de espigas verdes de milho em função da adubação nitrogenada. Rev Bras Milho Sorgo. 2010; 9(3):213-222.
4. Fidelis RR, Miranda GV, Santos IC, Galvão JCC, Peluzio JM, Lima SO. Fontes de germoplasma de milho para estresse de baixo nitrogênio. Pesq Agropec Trop. 2007; 37(3):147-153.

* Tel.: +55 84 3312 6222.

E-mail address: leilson@ufersa.edu.br.

- 261 5. Carvalho RP, Von Pinho RG, Davide LMC. Desempenho de cultivares de milho quanto
262 à eficiência de utilização de nitrogênio. Rev Bras Milho Sorgo. 2011; 10(2):108-120.
- 263 6. Santos, HG et al. Sistema brasileiro de classificação de solos. 3. ed. rev. ampl. Brasília:
264 Embrapa, 2013
- 265 7. Cavalcanti FJA. Recomendações de adubação para o Estado de Pernambuco: 2ª
266 aproximação. 3 ed. IPA: Recife, 2008.
- 267 8. Malavolta, E et al. Avaliação do estado nutricional das plantas: princípios e aplicações.
268 2. ed. Piracicaba: Associação Brasileira para Pesquisa da Potassa e do Fosfato, 1997.
- 269 9. Fageria NK, Santos AB, Cutrim VA. Produtividade de arroz irrigado e eficiência de uso
270 do nitrogênio influenciadas pela fertilização nitrogenada. Pesq. Agropec. Bras. 2007;
271 42(7):1029-1034.
- 272 10. Ferreira DF. Sisvar: a computer statistical analysis system. Ciênc Agrotec.
273 2011;35(6):1039-42.
- 274 11. Silva EC, Buzetti S, Guimarães GL, Lazarini E, Sá ME. Doses e épocas de aplicação
275 de nitrogênio na cultura do milho em plantio direto sobre Latossolo Vermelho. Rev Bras
276 Ciênc Solo. 2005; 29(3): 353-362.
- 277 12. Carmo MS, Cruz SCS, Souza EJ, Campos LFC, Machado CG. Doses e fontes de
278 nitrogênio no desenvolvimento e produtividade da cultura de milho doce (*Zea mays*
279 *convar. saccharata* var. *rugosa*). Biosci. J.v. 2012;28(1): 223-231.
- 280 13. Oliveira Júnior LFG, Deliza R, Pereira MG, Chiquieri TB, Bressan-Smith, R. Seleção de
281 genótipos de milho mais promissores para o consumo in natura. Ciênc Tecnol Aliment.
282 2006; 26(1):159-166.
- 283 14. Okumura RS, Mariano DC, Zaccheo PVC. Uso de fertilizante nitrogenado na cultura do
284 milho: uma revisão. Rev Bras Tecnol Aplicada Ciênc Agrárias. 2011; 4(2): 226–244.
- 285
- 286
- 287
- 288