

Original Research Article

Genetic diversity among accessions of *Calotropis procera* based on agronomic characters

ABSTRACT

This study evaluated the genetic diversity among accessions of *Calotropis procera* based on agronomic characters. Seeds of 70 *C. procera* accessions were collected in the Northeast region of Brazil. The experiment was conducted in a greenhouse in Campina Grande, Paraíba State, between January and September 2016. The plants were cultivated for 240 days after sowing in plastic pots filled with soil. Data regarding 23 characteristics were obtained; the following analyses were conducted: analysis of variance by the F test; canonical variables and cluster by the Tocher method and by the unweighted-pair-group method with arithmetic mean (UPGMA). The experiment indicated that genetic variability exists among the 70 *C. procera* accessions. Accessions 1, 12, 20, 39, 48, 50, 57, 67 and 70 are promising for use in the *C. procera* breeding program, with agronomic and forage potential, and can be used as parents in hybridization programs as well as for future studies.

Keywords: Genetic improvement, xerophilous plant, genetic variability, cluster analysis, *Calotropis procera*

1. INTRODUCTION

Calotropis procera [Ait.] WT Aiton is a species native to Africa, Madagascar, the Arabian Peninsula, Southwest Asia and India as well as from China to Malaysia (Rahman and Wilcock 1991). The species is a perennial, xerophilous plant that grows in arid to semiarid regions and stands out because of its drought resistance and salinity tolerance (Hassan et al., 2015). The economic applications of the species include its use as a medicinal plant due to its phytochemical and pharmacological properties (Ranjit et al., 2012) as well as in the production of bioenergy and biofuels in semiarid regions (Rathore and Menna 2010; Barbosa et al., 2014).

In this context, due to the economic potential and the need for new studies with the species, research has recently been conducted on its phenology (Farahat et al., 2015), seed storage (Oliveira-Bento et al., 2015), oil and fatty acid contents (Barbosa et al., 2014), metabolomic response to changes in water availability (Ramadan et al., 2014) and genetic diversity (Almeida et al., 2017b; El-Bakry et al., 2014; Yao et al., 2015), along with a literature review on the species (Hassan et al., 2015).

In Brazil, research aimed at forage production has been conducted, due to the potential of this species for use as ruminant feed. Further studies are needed on the rational cultivation of the crop, especially genetic improvement studies (Costa et al., 2009), to evaluate the genetic divergence between various materials.

Genetic divergence is one of the most important parameters evaluated by plant breeders during the initial phase of a breeding program (Negreiros et al., 2008), with the purpose of evaluating the genetic variability in the species to select promising materials. According to Cruz (2006), the success of a plant breeding program is directly related to the existence of variability in the base population, with superior and divergent individuals recommended for crossbreeding.

Researchers have conducted genetic diversity studies with *C. procera* using molecular markers (El-Bakry et al., 2014; Yao et al., 2015) and observed the occurrence of genetic variability. Nevertheless, Yao et al. (2015) indicate that is known regarding the genetic diversity in *C. procera*; thus, new studies are necessary to increase knowledge concerning this diversity.

Although the volume of genetic information from molecular markers has increased significantly in genetic diversity studies, emphasis is still placed on the study of diversity through phenotypic characteristics, mainly of a quantitative nature, due to their economic importance in plant breeding programs (Cruz et al., 2011). Almeida et al. (2017a) started a breeding program for *C. procera*, with the early selection of seedlings from estimates of genetic parameters. Recently, Almeida et al. (2017b) performed a study of genetic diversity based on seed physiological quality.

However, the literature contains no genetic diversity studies of the species using phenotypic characters, and research is needed to identify divergent accessions with potential for forage production. Thus, preliminary research involving the germplasm occurring naturally in the Brazilian territory becomes an important step in the *C. procera* breeding program. Therefore, considering the economic potential of the species, especially its forage potential, the present study aimed to evaluate the genetic diversity among *C. procera* accessions based on agronomic traits to support the selection of genotypes with forage potential.

2. MATERIAL AND METHODS

Seeds from 70 accessions of *C. procera* were collected between October and January 2016 in the Northeast region of Brazil within areas where the species occurs naturally, as shown in Figure 1. Currently, these accessions are included in the germplasm collection of the Instituto Nacional do Semiárido (National Institute for the Semi-arid region, INSA). The experiment was installed and conducted in a greenhouse at INSA in Campina Grande, Paraíba State, Brazil (7°16'23.25"S, 35°58'17.06"W and an elevation of 531 m), between January and September 2016.




Figure 1. Geographic location of the seed collection sites for the 70 *C. procera* accessions in the Northeast region of Brazil.

Physiologically mature *C. procera* fruit with open capsules and brown seeds were collected. The seeds were extracted manually and dried in the shade. After drying, the seeds were stored in paper bags at room temperature until sowing. Next, 20 of the seeds were sown per pot, and after an evaluation of emergence, thinning was performed, leaving only five seedlings per pot. Subsequently, the plants were collected for weighing throughout the experiment, leaving only one plant per pot until 240 days after sowing (DAS).

The plants were grown for 240 DAS in plastic 10-liter pots with the following dimensions: 25 cm in height, 27 cm in diameter at the opening and 18 cm in diameter at the base. A drainage system was installed at the base of the pots, with four holes to facilitate water drainage. The pots were filled with sandy soil of the following chemical composition: pH 4.9, 2.55 mg/dm³ of P, 71.6 mg/dm³ of K⁺, 0.09 cmolc/dm³ of Na⁺, 1.82 cmolc/dm³ of H⁺ + Al³⁺, 0.4 cmolc/dm³ of Al³⁺, 0.34 cmolc/dm³ of Ca²⁺, 0.8 cmolc/dm³ of Mg²⁺, 1.41 cmolc/dm³ for the sum of exchangeable bases, 3.22 cmolc/dm³ of cation exchange capacity and 1.79 g/kg of organic matter. The soil physical composition consisted of 900 g/kg of sand (0.05-2 mm), 60 g/kg of silt (0.002-0.05 mm), 40 g/kg of clay (<0.002 mm) and sand, as textural

class. Soil fertilization was performed as recommended by the chemical and physical analysis, and the plants were irrigated at 3 day intervals, based on water potential and soil field capacity.

A randomized block experimental design was used, with 70 accessions in three replicates. The number of seedlings emerged between the sixth and the tenth DAS were counted, and the following characteristics were evaluated: the final emergence percentage (FEP) (Brazil, 2009); speed of emergence index (SEI) (Maguire, 1962); and seedling vigor index (SVI) (Maroufi, Farahani and Moaveni 2011). The plant height (HEI), stem diameter (SD), number of leaves (NL), average leaf length (ALL), average leaf width (ALW), leaf area (LA) and total leaf area (TLA) at 210 DAS were also measured, as were the stem fresh mass (SFM), leaf fresh mass (LFM), total fresh mass (TFM), leaf dry mass (LDM), stem dry mass (SDM) and total dry mass (TDM) at 120 DAS. The fresh mass and dry mass data were obtained in this evaluation period, when two plants were present per pot, after which only one plant remained in the pot for the analysis of the next phenological stages, that is, of the characteristics described previously, as well as the beginning of flowering (BF), the number of flowers per inflorescence (NFI) and the number of fruit per inflorescence (NFrI).

The HEI was obtained by measuring the distance from the root crown to the apical bud with the aid of a graduated ruler (in centimeters). For the measurements of the SD (measured at the stem base, corresponding to the root crown of the plants), a digital caliper (in millimeters) was used. For the NL count, only fully expanded leaves were considered, and immature leaves were not counted; length (cm) and width (cm) were measured on three leaves obtained in the lower, middle and upper parts of the plant to generate the ALL and ALW values. The individual LA was obtained from the equation $LA (cm^2) = W \times L \times 0.75$, proposed by Moreira Filho et al. (2007) for *C. procera*; and the TLA was obtained by summing the individual LAs over all the leaves.

For the phytomass analysis, the plants were divided into leaves and stems for the determination of LFM, SFM and TFM on a precision scale ($e = 0.0001$ g). The material was then allowed to dry in a forced-air oven at 65 °C for 48 hours until reaching a constant mass, at which point the LDM, SDM and TDM were obtained, using the same scale.

In addition, chlorophyll a and b fluorescence was measured on base leaves (CAB and CBB) and on apex leaves (CAA and CBA) at 210 DAS, using a chlorophyll meter (Clorofilog Falker CFL 1030).

The data on the agronomic characteristics were subjected to analysis of variance (ANOVA) via the F test ($p < 0.05$). The genetic parameters were estimated based on the expected mean squares of the ANOVA. In addition, canonical variables analysis and cluster analysis by the Tocher method and by the unweighted-pair-group method with arithmetic mean (UPGMA) were performed with the GENES software (Cruz, 2013).

3. RESULTS AND DISCUSSION

The analysis of variance and estimates of the genetic parameters for the 23 agronomic characters are described in Table 1. The F test showed a significant difference ($P = 0.01$) among the 70 *C. procera* accessions for all the evaluated characters. Thus, the genetic parameters were estimated to support the selection of promising materials. According to Bárbaro et al. (2004), estimates of genetic parameters assume an important predictive role for directing breeding programs during the process of selecting the most promising genotypes.

Table 1. Summary of analyses of variance and estimates of genetic parameters for the 23 agronomic characters of the 70 *C. procera* accessions.

SV	DF	Average squares							
		SEI	FEP	SVI	CAA	CBA	CAB	CBB	SD
Blocks	2	39.05	124.64	0.351	11.32	6.63	0.052	0.252	16.28
Accessions	69	28.93**	677.4**	0.046**	9.27**	6.82**	16.96**	3.37**	9.26**

Residue	138	5.27	49.16	0.025	2.30	1.95	5.65	1.18	3.96
Genetic parameters									
Average		14.73	92.92	0.37	38.74	10.69	32.97	6.75	27.41
CVe (%)		15.57	7.54	41.39	3.91	13.07	7.21	16.06	7.26
h ²		81.78	92.74	45.95	75.15	71.32	66.67	65.07	57.17
CVg (%)		19.05	15.57	22.03	3.93	11.9	5.88	12.65	4.84
CVg/CVe		1.22	2.06	0.53	1.00	0.91	0.81	0.78	0.66
Average squares									
SV	DF	HEI	NL	ALW	ALL	LA	TLA	BF	NFI
Blocks	2	615.24	8.93	1.96	3.74	1284.4	1124212.4	6104.16	1515.1
Accessions	69	354.57**	7.91**	1.26**	1.68**	583.6**	151842.4**	948.4**	132.0**
Residue	138	105.19	2.06	0.27	0.6	135.36	70671.51	203.35	24.1
Genetic parameters									
Average		113.68	19.87	8.4	17.54	113.3	1678.8	168	8.73
CVe (%)		9.02	7.22	6.19	4.41	10.27	15.84	86.95	56.25
h ²		70.33	73.96	78.47	64.5	76.81	52.73	78.56	81.75
CVg (%)		8.02	7.03	6.82	3.43	10.79	9.73	96.09	68.73
CVg/CVe		0.89	0.97	1.1	0.78	1.05	0.61	1.11	1.22
Average squares									
SV	DF	NFrl	SFM	LFM	TFM	SDM	LDM	TDM	
Blocks	2	28.6	190.99	386.38	1115.52	9.9	1.71	19.26	
Accessions	69	3.50**	381.68**	685.93**	1866.43**	32.45**	14.36**	79.57**	
Residue	138	0.56	19.43	39.29	25.42	2.39	1.17	5.85	
Genetic parameters									
Average		1.25	35.68	70.56	106.24	9.03	8.15	17.17	
CVe (%)		59.56	12.36	8.88	4.75	17.12	13.28	14.08	
h ²		84.11	94.91	94.27	98.64	92.64	91.85	92.65	
CVg (%)		79.12	30.8	20.81	23.32	35.08	25.74	28.87	
CVg/CVe		1.33	2.49	2.34	4.91	2.05	1.94	2.05	

SV - source of variation; DF - degrees of freedom; CVe - coefficient of environmental variation; h² - heritability; CVg - coefficient of genetic variation; **significant (p < 0.01) by the F test.

As a result, genetic variability was observed in this study, as the genetic parameter estimates demonstrate a genetic effect on the expression of the characters, with the CVe values indicative of good experimental precision. Additionally, the CVg values for most of the characters were high, higher than their CVe values, thus leading to a CVg/CVe ratio above 1—except for the characters SVI, CBA, CAB, CBB, SD, HEI, NL, ALL and TLA—a situation favorable to genetic gains in the selection of promising materials, according to Vencovsky (1987).

However, these parameters should not be considered in isolation, with heritability being a better indicator of the success of a selection process in breeding programs. Because the h² estimates were high (Cruz, 2005), the possibility exists for the selection of promising accessions, with potentially significant genetic gains during the selection process. Almeida et al. (2017b) also observed genetic variability in *C. procera* genotypes, which supported an early selection of more productive seedlings. However, the study of genetic diversity must account for information from all the phenological stages of the species, as a way of aiding the selection process.

Yao et al. (2015) reported that little is known regarding genetic diversity in *C. procera*, and Barbosa et al. (2014) added that no existing genetic studies investigate the occurrence

of different genotypes in the Brazilian territory. Therefore, knowing that genetic variability was present among the 70 *C. procera* accessions, a study of genetic diversity was conducted through multivariate analysis, canonical variables analysis, and the measurement of dissimilarity between accessions as a way of generating a cluster structure and identifying similar and divergent materials to select parents for hybridization programs (Cruz et al., 2011).

Furthermore, Santos et al. (2000) reported that studies involving the analysis of genetic diversity, based on multivariate analysis techniques, have offered effective contributions in the discrimination and indication of potential parents for use in breeding programs, besides providing a greater knowledge of the accessions in germplasm collections.

In addition, the study of genetic diversity based on agronomic characters helps to distinguish the accessions, promoting the conservation of the genetic resources of the species; this protection occurs because the materials are conserved after their characterization in the germplasm bank as a way to preserve the genetic variability of the species (Costa et al., 2012; Ramos et al., 2007).

Thus, a cluster analysis of the 70 *C. procera* accessions by the Tocher optimization method was performed, based on Mahalanobis' generalized distance (D^2). This analysis generated five distinct groups (Table 2), with three groups consisting of only one accession (groups 3, 4 and 5) and considered the most isolated among those evaluated accessions. The first group contained 88.57% of the accessions and was subdivided into three subgroups: subgroups 1, 2 and 3 formed by 25.72, 41.42 and 21.42% of the accessions, respectively.

Table 2. Clustering of the 70 *C. procera* accessions by the Tocher optimization method.

Group		Accessions																	
	I.1	25	64	4	15	7	63	43	33	5	55	16	34	54	2	26	9	37	35
I	I.2	62	56	68	38	24	17	66	58	32	45	11	51	18	69	59	3	42	27
		41	44	65	61	49	46	40	14	8	52	36							
	I.3	19	29	10	31	6	13	30	23	60	47	22	57	67	12	21			
II	1	50	39	70	48														
III	20																		
IV	53																		
V	28																		

During the collection of the seeds, a greater genetic divergence was assumed to occur with a greater geographic distance between the accessions. However, the results indicate that the clustering pattern was unrelated to the geographic origin of the materials, since accessions from distant regions showed relatively small genetic distances. Thus, future studies with molecular markers are needed to explain the observed clustering pattern.

The second group was formed by accessions 1, 50, 39, 70 and 48, which are differentiated by mean values above the overall means for all the characters and thus constitute materials with crop potential. These accessions are promising for species breeding programs and can be used in hybridization programs, considering their production potential and genetic distance from the other groups. Cruz (2006) noted that the success of a plant breeding program is directly related to the existence of variability in the base population, with superior and divergent individuals recommended for crossbreeding.

Group 3, composed of accession 20, also obtained mean values above the overall mean for all the characters, together with early flowering and fruiting, showing promise as a material for seed production. This group can also be used as a parent in hybridization programs, given its genetic complementarity.

The fourth group, consisting of accession 53, stands out for its high phytomass production but reached a mean value below the overall mean for the SDM, LDM and TDM characters. Therefore, despite its genetic value, this genotype has no importance for the species breeding program because, according to Silva and Queiroz (2002), dry matter is an important characteristic in forage plants, composing the feed portion that contains all the nutrients and corresponding to the total mass minus the moisture.

By contrast, group 5, comprising accession 28, represents the material with the greatest genetic divergence from the accessions of group I, with lower results obtained for all the examined characters. **Despite the lower performance, access 28 should be preserved in the germplasm bank as a genetic resource.**

In addition, cluster analysis via the UPGMA hierarchical method was performed to detail the dissimilarity between the accessions through a dendrogram (Figure 2).




Figure 2. UPGMA dendrogram obtained from the dissimilarity matrix (D^2) of the 70 *C. procera* accessions based on 23 agronomic characters. The cophenetic correlation coefficient was 0.777. The dotted line represents the cutoff point based on the methodology proposed by MOJENA (1977).

The UPGMA analysis was able to significantly represent the genetic diversity between the accessions, with a satisfactory cophenetic correlation coefficient (CCC) (0.7777). This coefficient indicates the degree of distortion regarding the representation of similarity between individuals in a dendrogram, where values close to unity represent low distortion.

According to Rohlf (1970), a CCC below 0.7 indicates inadequacy of the cluster method, that is, the actual genetic divergence is not reliably represented in the two-dimensional plane.

A cutoff point at 2.1268 for dissimilarity by the Mojena criterion (Mojena, 1977) corresponds to a total dissimilarity of approximately 35%, which allowed the formation of six groups, with group 1 subdivided into four subgroups. The results observed by the Tocher method were similar to those obtained by the UPGMA analysis, differing mainly in the clustering of accessions 60, 22, 57, 67, 12, 48, 21, 6, 8 and 53, which were placed in different groups. In addition, three subgroups were formed within group 1 with the Tocher method, whereas that same group was subdivided into four subgroups with the UPGMA analysis.

With the UPGMA analysis, accessions 12, 57 and 67 were placed in different groups, along with promising accessions in the collection, namely, 1, 39, 48, 50 and 70. However, the Tocher method clustered accessions 12, 57 and 67 within the last positions of group I.3. This positioning indicates that these are materials with a greater genetic distance from the other accessions in the same group and are therefore important from the agronomic point of view and can be recommended as potential progenitors in hybridization programs.

As in the cluster analysis by the Tocher method, the UPGMA clustering pattern was unrelated to the geographic origin of the materials, since accessions from distant regions showed relatively small genetic distances.

In addition to the clustering methods, genetic diversity was evaluated by the canonical variables method, whose basic objective is to provide a structural simplification of the data in two or three dimensional space. The viability of its interpretation is restricted to the concentration of the variability among the first variables, usually above 80% (Cruz et al., 2011).

However, other authors recommend as viable a cumulative variance above 70% in the first canonical variables (Härdle and Simar, 2003; Rencher, 2002). Note also that cluster analysis can be complemented by other visualization techniques, such as canonical variables, as a way of facilitating geometric interpretation (Cruz et al., 2011). That facilitation justifies the use of canonical variables analysis in the present study, since the first two variables explain 72.59% of the total variation (Table 3), and this analysis additionally complements the cluster analysis and provides a structural simplification of the data.

Table 3. Canonical variables and their relative (root (%)) and cumulative (cumulative root (%)) importance for the 23 agronomic characters evaluated in the 70 *C. procera* accessions.

Canonical variable	Root	Root (%)	Cumulative %
1	68.864273	64.762112	64.762112
2	8.327747	7.831673	72.593785
3	5.369828	5.049954	77.643739
4	4.69763	4.417798	82.061537
5	3.136434	2.949601	85.011138
6	2.542641	2.391179	87.402317
7	2.23268	2.099682	89.501998
8	1.931751	1.816679	91.318678
9	1.468157	1.3807	92.699378
10	1.325425	1.246471	93.945849
11	1.149742	1.081254	95.027103
12	0.877134	0.824884	95.851987
13	0.805589	0.757601	96.609588
14	0.692235	0.650999	97.260587
15	0.665455	0.625815	97.886402
16	0.566546	0.532797	98.419199

17	0.490764	0.46153	98.880729
18	0.394032	0.37056	99.251289
19	0.245188	0.230583	99.481871
20	0.213711	0.20098	99.682852
21	0.151786	0.142745	99.825597
22	0.098641	0.092765	99.918362
23	0.086809	0.081638	100.00

Figure 3 shows the scatterplot of the 70 *C. procera* accessions based on the first two canonical variables, with the formation of six distinct groups, taking as reference the agreement with the groups formed by the Tocher and UPGMA methods, indicating the characters' potential to represent the genetic diversity among the genotypes studied.




Figure 3. Scatter plot of scores of the 70 *C. procera* accessions for the first two canonical variables, established by the 23 agronomic characters. Circles delimit the groups formed.

The group compositions obtained by the different analyses, mainly supported by the estimates of genetic parameters, provide support for the identification of promising accessions. These accessions are indicated for use in breeding programs focused on forage production, considering the importance of the agronomic characteristics in the evaluation of productive potential.

For quantifying the relative contribution of the agronomic characters to the genetic diversity, the method proposed by Singh (1981) was used, when possible, to obtain the values of S_j (where S is the measure of relative importance for variable j) for the 23 characters from the dissimilarity measure D^2 . The variables TFM, LFM and SFM were observed to contribute the most to the differentiation of the 70 accessions, with 85.23% of the total variance; and the SDM, LDM and TDM characters contributed with 14.62%. These results identify the agronomic characters to be used in future studies aimed at forage production. Note that plant phytomass is one of the main characteristics of forage plants because this character represents the productive potential of the plant. Therefore, plants

with the potential to reach relatively high phytomass, and therefore relatively high percentages of dry matter, are of great importance for future *C. procera* breeding programs.

The other characters, despite their importance, yielded the lowest estimates of **explain S.i.** This result highlights the need to perform future research with the accessions of the present study, aiming to investigate other characteristics, such as secondary metabolites, as a means to optimize the species breeding program. Mohamed et al. (2015) reported that *C. procera* latex has a high content of active compounds, including cardiotonic glycosides, alkaloids, terpenes, resins, lipids, flavonoids, tannins and steroids. Among these secondary metabolites, cardiotonic glycosides are the most important, **toxic action** in humans and animals. Thus, research that investigates these glycosides in the accessions of the present study is an important future priority.

4. CONCLUSION

Genetic variability in agronomic characters exists among the 70 *C. procera* accessions.

The phytomass characters are the ones that contribute most strongly to the genetic divergence among the *C. procera* genotypes.

Accessions 1, 12, 20, 39, 48, 50, 57, 67 and 70 show promise for inclusion in the *C. procera* breeding program, with agronomic and forage potential, and are indicated for use as parents in hybridization programs and for inclusion in future studies.

REFERENCES

Almeida IVB, Neder DG; Batista FRC and Dutra WF. (2017a). Characterization and early selection of silk blossom (*Calotropis procera*) genotypes with forage potential. *Revista Caatinga* 30 (3): 794-801.

Almeida IVB, Ramos JPC, Dutra WF, Bruno RLA and Alves EU. (2017b). Genetic diversity among *Calotropis procera* (Aiton) WT Aiton genotypes according to seed physiological quality. *Revista Caatinga* 30(4): 912-919.

Bárbaro IM, Mauro A, Centurion M, Perecin D, Unêda-Trevisoli SH, Muniz F and Gavioli, E. (2004). Seleção em populações F3 de soja visando à resistência ao cancro da haste e bons atributos agrônômicos. *Revista Ceres* 51: 619-634.

Barbosa MO, Almeida-Cortez JS, Silva SI and Oliveira AFM. (2014). Seed oil content and fatty acid composition from different populations of *Calotropis procera* (Aiton) WT Aiton (Apocynaceae). *Journal of the American Oil Chemists' Society* 91(8): 1433-1441.

BRASIL. Ministério da Agricultura, Pecuária e Abastecimento. (2009). Regras para análise de sementes. Ministério da Agricultura, Pecuária e Abastecimento. Secretaria de Defesa Agropecuária, Brasília: MAPA/ACS, 399p.

Costa AM, Spehar CR and Sereno JRB. (2012). Conservação de recursos genéticos no Brasil. Brasília, DF: Embrapa, 628p.

Costa RG, Medeiros AN, Alves AR and Medeiros GR. (2009). Perspectivas de utilização da flor-de-seda (*Calotropis procera*) na produção animal. *Revista Caatinga* 22(1): 1-9.

Cruz CD. (2005). Princípios da genética quantitativa. Viçosa, MG: UFV, 394p.

Cruz CD. (2006). Programa GENES: análise multivariada e simulação. Viçosa-MG, Ed. UFV, 175 p.

- Cruz CD. (2013). Genes: a software package for analysis in experimental statistics and quantitative genetics. *Acta Scientiarum Agronomy* 35(3): 271-276.
- Cruz CD, Ferreira FM and Pessoni LA. (2011). *Biometria aplicada ao estudo da diversidade genética*. Viçosa: Editora UFV, 620p.
- El-Bakry AA, Hammad IA and Rafat FA. (2014). Polymorphism in *Calotropis procera*: preliminary genetic variation in plants from different phytogeographical regions of Egypt. *Rendiconti Lincei* 25(4): 471-477.
- Farahat EA, Galal TM, El-Midany MM and Hassan LM. (2015). Phenology, biomass and reproductive characteristics of *Calotropis procera* (Aiton) WT Aiton in South Cairo, Egypt. *Rendiconti Lincei* 27(2): 197-204.
- Härdle W and Simar L. (2003). *Applied Multivariate Statistical Analysis*. Berlin: MD Tech, 488p.
- Hassan LM, Galal TM, Farahat EA and El-Midany MM. (2015). The biology of *Calotropis procera* (Aiton) WT. *Trees* 29(2): 311-320.
- Maguire JD. (1962). Seed of germination, aid in selection and evaluation for seedling emergence and vigor. *Crop Science* 2: 176-177.
- Maroufi K, Farahani HA and Moaveni P. (2011). Effects of hydropriming on seedling vigour in spinach (*Spinacia oleracea* L.). *Advances in Environmental Biology* 5 (8): 2224-2227.
- Mohamed NH, Liu M, Abdel-Mageed WM, Alwahibi LH, Dai H, Ismail MA and Shoreit AA. (2015). Cytotoxic cardenolides from the latex of *Calotropis procera*. *Bioorganic & medicinal chemistry letters* 25(20): 4615-4620.
- Mojena R. (1977). Hierarchical grouping methods and stopping rules: an evaluation. *The Computer Journal* 20: 359-363.
- Moreira Filho EC, Silva DS, Pereira WE, Cabral Júnior CR, Andrade MVM, Silva GE and Viana BL. (2007). Estimacão da área foliar da flor de seda (*Calotropis procera*). *Archivos de Zootecnia* 56: 245-248.
- Negreiros JRDS, Alexandre RS, Álvares VDS, Bruckner CH and Cruz CD. (2008). Divergência genética entre progênies de maracujazeiro-amarelo com base em características das plântulas. *Revista brasileira de Fruticultura* 30(1): 197-201.
- Oliveira-Bento SRS, Torres SB, Bento DAV, Silva BKA, Dantas FJC and Melo VC. (2015). Armazenamento de sementes de flor-de-seda [*Calotropis procera* (AITON) W.T. AITON]. *Revista Caatinga* 28(1): 39 - 47.
- Rahman MA and Wilcock CC. (1991). A taxonomic revision of *Calotropis* (Asclepiadaceae). *Nordic Journal of Botany* 11(3): 301-308.
- Ramadan A, Sabir JS, Alakilli SY, Shokry AM, Gadalla NO, Edris S and Baker NR. (2014). Metabolomic response of *Calotropis procera* growing in the desert to changes in water availability. *Plos One* 9(2): e87895.
- Ramos SRR, Queiroz MD and Pereira TNS. (2007). Recursos genéticos vegetais: manejo e uso. *Magistra* 19: 265-273.

389 Ranjit PM, Eswara RG, Krishnapriya M, Nagalakshmi V, Silpa P and Anjali M. (2012). An
 390 overview of phytochemical and pharmacological activities of *Calotropis procera*. FS Journal
 391 of Pharmacy Research 1(2): 18-25.
 392
 393 Rathore M and Meena RK. (2010). Potential of utilizing *Calotropis procera* flower biomass as
 394 a renewable source of energy. Journal of Phytology 2(1): 18-23.
 395
 396 Rencher AC. (2002). Methods of Multivariate Analysis. New York: Wiley-Interscience, 740p.
 397
 398 Rohlf FJ. (1970). Adaptive hierarchical clustering schemes. Systematic Zoology 19(1): 58-
 399 82.
 400
 401 Santos RC, Moreira JAN, Farias RH and Duarte JM. (2000). Classificação de genótipos de
 402 amendoim baseada nos descritores agromorfológicos e isoenzimáticos. Ciência Rural 30(1):
 403 55-59.
 404
 405 Silva DJ and Queiroz AC. (2002). Análises de alimentos: métodos químicos e biológicos.
 406 3.ed. Viçosa, MG: UFV, 253p.
 407
 408 Singh D. (1981). The relative importance of characters affecting genetic divergence. The
 409 Indian Journal of Genetics e Plant Breeding 41: 237-245.
 410
 411 Vencovsky R. (1987). Herança quantitativa. In: PATERNIANI, E. (Ed.) Melhoramento e
 412 produção de milho no Brasil. Piracicaba: Fundação Gargill.
 413
 414 Yao DARA, Sprycha Y, Porembski S and Horn R. (2015). AFLP assessment of the genetic
 415 diversity of *Calotropis procera* (Apocynaceae) in the West Africa region (Benin). Genetic
 416 Resources and Crop Evolution 62(6): 863-878.