

Addition of charcoal waste to earthworm humus promotes better results in the production of papaya seedlings

ABSTRACT

Aimed at improving the physical and chemical attributes of the substrates for seedling production, several agro-industrial wastes have been used. One of the waste with potential of substrate incorporation is the charcoal waste. The objective of this study was to evaluate the effect of increasing doses of charcoal waste (CR) on the production of papaya seedlings (*Carica papaya* L.). The experimental design was a randomized block design, as an experimental model. The treatments were composed of 5 (five) concentrations of CR added to earthworm humus based on volume, and the following treatments were evaluated: T1: 100% earthworm humus + 0% CR (control); T2: 97.5% humus + 2.5% CR; T3: 92.5% humus + 7.5% CR; T4: 85% humus + 15% CR and T5: 70% humus + 30% CR. Each of the 5 (five) treatments had 4 (four) replicates (blocks). The evaluations consisted in plant height measurement, collection diameter, root length, dry mass of leaves and roots, and Dickson quality index. Regression models were adjusted ($P < 0.05$) to treatments with increasing levels of CR. CR doses around 10% promoted better results of collection diameter, root length and root dry mass. For the other variables, CR doses around 13% promoted the best results. It is believed that the addition of CR promoted significant improvements in earthworm humus; however, high doses promoted elevation of pH at levels considered critical for the availability of some nutrients for the crop. In general, the best values obtained for the analyzed variables were those in which about 9.25% to 14.25% of the CR addition to the humus used.

Keywords: *Carica papaya* L., charcoal, humus, alternative substrate

1. INTRODUCTION

Papaya (*Carica papaya* L.) is one of the most consumed fruits in the world [1]. Brazil ranks second place as a world fruit producer with a production of 1,517.696 to per year⁻¹, ranking among the main exporting countries [2].

One of the problems related to the production of papaya fruit in Brazil is the culture establishment in the field because of the hard time to obtain quality seedlings within desirable phytosanitary characteristics [3]. The productivity and quality of the papaya fruits are directly related to the cultural practices, which have begun since the formation and implantation of the seedlings in the field [4].

Several factors can contribute to the production of seedlings with quality, among them, the substrate has great relevance. To be considered a good substrate, the material must have attributes such as aeration, porosity, nutrient availability and water retention, being these attributes essential for the good development of the plants [5]. In addition, the use of the substrate is due to its availability, cost and obtaining easiness and handling [6].

One of the substrates commonly used by farmers in the production of seedlings, mainly those working in organic scheme, is earthworm humus or vermicompost. The humus can be considered a plant growth biostimulator, moreover, it also stimulates the present microbiota in the substrate, improving plant health [7]. However, the same author defends that the humus concentration in the substrate, to produce forest seedlings, it should not exceed 80%, and it is necessary to incorporate some other element with good availability and low cost so that it acts as a good substrate.

To improve the physical and chemical attributes of the substrates for seedlings production, several agro-industrial waste has been used, so their accumulation in the industrial yards is reduced, and they have a more sustainable disposal [8]. One of the waste with potential for incorporation into substrate to produce papaya seedlings is the charcoal waste. The carbonized biomass (charcoal) has some

physico-chemical characteristics, such as basically aromatic molecular structure, high specific surface area and hydrophobicity [9] and, depending on the active sites, gives greater protection to soil microorganisms [10]. Thus, coal waste can act as a substrate conditioner [11]. According to the scientific literature, there are promising results of the use of charcoal waste for crops such as lettuce [12], rice [12], soybean and rice [13], eucalyptus [14], among others. Although it is used in the production of seedlings of some crops, little is known about the use of charcoal waste in the production of papaya seedlings. The objective of this work was to evaluate the effect of the addition of increasing doses of charcoal waste on earthworm humus in the production of papaya seedlings (*Carica papaya* L.).

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The experiment was installed and conducted in the Fruticulture Sector of the Federal Institute of Education, Science and Technology of Espírito Santo (Ifes), Santa Teresa campus, from November 2016 to January 2017, in a seedling nursery covered with polyolefin screen with 50% of shading, located in the municipality of Santa Teresa, state of Espírito Santo, with altitude of 155 m, latitude of 18° 48'South and longitude of 40° 40' West. The climate of the region, according to the Köppen climate classification of, fits in the Cwb type (temperate maritime climate/altitude tropical climate), with average annual temperature of 24.6°C and annual precipitation ranging from 700 to 1200 mm.

During the seedling production, were used Solo Gropu's papaya seeds from healthy and ripe fruits, as propagative material. The fruits of this group present the fruit size, with a weight varying from 300 g to 650 g. In addition, the shape of the well rounded fruits contribute to make them one of the most requested for export. Another characteristic is the precocity in the production, the orange-red pulp, which attracts the consumers, and with excellent taste [1]. The fruits were cut in half, separating pulp and seed. The seeds were washed in running water on fine mesh sieve to eliminate mucilage and pulp and bark waste. The selection was made through manual harvesting for the small and damaged seeds disposal. The drying was held in an airy and shaded place, during 6 (six) days.

The materials used in the formulation of the alternative substrate to produce papaya seedlings were earthworm humus and charcoal waste (CR). The earthworm humus was used in the experimental area of the Nucleus of Studies in Agroecology of the same Campus, from the digestion of cattle manure tanned by the intestinal tract of the rural Californian earthworms (*Eisenia foetida*) in, following the specifications of the EMBRAPA model, adapted by [15]. The charcoal waste (CR) used, originating from eucalyptus wood (*Eucalyptus* spp.), came from charcoal farms around the region. The CR was ground and sieved until a fine powder was obtained.

A randomized complete block design was used as the experimental model. The treatments were composed of 5 (five) CR concentrations added to the earthworm humus based on the volume, and the following treatments were evaluated: T1: 100% earthworm humus + 0% of CR (control); T2: 97.5% humus + 2.5% CR; T3: 92.5% of humus + 7.5% of CR; T4: 85% of humus + 15% of CR and T5: 70% of humus + 30% of CR. Each one of the 5 (five) treatments had 4 (four) replicates (blocks), considering 25 seedlings each, resulting in 500 seedlings throughout the experiment. Six central plants of each experimental unit were considered useful.

The humus physico-chemical analysis and charcoal waste consisted of determination of the electrical conductivity (EC) through bench conductivity meter. The chemical analyzes consisted of the determination of the pH by means of a bench potentiometer and the quantification of total organic carbon (TOC), organic matter (OM), total nitrogen (TN), phosphorus (P), potassium (K) and calcium (Ca), performed at the Soil and Solid Waste Laboratory of the Department of Agricultural Engineering of the Federal University of Viçosa (Table 1).

Table 1 presents the physical-chemical and chemical characteristics of the materials used in the formulation of the substrates.

Table 1. Physical-chemical (EC) and chemical characteristics of humus and charcoal waste (CR) used in the experiment

Materials	PH	EC	TOC	OM	TN	P	K	CA
		dS m ⁻¹				dag kg ⁻¹		
Humus	6,67	280	74,1	46,79	0,75	1,64	0,83	1,40
CR	7,77	283	76,95	-*	0,41	0,023	0,66	0,330

90 EC: electrical conductivity; TOC: total organic carbon; OM: organic matter; TN: total nitrogen; P: Phosphorus; K:
 91 Potassium; Ca: Calcium.
 92 * Organic matter: Charcoal waste is considered organic matter.
 93

94 For the papaya seedlings production was used suspended trays, allocated in masonry benches,
 95 containing 500 mL tubes suitable for germination. The mixture of CR to the humus happened
 96 manually until obtaining a homogeneous material, being held 7 (seven) days before sowing.
 97 According to nursery routine, the irrigation was performed manually. No pesticides types were
 98 applied, nor was any other correction process performed on the substrates.

99 Evaluations occurred at 60 (sixty) days after sowing (DAS). The seedlings were removed from the
 100 core and were carefully washed. The evaluation of seedling development was held by the following
 101 parameters: plant height (PH), collection diameter (CD), root length (RL), dry leaf mass (DLM) and
 102 root (DRM) and determination of the Dickson quality index (DQI).

103 To obtain the height of the aerial part was used a ruler millimeter, measuring from the stem base to
 104 the apical bud that gave rise to the last leaf. The sample diameter was measured by 0.01 mm
 105 precision digital caliper. To calculate the aerial part dry mass, it was considered only leaves. These
 106 leaves were extracted during the insertion of the petiole of the plants period. To obtain the root dry
 107 mass, the roots were carefully washed in running water, on a 5-mm sieve. The roots were measured
 108 from the collecting region of the root with the aid of a millimeter ruler. Subsequently, both plant
 109 materials were packed in paper bags, properly identified and dried in a greenhouse with forced air
 110 circulation at 65°C until reaching constant weight. Soon afterwards, the materials were weighed in a
 111 precision electronic scale of 0.01 g. To determine the DQI, the method proposed by Dickson et al.
 112 (1960), (Equation 1).
 113
 114


$$DQI = \frac{TDM (g)}{[PH(cm)/CD(mm)] + (DLM (g)/DRM (g))} \quad \text{Eq. 1}$$

115
 116
 117 where: DQI = Dickson quality index; TDM = Total dry matter mass (g); PH = Plant height (cm); CD =
 118 Collection diameter (mm); DLM = Dry leaf mass (g), and DRM = Dry root mass (g).
 119

120 All the evaluated variables were submitted to the normality tests (Lilliefors) and homoskedasticity
 121 (Bartlett) tests, which are the assumptions for the analysis of variance, using adjustments in
 122 regression models for the orthogonal polynomials method. A "α" of 0.05 was used for all procedures.
 123

124 3. RESULTS AND DISCUSSION

125
 126 The Figure 1 (A, B, C, D, E and F) shows the plant height values (PH), root length (RL), collection
 127 diameter (CD), dry leaf mass (DLM), dry root mass (DRM) and Dickson quality index (DQI) of papaya
 128 seedlings respectively, in response to the addition of increasing doses of charcoal waste on the
 129 substrate.
 130






Figure 1. Plant height (A), root length (B), collection diameter (C), dry leaf mass (D), dry root mass (E) and Dickson quality index of papaya seedlings respectively, in response to the addition of increasing doses of charcoal waste in the substrate.

The plant height variable showed increase with the CR increasement of to the humus to a concentration of 12.75%, providing a maximum height of 20.48 cm (Figure 1 (A)). Lower results of this study were found by [16], which evaluated different substrates to produce papaya seedlings. It was found greater growth, about 7 cm, with substrate composed by a mixture of Plantmax + humus + cattle manure (1: 2: 1); however, the evaluation took 37 DAS. Similarly, to the variable plant height, the variables root length and collection diameter presented the best results of 13.57 cm and 6.08 mm with the additions of CR to humus of 9% and 9.75%, respectively.

For the variables dry leaf mass and root dry mass (Figure 1 (D) and (E)) it is observed that the increase of CR to humus promoted significant results similar to the other variables, with the best concentrations of CR to humus at 11.25% and 10.75%, respectively, promoting a dry biomass gain of leaves and roots of 0.87 g and 0.55 g, respectively. [17], studying different substrates in the formation of papaya seedlings of the soil group, found better results for plant height of 31.45 cm with substrates containing bovine manure, soil, sand and vermiculite (2: 1: 1: 1 v / v) and Plantmax, corral dung, soil and sand (1: 1: 1: 1 v / v). In the same experiment, [17], found average values of 1.64 g evaluating the whole aerial part of the plants at 90 DAP and not only leaves at 60 DAP herein. These authors attribute the best constitution results of substrates rich in macronutrients such as P and K, these are of foremost importance in the development of papaya cultures. Moreover, the physical substrates characteristics are mentioned by [17] as indicative of satisfactory results because they present higher water retention capacity, aeration and nutrients, thus producing better quality seedlings.

The P element is very important in papaya crops, positively influencing the increase of the aerial and root system of the plants, mainly in the initial phase of plant growth [18]. According to Table 1, this element presents in good amount of earthworm humus, but an incipient amount in CR. Thus, whereas the addition of CR increased to the humus, it means a reduction in the proportion of this and other nutrients in the total volume of the substrate was promoted, which may have contributed to the reduction of the response in higher CR concentrations. At lower concentrations, the addition of CR to humus may have promoted conditioning action of the substrate, making it more suitable for production of papaya seedlings, contributing to greater retention of nutrients and water, and reducing physical barriers, which favors better plant development [11].

Some studies report positive effects of the addition of coal in the production of various seedlings. [19] evaluating the effect of growing doses of charcoal in the production of *carvoeiro* (*Tachigali vulgaris*) seedlings found increasing dry shoot and root biomass growing doses of up to 50% of charcoal. [20], when evaluating the effect of charcoal application on different substrates, they observed a major increase of dry shoot biomass and roots in *crajiru* seedlings (*Arrabidaea chica* Verlot.). These results corroborate with the results obtained herein, evidencing that the addition of CR promotes physical-chemical improvements of the substrate, leading to greater retention of moisture and nutrient availability.

For DQI, a variable includes several correlated elements to the quality of formed seedlings and the addition of CR in 11% to humus promoted the best result (0.2896). On contrary, higher doses promoted the reduction of this index (Figure 1 (F)). The value of DQI found herein points out that the CR addition at 11% to the earthworm humus that provides better papaya seedlings. [21] evaluating different substrates in the production of papaya seedlings, except for the treatment containing vermiculite, (IQD = 0,334), the other treatments presented lower IQD than those found in this study. However, in [21], organic and mineral fertilization were used to enrich the substrate. It is worth to point out that no nutritional correction or addition was used to the substrate in this study and, therefore, it is believed that the superiority of the DQI result found in this work in relation to those found by [21], is mainly a result of the interaction of the CR with the worm humus, indicating the combination of earthworm humus and CR may be an alternative in the substitution of commercial substrates.

In general, the addition of CR to the humus is between 9.25% and 14.25% promoted better results in the evaluated variables. [19] state that the addition of coal fines or charcoal waste can promote major improvements in the physical and chemical properties of the soil, correcting its acidity, increasing the levels of exchangeable Ca, extractable P, CTC, besides soil aggregation capacity. Although CR is considered an inert material, its structure has sites capable of performing ionic exchanges, thus promoting better availability of nutrients present in the substrate [22]. Thus, the addition of CR up to certain concentrations to earthworm humus may have provided greater stability of the substrate as well as favoring the absorption of nutrients by plants.

Figure 2 shows the plot of substrate pH values as a function of the various levels of charcoal waste insertion incorporated into the humus in the production of papaya sludge.




Figure 2. pH values of the substrate as a function of the various levels of charcoal waste incorporated into the humus in the production of papaya seedlings.

The CR addition promoted a linear pH increase of the substrate solution (Figure 2). According to [23] the ideal pH range is between 6.0 and 6.5 but there may be a behavioral variation of each plant species in relation to pH and its effects on the substrate. This alteration (acidity/alkalinity) in the substrate solution may promote the unavailability of some nutrients, such as P, N, S and B, which remain with high availability up to a pH range of 7.3 [24]. Thus, higher ratios than 11% promoted elevation of soil alkalinity (Figure 2) at indexes that may promote precipitation, especially P, making it unavailable to plants [25]. P is one of the main elements in initial plant growth, acting in storage, energy transfer and absorption of other nutrients [26]. Therefore, its lack can severely compromise the development of plants. [27], evaluating papaya seedlings observed an increase in the dry matter of plant roots due to the positive effect of P, which increased the capacity of nutrient absorption due to the greater root development.

According to the data obtained in this research, the CR addition can make earthworm humus more suitable to produce papaya seedlings. It is therefore believed that such fact is due to the increase of porosity and greater retention of water, besides the reduction of the physical barriers in the substrate that are propitiated by the incorporation of CR to the humus. Thus, the CR addition can promote significant improvements in earthworm humus, making it an important substrate for production of papaya seedlings, minimizing the production costs, increasing the profitability of the productive chain of seedlings production.

In addition to the quality and sanity with which the seedlings are formed, the addition of CR to the substrate can promote the sustainability of the coal sector, being an option for the final waste destination waste, also contributing to the carbon storage in the soil, reducing the harmful gas emission to the greenhouse effect [28], considering that [29] state that a great part of the C emitted by anthropic activities (12%) could be eliminated by applying C in soil in the form of carbonized biomass.

4. CONCLUSION

The best values obtained from the analyzed variables of papaya plants were 9% to 14% of the insertion of charcoal waste to the humus.

The addition of CR to the earthworm humus promoted elevation of the substrate pH.

REFERENCES

1. Serrano LAL, Cattaneo LF. O cultivo do mamoeiro no Brasil. Revista Brasileira de Fruticultura. 2010; 32(3): 657-659.
2. Empresa Brasileira De Pesquisa Agropecuária. Mandioca e Fruticultura. Brasília: EMBRAPA. Disponível em: <https://www.embrapa.br/mandioca-e-fruticultura/cultivos/mamoa>. Acesso em: 04 de Jul 2018.
3. Francisco MGS, Maruyama WI, Mendonça V, Silva EA, Reis LL, Leal ST. Composition of substrates and volume of recipients in the production of papaya 'sunrise solo'. Revista Agrarian. 2010; 3(9): 267-274.
4. Araújo WBM, Alencar RD, Mendonça V, Medeiros EV, Andrade RC, Araújo RR. Goat manure used in substrate composition for formation of papaya seedlings. Ciência e Agrotecnologia. 2010; 34(1): 68-73.
5. Terra SB, Ferreira AAF, Peil RMN, Stumpf ERT, Beckmann cavalcante MZ, Cavalcante IH L. Alternative substrates for growth and production of potted chrysanthemum (cv. Funny). Acta Scientiarum Agronomy. 2011; 33(3): 465-471.
6. Silva JDC, Leal TTB, Araujo RM, Gomes RLF, Araujo ASF, Melo WJ. Germination and initial growth of ornamental Capsicum and Celosia in substrate of composted tannery sludge. Revista Ciência Rural. 2011; 41(3): 412-417.
7. Steffen GPK, Antonioli ZI, Steffen RB, Schiedeck G. Use of vermicompost as a substrate for the production of Eucalyptus grandis and Corymbia citriodora seedlings. Revista Brazilian Journal of Forestry Research. 2011; 31(66): 75-82.

8. Silva Júnior JV, Cavalcante MZB, Brito LPS, Avelino RC, Cavalcante ÍHL. Aproveitamento de materiais alternativos na produção de mudas de tomateiro sob adubação foliar. *Revista Ciência Agronômica*. 2014; 45: 528-536.
9. Carvalho MTM, Maia AHN, Madari BE, Bastiaans L, Van Oort LPA, Heinemann AB, Silva MAS, Petter FA, Marimon Jr. BH, Andmeinke, H. Biochar increases plant-available water in a sandy loam soil under anaerobic rice crop system. *Solid Earth*. 2014; 5:939-952.
10. Brady NC, Weil RR. The nature and properties of soils. 14. ed. Prentice Hall, Upper Saddle River, NJ, USA. 2008; 965 p.
11. Petter FA, Marimon BHJ, Andrade FR, Schossler TR, Golçalves LG, Marimon BS. Biochar como condicionador de substrato para produção de mudas de alface. *Revista Agrarian*. 2012; 5(17): 243-250.
12. Madari BE, Cunha TJF, Novotny EH, Milori DMBP, Martin Neto L, Benites VM, Coelho M R, Santos GA. Matéria orgânica dos solos antrópicos da Amazônia (terra preta de índio). Suas características e papel na sustentabilidade da fertilidade do solo. In: Teixeira WG, Kern DC, Madari BE, Lima EM, Woods WI. *As Terras Pretas de Índio da Amazônia: Sua Caracterização e Uso deste Conhecimento na Criação de Novas Áreas*. 1º ed. Manaus: Embrapa Amazônia Ocidental. 2009; 172-188.
13. Petter FA. Biochar as soil amendment: agronomic and environmental aspects of its application in soils of the Brazilian savana (cerrado). Universidade Federal de Goiás. Tese Doutorado em Produção Vegetal. 2010; 130 p.
14. Petter FA, Andrade FR, Marimon BHJ, Golçalves LG, Schossler TR. Biochar como condicionador de substrato para produção de mudas de eucalipto. *Revista Caatinga*. 2011; 25(4): 44-51.
15. Schiedeck G, Gonçalves MM, Schwengber JE. Minhocultura e produção de húmus para a agricultura familiar. *Circular técnica*. 2006; 57.
16. Góes GB, Mendonça V, Medeiros PVQ, Tosta MS, Medeiros LF. Diferentes substratos na produção de mudas de mamoeiro em bandejas. *Revista Verde*. 2010; 5(1):178-184.
17. Negreiros J, Braga L, Alvares V, Bruckner C. Different substrata for a production of papaya tree (*Carica papaya* L.) group solo seedlings. *Revista Brasileira de Agrociência*. 2005; 11(1): 101-103.
18. Taiz L, Zeiger E. *Physiology Vegetal*. ed. 5. Porto Alegre, Artmed. 2013.
19. Souchie FF, Junior BHM, Petter FA, Madari BE, Marimon BS, Lenza E. Charcoal as amendment to substrate of seedlings of *Tachigali vulgaris* L.G. Silva & H.C. Lima. *Ciência Florestal*. 2011; 21(4): 811-821.
20. Souza GKA, Teixeira WG, Reis RA, Chaves FCM, Xavier JJBN. Growth of cajuru (*Arrabidaea chica* Verlot.) on different growing media. *Revista Brasileira de Plantas Medicinais*. 2006; 8(n. esp): 62-65.
21. Costa E, Mesquita VDAG, Leal PAM, Fernandes CD, Abot AR. Formação de mudas de mamão em ambientes de cultivo protegido em diferentes substratos. *Revista Ceres*. 2010; 57(5): 679-685.
22. Novotny EH, Hayes MHB, Madari BE, Bonagamba TJ, De Azevedo ER, De Souza AA, Song G, Nogueira CM, Mangrich AS. Lessons from the terra preta de índios of the amazon region for the utilization of charcoal for soil amendment. *Journal of the Brazilian Chemical Society*. 2009; 20(6): 1003-1010.
23. Raij BV. Fertilidade do solo e adubação. São Paulo, Agronômica Ceres. 1991; 343p.

- 316 24. Souza DMG, Miranda LN, Oliveira AS. Acidez do solo e sua correção. In: NOVAIS RF, Alvarez
317 VVH, Barros NF, Fontes RLF, Cantarutti RB, Neves JCL (Eds.). Fertilidade do solo. Viçosa,
318 Sociedade Brasileira de Ciência do Solo. 2007; 471-550.
319
- 320 25. Novais RF, Smyth TJ, Nunes FN. Fósforo. In: Novais RF, Alvarez VVH, Barros NF, Fontes RLF,
321 Cantarutti RB, Neves JCL (Eds.). Fertilidade do solo. Viçosa, Sociedade Brasileira de Ciência do
322 Solo. 2007; 471- 550.
323
- 324 26. Brasil EC, Nascimento EVS. Influência de calcário e fósforo no desenvolvimento e produção de
325 variedades de maracujazeiro-amarelo. Revista Brasileira de Fruticultura. 2010; 32:892-902.
326
- 327 27. Saraiva KR, Nascimento RS, Sales FAL, Araújo HF, Fernandes CNV, Lima AD. Papaya seedling
328 production with doses of phosphoric fertilization using superphosphate as source. Revista Brasileira
329 de Agricultura Irrigada. 2011; 5:376-383.
330
- 331 28. Madari BE, Costa AR, Castro LM, Santos JL, Benites VM, Rocha AO, Machado PLOA.
332 Comunicado Técnico 125. Goiânia, GO. EMBRAPA. Centro Nacional de Pesquisa de Arroz e Feijão.
333 2006; 4p.
334
- 335 29. Lehmann J, Gaunt J, Rondon M. Bio-char sequestration in terrestrial ecosystems a review.
336 Mitigation and Adaptation Strategies for Global Change, Netherlands. 2006; 11(2): 403 427.
337