

Original Research Article

Resistance induction in *Zea mays* L. cultivars to *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

ABSTRACT

The maize crop (*Zea mays* L.) faces serious limitations in cultivation due to attack by the *Spodoptera frugiperda* insect-pest. The objective of this work was to evaluate the resistance induction in maize cultivars with *S. frugiperda* caterpillars due to the application of Silicon and Calcium sources. Two cultivars of maize, Genotype A simple hybrid and the Creole Genotype B variety were used. These cultivars were submitted to two treatments of foliar fertilization, calcium chloride and potassium silicate, and a treatment without the application of an elicitor. The evaluations of the damages caused by the *S. frugiperda* caterpillar were carried out weekly, from the seventh day after the first application of the products (Si and Ca). In the bolting phase (VT), physiological evaluations of gas exchanges and fluorescence of Chlorophyll *a* were carried out. For the Creole variety, there was a predominance of plants that did not present any type of damage. The Creole variety presented the lowest mean of infestation by *S. frugiperda*, regardless of the product applied. No statistical differences were observed between the products applied in each cultivar. The maximum quantum efficiency of FSII (F_v/F_m) was not statistically significant for the cultivars nor for the cultivars x treatments interaction. For GS, a statistical difference between cultivars was found. In the Creole genotype B variety, superior results were obtained in relation to the simple hybrid. The TF variable did not present statistical difference ^{between} among the evaluated cultivars. Different behaviors were observed in the transpiration variable (E) among the cultivars. In the Creole genotype B variety when submitted to silicon application and in the control, a higher transpiration was obtained comparing to the simple hybrid. The sources of Silicon and Calcium do not promote the induction of resistance to *S. frugiperda* in the studied variables. The Creole variety Genotype B of *Zea mays* suffers minor damage due to infestation of *Spodoptera frugiperda* in relation to the simple hybrid maize Genotype A.

Keywords: control method, fall armyworm, corn

1. INTRODUCTION

Zea mays L. maize is one of the world's largest cereals grown with the United States, China and Brazil as the world's largest producers. In the export ranking, Brazil occupies the second position [1], the Central West and South being the main producing regions of this cereal [2].

Although the country is prominent in the world production, it still presents limitations in the cultivation, which implies low productivity mainly due to insect pest attacks. Among the insect complexes that infest the maize crop, the *Spodoptera frugiperda* caterpillar (J. E. Smith, 1797) is the main cause of damage and represents a high cost of control annually [3].

20 The fall armyworm becomes an important pest of this crop, both by reducing productivity and
 21 decreasing the quality of the final product, besides the difficulty of control [4]. The
 22 considerable economic losses caused by this pest cause farmers to intensify the use of
 23 chemical insecticides and consequently lead to the development of resistance to many
 24 insecticide groups, making their control even more difficult [5]

25 Different control methods are used to reduce populations of *S. frugiperda*, such as cultural,
 26 chemical and biological tactics [6]. Plant resistance is also easily incorporated into integrated
 27 pest management because it maintains the pest population below the level of economic
 28 damage without unbalance to the environment and has cumulative and persistent effect
 29 beyond the lowest cost of production [7].

30 Due to the different biochemical, morphological and physiological characteristics among the
 31 genotypes of the same species, it is possible that some plants suffer a greater or lesser
 32 impact of phytophagy [8]; [9]. Plant resistance to pests can also be obtained by induction,
 33 which is a temporary manifestation, where a plant becomes less favorable to insect attack
 34 due to a certain condition that affects its physiology [7]. Among the various induction
 35 techniques, fertilizers have been widely used, mainly micro and macronutrients.

36 Silicon (Si) is considered a beneficial element for plants [10], it acts on the reinforcement of
 37 the cell wall by depositing solid silica, making the plant more rigid, with better growth and
 38 yield, for suppressing diseases and conferring characteristics of resistance to insects [11];
 39 [12]. Calcium (Ca) is a macronutrient that is involved in a large number of signaling
 40 pathways in plants, from symbiotic interactions, defense responses and various hormones,
 41 functioning as a secondary messenger in response to biotic and abiotic signals [13]. Calcium
 42 participates in lignin biosynthesis, which favors plant self-defense and can be used as an
 43 inducer of pest resistance [14].

44 Other biomorphological factors are also associated with plant defense in relation to insect
 45 infestation. As a consequence of the stress caused by pest injury, there are reductions in the
 46 physiological parameters such as photosynthetic rate, transpiration and stomatal
 47 conductance [15]; [16]. For this purpose, the gas exchange and fluorescence analysis of
 48 chlorophyll "a" are able to detect, with safety and reliability, stress and injury effects in the
 49 photosynthetic process caused by biotic or abiotic factors [17]. In view of the above, the
 50 objective of this work was to evaluate induction resistance in corn cultivars due to the
 51 application of Silicon and Calcium sources in the infestation of the *Spodoptera frugiperda*
 52 caterpillar.

53

54 2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

55

56 The experiment was conducted at the Chã de Jardim experimental farm, belonging to the
 57 Agricultural Sciences Center (CCA) of the Federal University of Paraíba (UFPB), Areia-PB,
 58 Brazil. The climate of the region is classified as tropical humid, presenting average
 59 temperature, relative humidity and precipitation of 23°C, 80% and 1,400 mm annually,
 60 respectively [18]. The soil is classified as Yellow Latosol [19].

61 The simple hybrid corn denominated Genotype A and the Creole variety denominated
 62 Genotype B were used. These materials were submitted to two treatments of foliar
 63 fertilizations, being applied commercial sources of Calcium Chloride and Potassium Silicate,
 64 and a control treatment. Three applications of the products were carried out in the vegetative
 65 growth phase of the maize cultivars, specifically in the V3, V6 and V12 stages, following the
 66 recommended dosages for culture suggested by the manufacturer of 2 L ha⁻¹. The products

67 were always applied from 4pm, this being the most recommended time for loss reduction,
68 and manually by a costal spray with a 20-liter capacity.

69 The plots were constituted by four rows of 5 m in length, where after the mechanical
70 preparation of the soil with a harrowing, furrows were opened and manual sowing was done,
71 allowing a spacing of 0.50 m between rows and 0.20 m between plants, totaling 100 plants
72 per plot and a density of 100,000 plants/ha. The useful area of the plot was constituted by
73 the two central rows, scattering 0.5 m of the border. The nutritional correction was carried
74 out with the fertilization of macronutrients in the foundation and cover, and soil acidity, with
75 dolomitic limestone application, according to results of soil chemical analysis. Weed control
76 was performed with manual weeding.

77 2.1 Evaluation of the damage and infestation of *Spodoptera frugiperda*

78 The evaluations of the damages caused by *S. frugiperda* were carried out weekly, from the
79 seventh day after the first application of the products (Si and Ca), with this done at 7, 14, 21,
80 35 and 42 days, ending at the beginning of flowering. Ten plants per plot were randomly
81 selected, with individual scores varying between 0 and 5 (Table 1) [20].

82 **Table 1. Scale of damages caused by *Spodoptera frugiperda* in *Zea mays* plants [20].**

Score	Damage description
0	No damage
1	Scraped leaves
2	Bored leaves
3	Ripped leaves
4	Stem damage
5	Stem destroyed

83

84 To evaluate the level of infestation of *S. frugiperda*, the two central rows of each plot were
85 used, relating the number of plants attacked by the total number of plants in the crop line.

86 2.2 Evaluation of physiological characteristics

87 The Fluorescence of Chlorophyll a, was determined from a portable fluorometer, model QS-
88 30p from Opti-Sciences®. Measurements were made on three plants per plot, between 9
89 and 12 hours, at 2 cm from the margin of the leaf, excluding its veins. To that end, one leaf
90 per plant, randomly selected was adapted to the dark with the use of tweezers, for
91 approximately 30 minutes. The maximum FSMI photochemical quantum (Fv/Fm) parameter,
92 often evaluated by the maximum fluorescence rate of chlorophyll "a", was evaluated to
93 compare healthy and damaged tissues due to biotic factors [21].

94 In the maize bolting (VT) phase, gas exchange and photosynthetic efficiency of maize leaves
95 were evaluated using an Infrared Gas Analyzer (IRGA), LCpro-SD model from
96 BioScientific®. Two plants were used per plot in the morning, between 9am and 11am,

97 sampling the first leaf opposite and below the ear. The following physiological parameters
 98 were evaluated: Transpiration (E), Internal CO₂ concentration in leaf (Ci), Stomatal
 99 conductance (gs), Photosynthesis (A), Carboxylation efficiency (EC), Pressure Deficit (DP)
 100 and Efficiency of use of water (EFUA).


101 2.3 Statistical analysis

102 The experimental design was a randomized complete block (DBC), in a factorial scheme
 103 (2x3x4), two maize cultivars (genotype A and genotype B), two sources of resistance
 104 induction (Si and Ca) plus one control (without application), with four replicates per
 105 treatment, totalizing 24 experimental plots.

106 Damage probability data were submitted to multinomial distribution analysis. The
 107 percentage of infestation, data related to gas exchange and chlorophyll a fluorescence were
 108 subjected to variance analysis, and the means of the treatments were compared by the
 109 Tukey test at 5% probability, to evaluate both the effect of the treatments and the sensitivity
 110 differences between the cultivars to said treatments, all the variation factors were processed
 111 by SAS [22].

112 3. RESULTS AND DISCUSSION

113 For leaf damage caused by *Spodoptera frugiperda*, it was verified according to the
 114 evaluation periods (Figure 1A) that there was a statistical difference between the simple
 115 hybrid and Creole variety (Figure 1B). It was observed that in the interval of 14 to 42 days,
 116 the simple hybrid was the one that presented the highest probability of occurrence of score
 117 4, characterized by the stem of the partially destroyed plant, followed by score 3 that is the
 118 plant with symptoms of ripped leaf. In contrast, it was observed that from the second
 119 evaluation (14 days), the probability of occurrence of score 0, plants without symptoms of
 120 attack, was decreasing as a function of time, this characterizes an increase in the intensity of
 121 the attack of the pest.






Figure 1. Damage caused by *Spodoptera frugiperda* on the simple hybrid Genotype A (2A) and Creole variety Genotype B (2B) in the evaluation period.

For the cultivar Creole Genotype B in all the evaluation periods, there was a predominance of plants that did not present any type of damage, that is, a higher probability of occurrence of score 0 was observed, showing that this variety in the study conditions was larger degree of resistance than the simple hybrid. Little improved plants may present high levels of resistance, but tend to be less productive, which may justify the fact that the Creole variety has presented more plants with no damages when compared to the simple hybrid. Improved plants, such as the hybrid, invest little of their resources in synthesis of defenses against the attack of pest insects, because most of the times they are manipulated with the intention of achieving high productivity [23].

When evaluating 15 maize genotypes, 9 of which were Creole and 6 hybrids (1 Bt hybrid), found that 4 Creole genotypes presented moderate resistance to *S. frugiperda* in comparison to the other evaluated genotypes, based on visual observations [8].

There are different performances among maize cultivars in relation to *S. frugiperda* attacks, verified that higher damage scores were observed in hybrids when compared to their transgenic isogenic version [24]. [25] when studying different genotypes of maize against *S. frugiperda* attacks, found similar results that corroborate to this research, where the hybrids suffered a higher level of attack, when compared to the transgenic cultivar.

In terms of perennial infestation, there was a statistical difference between the evaluated cultivars (Table 2). The Creole Genotype B variety showed the lowest mean of *S. frugiperda* infestation, independent of the applied product, in relation to the simple hybrid Genotype A in all the evaluation periods. The highest percentage of infestation (76.0%) occurred at 28 days after application of the products for Genotype A while Genotype B at the same time presented an average percentage of infestation of 28.7%, which means a difference of 37.7% in the percentage of infestation between the two cultivars. This difference between the materials is also found in a study that evaluated the fertilization responses in different

" " ≠ to Creole cvs.

maize genotypes and the effect on *S. frugiperda* infestation. This difference between the materials is also found in a study that evaluated the fertilization responses in different maize genotypes and the effect on *S. frugiperda* infestation [26]

Table 2. Percentage of infestation of *Spodoptera frugiperda* in *Zea mays* submitted to treatments with Potassium Silicate and Calcium Chloride in five evaluation periods, Areia, 2017.

Cultivars	Treatments	<i>S. frugiperda</i> infestation (%) in five evaluation periods (days)					μ
		14	21	28	35	42	
Simple hybrid Genotype A	Si	79.8 Aa	74.6 Bc	76.0 Bb	69.7 Bd	54.9 Be	71.0 A
	Ca	77.8 Bb	76.5 Ad	83.4 Aa	76.9 Ac	60.0 Ae	74.9 A
	Test	67.07Cb	59.7 Cc	68.7 Ca	46.0 Cd	43.0 Gs	56.9 A
μ		74.87 A	70.3 A	76.0 A	64.2 A	52.6 A	-
Creole variety Genotype B	Si	16.31 Fd	15.2 Fc	23.8 Fb	40.2 Fa	17.8 Fe	22.7 A
	Ca	19.20 Ed	21.5 Ec	29.1 Ea	21.8 Eb	18.0 Ee	21.9 A
	Test	24.30 Dd	32.6 Db	33.3 Da	32.6 Dc	23.6 De	29.3 A
μ		19.93 B	23.1 B	28.7 B	31.5 B	19.8 B	-


¹Averages followed by the same capital letter in the column and lowercase in the row do not differ statistically from each other by Tukey's test at $P = 5$.
 μ =Infestation average

No statistical differences were observed in the mean percentage of infestation of *S. frugiperda* in plants where the products (Si and Ca) were applied (Table 2). This result differed from that found in another work with maize fertilization, which indicates that the types of fertilization possibly altered the response of the plant to the defense, in this work the sources of leaf fertilization did not alter any of the genotypes tested in this aspect [27]

In relation to the evaluation periods there was a tendency to reduce the percentage of infestation over time in all evaluated treatments. This observation may be related to the insect's own biology, which after its larval phase, around 15 days, leaves the plant to complete its cycle. Phenology of the plant may be another factor that influences the pest preference for younger tissues.

The results of the maximum quantum efficiency of FSII (F_V/F_M) were not statistically significant for the cultivars nor for the cultivars x treatments interaction (Si, Ca and control) (Figure 2). However, in this study, the presented results of the F_V/F_M ratio were close to the standard value for a plant with no damage to the photosynthetic apparatus, which is an approximate value of 0.8 [28]

control?
 treatments
 example? reference?






178
179

180 **Figure 2. Maximum quantum efficiency of PSII (F_v/F_m) in *Zea mays* submitted to**
 181 **treatments with Potassium Silicate, Calcium Chloride and control (without**
 182 **application), due to the stress caused by *Spodoptera frugiperda* attack.**

183 Evaluating the impact of infestation of *Coccus hesperidum* infestation (Linnaeus, 1758) on
 184 *Citrus limon*, ^{was} and found that the rates of F_v/F_m were significantly affected when the insect
 185 was fed from the plant [29].

186 For the stomatal conductance (g_s), a statistical difference was observed between the
 187 cultivars. In the Creole variety Genotype B, superior results were observed in relation to the
 188 simple hybrid Genotype A, specifically in silicon and control treatments, promoting to the
 189 hybrid a greater restriction to the stomatal opening. In addition, the Creole variety was
 190 different among the treatments, presenting a lower conductance in the treatment with
 191 calcium application (Figure 3A).



192

193

194

195

196

197

198

199

200

201

202

203

Figure 3. Evaluation of Stomatal Conductance (gs) (A); Photosynthesis (A) (B) and Transpiration (E) (C), in maize under the application of Potassium Silicate, Calcium Chloride and control. Values followed with the same lowercase letter did not differ among cultivars, and with the same capital letter they did not differ among the treatments of each cultivar by the Tukey test at 5% of probability.

The variable photosynthetic rate (A) did not present statistical difference among the evaluated cultivars. However, the treatments applied in the Creole variety were different, where a lower photosynthetic rate was observed when applied to the calcium source (Figure 3B). That order to compensate for the insect attack, in most cases the plant defense responses are associated with the reduction in the photosynthetic rate [30]

TF?
? and

between
?

Treatments are equal

just Calcium was $\neq$ to the control. $AB = B$

So maybe it is not effect of Silicon. Maybe it is effect of the cultivar.

204 Different transpiration behaviors (E) were observed among the cultivars, in the Creole variety
205 when submitted to silicon application and in the control, it was obtained a greater
206 transpiration compared to the simple hybrid (Figure 3C). This reduction may be related to the
207 results of stomatal conductance (Figure 3A). In the Creole variety, there was a significant
208 difference between the applied treatments, with lower transpiration when calcium was
209 applied (Figure 3C). These results differ from those found in a study with rice cultivation,
210 where increased transpiration under calcium fertilization was observed [31]

This is
? not true
AB = B

211 In this study, the results showed a significant reduction in the number of infestations (Table
212 2) and damage caused by *S. frugiperda*. The insect-plague in this one, causing greater
213 infestation and damage to the plants, being this similar result in work with cabbage and
214 beans that describes that the photosynthetic rate, the stomatal conductance and the
215 perspiration are significantly reduced under attack of plagues [32].

I loss!
What do
you mean?

216 Regarding the physiological parameters of Internal CO₂ Concentration (CI), Pressure Deficit
217 (DP), Water Efficiency (EFUA) and Carboxylation Efficiency (EC), no significant differences
218 were observed between the cultivars evaluated nor the treatments evaluated (Table 3). This
219 result may be related to the same cultural management and availability of water provided to
220 the cultivars, possibly not being these parameters sensitive or affected by the presence of
221 pests.

222 Table 3. Internal CO₂ Concentration (CI), Pressure Deficit (DP), Water Use Efficiency
223 (EFUA) and Carboxylation Efficiency (EC) in maize cultivars under the application of
224 potassium silicate, calcium chloride and control, Areia, 2017.
225

Cultivars	Parameters Evaluated			
	CI?	DP	EFUA	EC
Simple hybrid	72.52 a	2.48 a	4.57 a	0.26 a
Creole variety	68.37 a	2.40 a	4.36 a	0.38 a
Treatments				
Silicon	66.37 a	2.327 a	4.57 a	0.329 a
Calcium	76.75 a	2.528 a	4.44 a	0.219 a
Control	68.20 a	2.461 a	4.38 a	0.408 a
C.V. (%)	30.14	10.72	20.29	81.39

226 Means followed by the same lowercase letter in the column do not differ statistically from each other,
227 by Tukey's test at $P = .05$.

228 4. CONCLUSIONS

229 Silicon and Calcium sources do not promote the resistance induction to *S. frugiperda* in the
230 simple hybrid Genotype A and in the Creole variety Genotype B of maize.

231 The Creole variety of *Zea mays* suffers less damage by *Spodoptera frugiperda* infestation
232 than the simple hybrid Genotype A maize.

233

234 5. REFERENCES

- 235 1. USDA - United States Department of Agriculture. Grain: World Markets and Trade. 2017;
236 52.
- 237 2. CONAB - Companhia Nacional de Abastecimento. Acompanhamento de safra brasileira
238 de grãos. 2017; 4 (11): 1-171.
- 239 3. Nagoshi RN, Silvie P, Meagher LR, Lopez J, Machado V. Identification and comparison of
240 fall armyworm (Lepidoptera: Noctuidae) host strains in Brazil. Annals of the Entomological
241 Society of America. 2007; 100 (3): 394-402.
- 242 4. Nebo L, Matos AP, Vieira PC, Fernandes JB, Silva MFGF, Rodrigues RR. Atividade
243 inseticida dos frutos de *Trichilia clausenii* (Meliaceae) sobre *Spodoptera frugiperda*.
244 Química Nova. 2010; 33 (9): 1849-1852.
- 245 5. Carvalho RA, Omoto C, Field LM, Williamson MS, Bass C. Investigating the molecular
246 mechanism of organophosphate and pyrethroid resistance in the fall armyworm *Spodoptera*
247 *frugiperda*. Plos One. 2013; 8 (4): e62268.
- 248 6. Cruz I, Waquil JM. Pragas da cultura do milho para silagem. In: Cruz JC, Pereira FIA,
249 Rodrigues JAS, Ferreira JJ. Produção e utilização de silagem de milho e sorgo. Sete
250 Lagoas: Embrapa Milho e Sorgo. 2001; 41-207.
- 251 7. Lara FM. Princípios de Resistência de Plantas a Insetos, 1991; 2: 336.
- 252 8. Farias CA, Brewer MJ, Anderson DJ, Odvody GN; Xu W, Sétamou M. Native Maize
253 Resistance to Corn Earworm, *Helicoverpa zea*, and Fall Armyworm, *Spodoptera frugiperda*,
254 with Notes on Aflatoxin Content. Society of Southwestern Entomologists. 2014; 39 (2): 411-
255 426.
- 256 9. Gordy JW, Leonard BR, Blouin D, Davis JA, Stout MJ. Comparative effectiveness of
257 potential elicitors of plant resistance against *Spodoptera frugiperda* (J E Smith) (Lepidoptera:
258 Noctuidae) in four crop plants. Plos One. 2015; 10 (9): e0136689.
- 259 10. IPNI-International Plant Nutrition Institute. Nutri-Facts. Silicon, 2015; 14: 2.
- 260 11. Heckman J. Silicon: A Beneficial Substance, Better Crops. 2013; 97 (4): 14-16.
- 261 12. Liang Y, Nikolic M, Belanger R, Haijun G, Song A. Silicon in Agriculture. From Theory to
262 Practice. 1ª ed. Springer, Dordrecht, 2015.
- 263 13. Taiz L, Zeiger E, Moller IM, Murphy A. Fisiologia e desenvolvimento vegetal. 6ª ed
264 Artmed, Porto Alegre, 2017.
- 265 14. Firmino A, Abreu HS, Portugal ACP, Nascimento AM, Souza EL, Pereira RPW, Monteiro
266 MBO, Maêda JM. Alterações ligno-anatômicas em *Solanum gilo* Raddi por aplicação de
267 cálcio e boro como estratégia de defesa. Ciência Agrotécnica, Lavra. 2006; 30 (3).
- 268 15. Florentine SK, Raman A, Dhileepan K. Effects of gall induction by *Epiblema strenuana*
269 on gas exchange, nutrients, and energetics in *Parthenium hysterophorus*. BioControl,
270 Dordrecht. 2005; 50: 787-801.

- 271 16. Moore GM, Watts DA, Goolsby JA. Ecophysiological responses of giantreed (*Arundo*
272 *donax*) to herbivory. Invasive Plant science and management, Washington. 2010; 3: 521-
273 529.
- 274 17. Yusuf MA, Kumar D, Rajwanshi R, Strasser RJ, Tsimillimichael M, Govindje E, Sarin NB.
275 Overexpression of γ -totoopherol methyl transferase gene in transgenic *Brassica juncea* plants
276 alleviates abiotic stress: Physiological and chlorophyll a fluorescence measurements.
277 *Biochimica et Biophysica Acta*. Amsterdã. 2010; 1797: 1428-1438.
- 278 18. Peel MC, Finlayson BL, McMahon TA. Updated world map of the Köppen-Geiger climate
279 classification. *Hydrol. Earth System Science*, Bangalore. 2007; 11:1633-1644.
- 280 19. Embrapa - Empresa Brasileira de Pesquisa Agropecuária. Centro Nacional de Pesquisa
281 de Solos. Sistema brasileiro de classificação de solos. 2ª ed. Brasília, 2006.
- 282 20. Carvalho RPL. Danos, flutuação da população, controle e comportamento de
283 *Spodoptera frugiperda* (J. E. Smith, 1797) e suscetibilidade de diferentes genótipos de
284 milho, em condições de campo. Tese de Doutorado. Escola Superior de Agricultura Luiz de
285 Queiroz. 1970: 170.
- 286 21. Rousseau C, Belin E, Bove E, Rousseau D, Fabre F, Berruyer R, Guillaumés J,
287 Mangsau C, Jacques MA, Boureau T. High throughput quantitative phenotyping of plant
288 resistance using chlorophyll fluorescence image analysis. *Plant Methods*. 2013; 9 (17).
- 289 22. Cody R. Na introduction to SAS, University Edition, Cary, N. C., Statical Analysis System
290 Instituto, 2015; 366.
- 291 23. Agrawal AA, Hastings AP, Johnson MTJ, Maron JL, Salminen JP. Insect herbivores drive
292 real-time ecological and evolutionary change in plant populations. *Science*. 2012; 338
293 (6103): 113-116.
- 294 24. Moraes ARA, Lourenção AL, Paterniani MEAGZ. Resistência de híbridos de milho
295 convencionais e isogênicos transgênicos a *Spodoptera frugiperda* (Lepidoptera: Noctuidae).
296 *Bragantia*. 2015; 74 (1): 50-57.
- 297 25. Toscano LC, Fernandes MA, Rota MS, Maruyama WI, Andrade JV. Híbridos de milho
298 frente ao ataque de *Spodoptera frugiperda* em associação com adubação silicatada e o
299 efeito sobre o predador *Doru luteipes*. *Revista de Agricultura Neotropical*. 2016; 3 (1): 51-55.
- 300 26. Boiça JAL., Fernandes EB, Toscano LC, Lara FM. Influência de genótipos de milho,
301 adubação e inseticida sobre a população e danos de *Spodoptera frugiperda* (J.E. Smith,
302 1797) em duas épocas de semeadura. *Acta Scientiarum*. 2001; 23 (5): 1185-1190.
- 303 27. Roel AR, Soares JAL, Peruca RD, Pereira LC, Jadoski CJ. Occurrence in field and
304 development lab *Spodoptera frugiperda* (J.E. Smith) (Noctuidae) in corn with organic
305 fertilizer and chemical. *Applied Research & Agrotechnology*. 2017; 10 (1): 67-73.
- 306 28. Kalaji MH, GUO P. Chlorophyll fluorescence: a useful tool in barley plant breeding
307 programs. In: Sanchez, A. Gutierrez, S^J (Org.) Photochemistry research progress. Nova
308 publishers. 2008; 439-463.

- 309 29. Golan K, Rubinowska K, Kmiec K, Kot I, Górska-drabik E, Lagowska BM. Impact of
310 scale insect infestation on the content of photosynthetic pigments and chlorophyll
311 fluorescence in two host plant species. *Arthropod Plant Interactions*. 2015; 9: 55-65.
- 312 30. Bilgin DD, Zavala JA, Clough SJ, Ort DR, Delucia EH. Biotic stress globally
313 downregulates photosynthesis genes. *Plant, Cell and Environment*. 2010; 33 (10): 1597-
314 1613.
- 315 31. Silva LS, Bohnen H. Produtividade e absorção de nutrientes pelo arroz cultivado em
316 solução nutritiva com diferentes níveis de silício e cálcio. *Revista Brasileira de Agrociência*.
317 2003; 9 (1): 49-52.
- 318 32. Velikova V, Salerno G, Frati F, Peri E, Conti E, Colazza S, Loreto F. Influence of feeding
319 and oviposition by phytophagous pentatomids on photosynthesis of herbaceous plants.
320 *Journal of Chemical Ecology*. 2010; 36: 629-641.