

Original Research Article

Resistance induction in *Zea mays* L. cultivars against *Spodoptera frugiperda* (Lepidoptera: Noctuidae)

ABSTRACT

The maize crop *Zea mays* L., faces serious limitations in cultivation due to attack by the *Spodoptera frugiperda*. The objective of this work was to evaluate the resistance induction in maize cultivars with *S. frugiperda* caterpillars due to the application of Silicon and Calcium sources. Two cultivars of maize, Genotype A simple hybrid and the Creole Genotype B variety were used. These cultivars were submitted to two treatments of foliar fertilization, calcium chloride and potassium silicate, and a treatment without the application of an elicitor. The evaluations of the damages caused by the *S. frugiperda* caterpillar were carried out weekly, from the seventh day after the first application of the products (Si and Ca). In the bolting phase (VT), physiological evaluations of gas exchanges and fluorescence of Chlorophyll "a" were carried out. For the Creole variety, there was a predominance of plants that did not present any type of damage. The Creole variety presented the lowest mean of infestation by *S. frugiperda*, regardless of the product applied. No statistical differences were observed between the products applied in each cultivar. The maximum quantum efficiency of FSI (F_v/F_m) was not statistically significant for the cultivars nor for the cultivars x treatments interaction. For stomatal conductance (gs), a statistical difference between cultivars was found. In the Creole genotype B variety, superior results were obtained in relation to the simple hybrid. The TF variable did not present statistical difference between the evaluated cultivars. Different behaviors were observed in the transpiration variable (E) among the cultivars. In the Creole genotype B variety when submitted to silicon application and in the control, a higher transpiration was obtained comparing to the simple hybrid. The sources of Silicon and Calcium do not promote the induction of resistance to *S. frugiperda* in the studied variables. The Creole variety Genotype B of *Zea mays* suffers minor damage due to infestation of *Spodoptera frugiperda* in relation to the Genotype A simple hybrid maize.

Keywords: control method, fall armyworm, corn

1. INTRODUCTION

Zea mays L. maize is one of the world's largest cereals grown with the United States, China and Brazil as the world's largest producers. In the export ranking, Brazil occupies the second position [1], the Central West and South being the main producing regions of this cereal [2].

Although the country is prominent in the world production, it still presents limitations in the cultivation, which implies low productivity mainly due to insect pest attacks. Among the insect complexes that infest the maize crop, the *Spodoptera frugiperda* (J. E. Smith, 1797) caterpillar is the main cause of damage and represents a high cost of control annually [3].

Comment [m1]: Italic

Comment [m2]: Italic

Comment [m3]: Italic

The fall armyworm becomes an important pest of this crop, both by reducing productivity and decreasing the quality of the final product, besides the difficulty of control [4]. The considerable economic losses caused by this pest cause farmers to intensify the use of chemical insecticides and consequently lead to the development of resistance to many insecticide groups, making their control even more difficult [5].

Different control methods are used to reduce populations of *S. frugiperda*, such as cultural, chemical and biological tactics [6]. Plant resistance is also easily incorporated into integrated pest management because it maintains the pest population below the level of economic damage without unbalance to the environment and has cumulative and persistent effect beyond the lowest cost of production [7].

Due to the different biochemical, morphological and physiological characteristics among the genotypes of the same species, it is possible that some plants suffer a greater or lesser impact of phytophagy [8]; [9]. Plant resistance to pests can also be obtained by induction, which is a temporary manifestation, where a plant becomes less favorable to insect attack due to a certain condition that affects its physiology [7]. Among the various induction techniques, fertilizers have been widely used, mainly micro and macronutrients.

Silicon (Si) is considered a beneficial element for plants [10], it acts on the reinforcement of the cell wall by depositing solid silica, making the plant more rigid, with better growth and yield, for suppressing diseases and conferring characteristics of resistance to insects [11]; [12]. Calcium (Ca) is a macronutrient that is involved in a large number of signaling pathways in plants, from symbiotic interactions, defense responses and various hormones, functioning as a secondary messenger in response to biotic and abiotic signals [13]. Calcium participates in lignin biosynthesis, which favors plant self-defense and can be used as an inducer of pest resistance [14].

Other biomorphological factors are also associated with plant defense in relation to insect infestation. As a consequence of the stress caused by pest injury, there are reductions in the physiological parameters such as photosynthetic rate, transpiration and stomatal conductance [15]; [16]. For this purpose, the gas exchange and fluorescence analysis of chlorophyll "a" are able to detect, with safety and reliability, stress and injury effects in the photosynthetic process caused by biotic or abiotic factors [17]. In view of the above, the objective of this work was to evaluate induction resistance in corn cultivars due to the application of Silicon and Calcium sources in the infestation of the *Spodoptera frugiperda* caterpillar.

2. MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY

The experiment was conducted at the Chã de Jardim experimental farm, belonging to the Agricultural Sciences Center (CCA) of the Federal University of Paraíba (UFPB), Areia-PB, Brazil. The climate of the region is classified as tropical humid, presenting average temperature, relative humidity and precipitation of 23°C, 80% and 1,400 mm annually, respectively [18]. The soil is classified as Yellow Latosol [19].

The simple hybrid corn denominated Genotype A and the Creole variety denominated Genotype B were used. These materials were submitted to two treatments of foliar fertilizations, being applied commercial sources of Calcium Chloride and Potassium Silicate, and a control treatment. Three applications of the products were carried out in the vegetative growth phase of the maize cultivars, specifically in the V3, V6 and V12 stages, following the recommended dosages for culture suggested by the manufacturer of 2 L ha⁻¹. The products

were always applied from 4pm, this being the most recommended time for loss reduction, and manually by a costal spray with a 20-liter capacity.

The plots were constituted by four rows of 5 m in length, where after the mechanical preparation of the soil with a harrowing, furrows were opened and manual sowing was done, allowing a spacing of 0.50 m between rows and 0.20 m between plants, totaling 100 plants per plot and a density of 100,000 plants/ha. The useful area of the plot was constituted by the two central rows, scattering 0.5 m of the border. The nutritional correction was carried out with the fertilization of macronutrients (NPK) in the foundation and cover, and soil acidity, with dolomitic limestone application, according to results of soil chemical analysis. Weed control was performed with manual weeding.

2.1 Evaluation of the damage and infestation of *Spodoptera frugiperda*

The evaluations of the damages caused by *S. frugiperda* were carried out weekly, from the seventh day after the first application of the products (Si and Ca), with this done at 7, 14, 21, 35 and 42 days, ending at the beginning of flowering. Ten plants per plot were randomly selected, with individual scores varying between 0 and 5 (Table 1) [20].

Table 1. Scale of damages caused by *Spodoptera frugiperda* in *Zea mays* plants [20].

Score	Damage description
0	No damage
1	Scraped leaves
2	Bored leaves
3	Ripped leaves
4	Stem damage
5	Stem destroyed

To evaluate the level of infestation of *S. frugiperda*, the two central rows of each plot were used, relating the number of plants attacked by the total number of plants in the crop line.

2.2 Evaluation of physiological characteristics

The Fluorescence of Chlorophyll a, was determined from a portable fluorometer, model OS-30p from Opti-Sciences®. Measurements were made on three plants per plot, between 9 and 12 hours, at 2 cm from the margin of the leaf, excluding its veins. To that end, one leaf per plant, randomly selected was adapted to the dark with the use of tweezers, for approximately 30 minutes. The maximum photochemical quantum FSMI (Fv/Fm) parameter, often evaluated by the maximum fluorescence rate of chlorophyll "a", was evaluated to compare healthy and damaged tissues due to biotic factors [21].

In the maize bolting (VT) phase, gas exchange and photosynthetic efficiency of maize leaves were evaluated using an Infrared Gas Analyzer (IRGA), LCpro-SD model from BioScientific®. Two plants were used per plot in the morning, between 9 am and 11 am,

sampling the first leaf opposite and below the ear. The following physiological parameters were evaluated: Transpiration (E), Internal CO₂ concentration in leaf (IC), Stomatal conductance (gs), Photosynthesis (A), Carboxylation efficiency (EC), Pressure Deficit (PD) and Water Use Efficiency of (WUE).

2.3 Statistical analysis


The experimental design was a randomized complete block (DBC), in a factorial scheme (2x3), two maize cultivars (genotype A and genotype B), two sources of resistance induction (Si and Ca) plus one control (without application), with four replicates per treatment, totaling 24 experimental plots.

Comment [m4]: RANDOMIZED COMPLETE BLOCK DESIGN (RCBD)

Damage probability data were submitted to multinomial distribution analysis. The percentage of infestation, data related to gas exchange and chlorophyll a fluorescence were subjected to variance analysis, and the means of the treatments were compared by the Tukey test at 5% probability, to evaluate both the effect of the treatments and the sensitivity differences between the cultivars to said treatments, all the variation factors were processed by SAS [22].

3. RESULTS AND DISCUSSION

For leaf damage caused by *Spodoptera frugiperda*, it was verified according to the evaluation periods (Figure 1 A) that there was a statistical difference between the simple hybrid and Creole variety (Figure 1 B). It was observed that in the interval of 14 to 42 days, the simple hybrid was the one that presented the highest probability of occurrence of score 4, characterized by the stem of the partially destroyed plant, followed by score 3 that is the plant with symptoms of ripped leaf. In contrast, it was observed that from the second evaluation (14 days), the probability of occurrence of score 0, plants without symptoms of attack, was decreasing as a function of time, this characterizes an increase in the intensity of the attack of the pest.






Figure 1. Damage caused by *Spodoptera frugiperda* on the simple hybrid Genotype A (1 A) and Creole variety Genotype B (1 B) in the evaluation period.

For the cultivar Creole Genotype B in all the evaluation periods, there was a predominance of plants that did not present any type of damage, that is, a higher probability of occurrence of score 0 was observed, showing that this variety in the study conditions was larger degree of resistance than the simple hybrid. Little improved plants may present high levels of resistance, but tend to be less productive, which may justify the fact that the Creole variety has presented more plants with no damages when compared to the simple hybrid. Improved plants, such as the hybrid, invest little of their resources in synthesis of defenses against the attack of pest insects, because most of the times they are manipulated with the intention of achieving high productivity [23].

When evaluating 15 maize genotypes, 9 of which were Creole and 6 hybrids (1 Bt hybrid), found that 4 Creole genotypes presented moderate resistance to *S. frugiperda* in comparison to the other evaluated genotypes, based on visual observations [8].

There are different performances among maize cultivars in relation to *S. frugiperda* attacks, verified that higher damage scores were observed in hybrids when compared to their transgenic isogenic version [24]. When studying different genotypes of maize against *S. frugiperda* attacks, found similar results that corroborate to this research, where the hybrids suffered a higher level of attack, when compared to the transgenic cultivar [25].

In terms of perennial infestation, there was a statistical difference between the evaluated cultivars (Table 2). The Creole Genotype B variety showed the lowest mean of *S. frugiperda* infestation, independent of the applied product, in relation to the simple hybrid Genotype A in all the evaluation periods. The highest percentage of infestation (76.0%) occurred at 28 days after application of the products for Genotype A while Genotype B at the same time presented an average percentage of infestation of 28.7%, which means a difference of

37.7% in the percentage of infestation between the two cultivars. This difference between the materials is also found in a study that evaluated the fertilization responses in different maize genotypes and the effect on *S. frugiperda* infestation [26]

Table 2. Percentage of infestation of *Spodoptera frugiperda* in *Zea mays* submitted to treatments with Potassium Silicate and Calcium Chloride in five evaluation periods, Areia, 2017.

Cultivars	Treatments	<i>S. frugiperda</i> infestation (%) in five evaluation periods (days)					μ
		14	21	28	35	42	
Simple hybrid Genotype A	Si	79.8 Aa	74.6 Bc	76.0 Bb	69.7 Bd	54.9 Be	71.0 A
	Ca	77.8 Bb	76.5 Ad	83.4 Aa	76.9 Ac	60.0 Ae	74.9 A
	Control	67.07Cb	59.7 Cc	68.7 Ca	46.0 Cd	43.0 Gs	56.9 A
μ		74.87 A	70.3 A	76.0 A	64.2 A	52.6 A	-
Creole variety Genotype B	Si	16.31 Fd	15.2 Fc	23.8 Fb	40.2 Fa	17.8 Fe	22.7 A
	Ca	19.20 Ed	21.5 Ec	29.1 Ea	21.8 Eb	18.0 Ee	21.9 A
	Control	24.30 Dd	32.6 Db	33.3 Da	32.6 Dc	23.6 De	29.3 A
μ		19.93 B	23.1 B	28.7 B	31.5 B	19.8 B	-

¹Averages followed by the same capital letter in the column and lowercase in the row do not differ statistically from each other by Tukey's test at $P = 0.05$.
 μ =Infestation average

No statistical differences were observed in the mean percentage of infestation of *S. frugiperda* in plants where the treatments (Si and Ca) were applied (Table 2). This result differed from that found in another work with maize fertilization, which indicates that the types of fertilization possibly altered the response of the plant to the defense, in this work the sources of leaf fertilization did not alter any of the genotypes tested in this aspect [27].

In relation to the evaluation periods there was a tendency to reduce the percentage of infestation over time in all evaluated treatments. This observation may be related to the insect's own biology, which after its larval phase, around 15 days, leaves the plant to complete its cycle. Phenology of the plant may be another factor that influences the pest preference for younger tissues.

The results of the maximum quantum efficiency of FSII (F_v/F_m) were not statistically significant for the cultivars nor for the cultivars x treatments interaction (Si, Ca and control) (Figure 2). However, in this study, the presented results of the F_v/F_m ratio were close to the standard value for a plant with no damage to the photosynthetic apparatus, which is an approximate value of 0.8 [28]






Figure 2. Maximum quantum efficiency of FSII (F_v/F_m) in *Zea mays* submitted to treatments with Potassium Silicate, Calcium Chloride and control (without application), due to the stress caused by *Spodoptera frugiperda* attack.

Evaluating the impact of infestation of *Coccus hesperidum* infestation (Linnaeus, 1758) on *Citrus limon*, was found that the rates of F_v/F_m were significantly affected when the insect was fed from the plant [29].

For the stomatal conductance (gs), a statistical difference was observed between the cultivars. In the Creole variety Genotype B, superior results were observed in relation to the simple hybrid Genotype A, specifically in silicon and control treatments, promoting to the hybrid a greater restriction to the stomatal opening. In addition, the Creole variety was different among the treatments, presenting a lower conductance in the treatment with calcium application (Figure 3A).





197

198 **Figure 3. Evaluation of Photosynthesis (A) (A); Stomatal Conductance (gs) (B) and**
 199 **Transpiration (E) (C), in maize under the application of Potassium Silicate, Calcium**
 200 **Chloride and control. Values followed with the same lowercase letter did not differ**
 201 **among cultivars, and with the same capital letter they did not differ among the**
 202 **treatments of each cultivar by the Tukey test at 5% of probability.**

203 The variable photosynthetic rate (A) did not present statistical difference among the cultivars
 204 evaluated. However, the treatment with calcium application was different from the control in
 205 the Creole variety, where a lower photosynthetic rate was observed when applied to the
 206 calcium source (Figure 3B). This order to compensate for insect attack, in most cases, plant
 207 defense responses are associated with the reduction in photosynthetic rate [30].

208 Different transpiration behaviors (E) were observed among the cultivars, in the Creole variety
 209 when submitted to silicon application and in the control, it was obtained a greater
 210 transpiration compared to the simple hybrid (Figure 3C). This reduction may be related to the
 211 results of stomatal conductance (Figure 3A). In the Creole variety, there was a significant
 212 difference between the applied treatments, with lower transpiration when calcium was
 213 applied (Figure 3C). These results differ from those found in a study with rice cultivation,
 214 where increased transpiration under calcium fertilization was observed [31]

215 In this study, the results showed a significant reduction in the percentage of infestations
 216 (Table 2) and damages caused by *S. frugiperda*. A similar result was observed in the work

with cabbage and beans which describes that the photosynthetic rate, stomatal conductance and transpiration are significantly reduced under pest attack [32].

Regarding the physiological parameters of Internal CO₂ Concentration (IC), Pressure Deficit (PD), Water Use Efficiency (WUE) and Carboxylation Efficiency (CE), no significant differences were observed between the cultivars evaluated nor the treatments evaluated (Table 3). This result may be related to the same cultural management and availability of water provided to the cultivars, possibly not being these parameters sensitive or affected by the presence of pests.

Table 3. Internal CO₂ Concentration (IC), Pressure Deficit (PD), Water Use Efficiency (WUE) and Carboxylation Efficiency (CE) in maize cultivars under the application of potassium silicate, calcium chloride and control, Areia, 2017.

Cultivars	Parameters Evaluated			
	IC	PD	WUE	CE
Simple hybrid	72.52 a	2.48 a	4.57 a	0.26 a
Creole variety	68.37 a	2.40 a	4.36 a	0.38 a
Treatments				
Silicon	66.37 a	2.327 a	4.57 a	0.329 a
Calcium	76.75 a	2.528 a	4.44 a	0.219 a
Control	68.20 a	2.461 a	4.38 a	0.408 a
C.V. (%)	30.14	10.72	20.29	81.39

Means followed by the same letter in the column do not differ statistically from each other, by Tukey's test at $P = 0.05$.

4. CONCLUSIONS

Silicon and Calcium sources do not promote the resistance induction to *S. frugiperda* attack in the simple hybrid Genotype A and in the Creole variety Genotype B of maize.

The Creole variety of *Zea mays* suffers less damage by *Spodoptera frugiperda* infestation than the simple hybrid Genotype A maize.

5. REFERENCES

1. USDA - United States Department of Agriculture. Grain: World Markets and Trade. 2017;52. Accessed 28 April 2018. Available: <https://www.fas.usda.gov/data/grain-world-markets-and-trade>.

2. CONAB - Companhia Nacional de Abastecimento. Acompanhamento de safra brasileira de grãos. 2017;4(11):1-171.

3. Nagoshi RN, Silvie P, Meagher LR, Lopez J, Machado V. Identification and comparison of fall armyworm (Lepidoptera: Noctuidae) host strains in Brazil. Annals of the Entomological Society of America. 2007;100(3):394-402.

Comment [m5]: do not promote or induction the resistance towards *S.frugiperda* attack

- 245 4. Nebo L, Matos AP, Vieira PC, Fernandes JB, Silva MFGF, Rodrigues RR. Atividade
246 inseticida dos frutos de *Trichilia clausenii* (Meliaceae) sobre *Spodoptera frugiperda*.
247 Química Nova. 2010;33(9):1849-1852.
- 248 5. Carvalho RA, Omoto C, Field LM, Williamson MS, Bass C. Investigating the molecular
249 mechanism of organophosphate and pyrethroid resistance in the fall armyworm *Spodoptera*
250 *frugiperda*. Plos One. 2013;8(4):1-11.
- 251 6. Cruz I, Waquil JM. Pragas da cultura do milho para silagem. In: Cruz JC, Pereira FIA,
252 Rodrigues JAS, Ferreira JJ. Produção e utilização de silagem de milho e sorgo. Sete
253 Lagoas: Embrapa Milho e Sorgo. 2001;41-207.
- 254 7. Lara FM. Princípios de Resistência de Plantas a Insetos. 1ª ed. 1991:336.
- 255 8. Farias CA, Brewer MJ, Anderson DJ, Odvody GN; Xu W, Sétamou M. Native Maize
256 Resistance to Corn Earworm, *Helicoverpa zea*, and Fall Armyworm, *Spodoptera frugiperda*,
257 with Notes on Aflatoxin Content. Society of Southwestern Entomologists. 2014; 39 (2): 411-
258 426.
- 259 9. Gordy JW, Leonard BR, Blouin D, Davis JA, Stout MJ. Comparative effectiveness of
260 potential elicitors of plant resistance against *Spodoptera frugiperda* (J E Smith) (Lepidoptera:
261 Noctuidae) in four crop plants. Plos One. 2015;10(9):1-14.
- 262 10. IPNI-International Plant Nutrition Institute. Nutri-Facts. Silicon. 2015;14:2. Accessed 12
263 May 2018. Available: <http://www.ipni.net/nutrifacts-northamerican>.
- 264 11. Heckman J. Silicon: A Beneficial Substance, Better Crops. 2013;97(4):14-16.
- 265 12. Liang Y, Nikolic M, Belanger R, Haijun G, Song A. Silicon in Agriculture. From Theory to
266 Practice. 1ª ed. Springer, Dordrecht; 2015.
- 267 13. Taiz L, Zeiger E, Moller IM, Murphy A. Fisiologia e desenvolvimento vegetal. 6ª ed.
268 Artmed, Porto Alegre; 2017.
- 269 14. Firmino A, Abreu HS, Portugal ACP, Nascimento AM, Souza EL, Pereira RPW, Monteiro
270 MBO, Maêda JM. Alterações ligno-anatômicas em *Solanum gilo* Raddi por aplicação de
271 cálcio e boro como estratégia de defesa. Ciência Agrotécnica, Lavras. 2006;30(3):394-401.
- 272 15. Florentine SK, Raman A, Dhileepan K. Effects of gall induction by *Epiblema strenuana*
273 on gas exchange, nutrients, and energetics in *Parthenium hysterophorus*. BioControl,
274 Dordrecht. 2005;50:787-801.
- 275 16. Moore GM, Watts DA, Goolsby JA. Ecophysiological responses of giantreed (*Arundo*
276 *donax*) to herbivory. Invasive Plant science and management, Washington. 2010;3:521-529.
- 277 17. Yusuf MA, Kumar D, Rajwanshi R, Strasser RJ, Tsimilimichael M, Govindje E, Sarin NB.
278 Overexpression of γ-totopherol methyl transferase gene in transgenic *Brassica juncea* plants
279 alleviates abiotic stress: Physiological and chlorophyll a fluorescence measurements.
280 Biochimica et Biophysica Acta. Amsterdã. 2010;1797:1428-1438.
- 281 18. Peel MC, Finlayson BL, McMahon TA. Updated world map of the Köppen-Geiger climate
282 classification. Hydrol. Earth System Science, Bangalore. 2007;11:1633-1644.

- 283 19. Embrapa - Empresa Brasileira de Pesquisa Agropecuária. Centro Nacional de Pesquisa
284 de Solos. Sistema brasileiro de classificação de solos. 2ª ed. Brasília. 2006: 286.
- 285 20. Carvalho RPL. Danos, flutuação da população, controle e comportamento de
286 *Spodoptera frugiperda* (J. E. Smith, 1797) e suscetibilidade de diferentes genótipos de
287 milho, em condições de campo. Tese de Doutorado. Escola Superior de Agricultura Luiz de
288 Queiroz.1970:170.
- 289 21. Rousseau C, Belin E, Bove E, Rousseau D, Fabre F, Berruyer R, Guillaumés J,
290 Mangsau C, Jacques MA, Boureau T. High throughput quantitative phenotyping of plant
291 resistance using chlorophyll fluorescence image analysis. *Plant Methods*. 2013;9(17):1-13.
- 292 22. Cody R. Na introduction to SAS, University Edition, Cary, N. C., Statical Analysis System
293 Instituto, 2015;366.
- 294 23. Agrawal AA, Hastings AP, Johnson MTJ, Maron JL, Salminen JP. Insect herbivores drive
295 real-time ecological and evolutionary change in plant populations. *Science*. 2012;338
296 (6103):113-116.
- 297 24. Moraes ARA, Lourenção AL, Paterniani MEAGZ. Resistência de híbridos de milho
298 convencionais e isogênicos transgênicos a *Spodoptera frugiperda* (Lepidoptera: Noctuidae).
299 *Bragantia*. 2015;74(1):50-57.
- 300 25. Toscano LC, Fernandes MA, Rota MS, Maruyama WI, Andrade JV. Híbridos de milho
301 frente ao ataque de *Spodoptera frugiperda* em associação com adubação silicatada e o
302 efeito sobre o predador *Doru luteipes*. *Revista de Agricultura Neotropical*. 2016;3(1):51-55.
- 303 26. Boiça JAL.; Fernandes EB, Toscano LC, Lara FM. Influência de genótipos de milho,
304 adubação e inseticida sobre a população e danos de *Spodoptera frugiperda* (J.E. Smith,
305 1797) em duas épocas de semeadura. *Acta Scientiarum*. 2001;23(5):1185-1190.
- 306 27. Roel AR, Soares JAL, Peruca RD, Pereira LC, Jadoski CJ. Occurrence in field and
307 development lab *Spodoptera frugiperda* (J.E. Smith) (Noctuidae) in corn with organic
308 fertilizer and chemical. *Applied Research & Agrotechnology*. 2017;10(1):67-73.
- 309 28. Kalaji MH, GUO P. Chlorophyll fluorescence: a useful tool in barley plant breeding
310 programs. In: Sanchez, A. Gutierrez, S.J (Org.) *Photochemistry research progress*. Nova
311 publishers. 2008;439-463.
- 312 29. Golan K, Rubinowska K, Kmiec K, Kot I, Górska-drabik E, Lagowska BM. Impact of
313 scale insect infestation on the content of photosynthetic pigments and chlorophyll
314 fluorescence in two host plant species. *Arthropod Plant Interactions*. 2015;9:55-65.
- 315 30. Bilgin DD, Zavala JA, Clough SJ, Ort DR, Delucia EH. Biotic stress globally
316 downregulates photosynthesis genes. *Plant, Gsll and Environment*. 2010;33(10):1597-1613.
- 317 31. Silva LS, Bohnen H. Produtividade e absorção de nutrientes pelo arroz cultivado em
318 solução nutritiva com diferentes níveis de silício e cálcio. *Revista Brasileira de Agrociência*.
319 2003;9(1):49-52.
- 320 32. Velikova V, Salerno G, Frati F, Peri E, Conti E, Colazza S, Loreto F. Influence of feeding
321 and oviposition by phytophagous pentatomids on photosynthesis of herbaceous plants.
322 *Journal of Chemical Ecology*. 2010;36:629-641.