

Original Research Article

Initial development of sunflower under weed competition with different species of grasses

ABSTRACT: Introduction: All cultivated plants are subject to intra and interspecific grass competition, once this competition may gradually increase within the crop cycle. **Objective:** In view of the above, this work aims to evaluate the initial development of sunflower, when submitted to competition with different species of grasses. **Material and methods:** A randomized block design was installed in a 2x4 factorial scheme, with two grass species, *Urochloa decumbens* cv. Basilisk and *Panicum maximum* cv. Mombasa, with four densities of competition with grasses, plus a control group (absence of grass) and four replications, total of 36 plots. **Discussion:** During the sowing of the grasses, the following densities were considered: low; average; high and very high density of competition. While there was an increase in density of competition, sunflower development was impaired. **Conclusion:** The initial development of the sunflower was impaired under a 49.06 cm² competition per weed and the *Panicum maximum* species caused greater damage to the crop.

Keywords: *Urochloa decumbens*; *Panicum maximum*; weed; stress

Introduction

The sunflower, due to the diversity of physical-chemical features of its by-products, has become an attractive option for the Brazilian agribusiness, due to its adaptability to different growing conditions [1,2]. However, all cultivated plants are subject to intra and interspecific grass competition, since this competition may gradually increase with the course of the crop cycle [3,4].

Inherent or weedy plants can influence crop yield decline through competition for water, light, nutrients and space, or even for the allelopathic action of some species [5]. This competition may affect the quality of the raw

Comment [s1]: D

Comment [s2]: S

Comment [s3]: W

Comment [s4]: C

Comment [s5]: D

Comment [s6]: S

Comment [s7]: G

Comment [s8]: BOLD

Comment [s9]: Results and

Comment [s10]: W

Comment [s11]: S

material, elevate the cost of producing the crop and can be a host of pests and diseases [6].

At the beginning of the productive cycle the invasive plants can coexist for a time without damage to the development of the cultivated crop, however, during the course of its phases it is possible to observe the damages caused by the grass competition, which makes necessary to understand the effects caused by this competition in order to set an ideal time for the control of invasive plants, mainly grass species, making necessary the knowledge of the period before weed interference (PBI) [7,8].

The damaging effect of grass competition on sunflower crop in the first 30 days after emergence is notorious due to its slow growth as compared to weeds, mainly grass species, which can vary from 23 to 70% of losses in the production. Some sunflowers cultivars showed a variation in PAI from 16 to 24 days after germination, in which a negative interference was verified in the stem diameter, grain yield and oil yield [9].

These negative responses to grass competition can be a reflection of the morpho-physiological changes in the cultivated plant, making necessary to understand these changes that can cause a greater foliar transpiration or even the reduction of the photosynthetic rate due to competition, which entails in a smaller accumulation of mass dry [4,10,11,12].

In view of the above, this work aims to evaluate the development of the sunflower, when submitted to competition with different grass species.

Material and methods

Comment [s12]: Methods

The experiment was carried out during the month of August 2018, at the Andradina Educational Foundation (FEA), located in the city of Andradina, state of São Paulo, Brazil. A randomized block design was installed in a 2x4 factorial scheme, two grass species, that is, *Urochloa decumbens* cv. Basilisk and *Panicum maximum* cv. Mombasa with four densities of competition with grasses, plus a control group, with absence of grass, and four replications, total of 36 plots.

Sunflowers seeds variety Agobel 960 was sowed at 3 cm deep in pots with a 7 dm³ capacity and 490.6 cm² area, filled with soil originated from the 0 –

0.3 layer with chemical attributes according to Table 1. The soil was fertilized with urea – 0.0777 g; triple superphosphate – 1.36 g; and potassium chloride – 0.36 g per pot [13].

Comment [s13]: g./pot.

Table 1: Chemical attributes of the soil.

Table 1. Chemical attributes of the soil.																
pH	MO	P	K	Ca	Mg	H+Al	Al	SB	CTC	V%	m%	B	Cu	Fe	Mn	Zn
CaCl ₂	g dm ⁻³	mg dm ⁻³	mmol _c dm ⁻³					mg dm ⁻³								
5.9	11	21	2.0	19	7	15	0	28	43	65	0	0.21	1	25	9	3.4

SB: Sum of bases; V%: Base saturation; m%: Aluminum saturation.

At the same time the grasses were sown and the following densities were considered: absence of the grass: only the sowing of the sunflower; low density of competition with grasses, where the presence of two grassy plants per pot was considered, providing 245.3 cm² per plant. For the medium density competing conditions, the presence of four grassy plants per pot was considered, providing 122.65 cm² per plant. In conditions of high competition density, the presence of eight grassy plants per pot was considered, being 61.32 cm² per plant. And for the conditions of very high competition density, the presence of sixteen grassy plants per pot was considered, being 30.66 cm² per plant. During the experiment, the vessels were irrigated whenever necessary, respecting the field capacity.

Comment [s14]: and

Comment [s15]: cm²/plant

Comment [s16]: cm²/ plant

Thirty days after the planting, the following parameters were set: PHS and PHG - plant height of sunflower and plant height of grass, determined by using a millimeter ruler; NLS – number of leaves of the sunflower determined by the direct count in the plant; DMS and DMAPG – dry mass of sunflower and dry mass of aerial part of grass; DMSR and DMGR – dry mass of sunflower root and dry mass grass root, set by drying the wet mass in a circulation oven and renewing air at a constant temperature of 65 ° C until reaching constant weight.

The impression was also made on the inferior or abaxial epidermal surface of the fragments collected using cyanoacrylate ester, for the determination functionality of the stomatal inferior or abaxial surface (SFS) and stomatal density of the lower or abaxial surface (SDS) [14,15]. For all the characteristics 10 measurements per slide were performed. The plots were represented by the average value obtained from the measurements of each characteristic.

All variables were submitted to the F test ($p < 0.05$) and the regression analysis was applied to the competition densities, where their models were tested: linear; quadratic and cubic, whereas for the grass species the Tukey test was applied at 5% probability [16]. The statistical program Assistat 7.7 was used [17].

Comment [s17]: Assistant

Results

Sunflower

U. decumbens species caused a greater reduction in the height of the sunflower as shown in Table 2.

Table 2: Mean values of plant height (PH); number of leaves (NL); dry mass of aerial part of sunflower and root (DMAPS and DMRS); stomatal sunflower density (SDS) and stomatal sunflower functionality (SSF) as submitted to competition with different grass densities.

	PH (cm)	NL	DMAPS (g)	DMRS (g)	SDS (n°S/mm ²)	SSF
<i>U. decumbens</i>	42.45b	15.45a	6.66a	2.28a	240.62a	3.17a
<i>P. maximum</i>	46.68a	15.80a	5.56b	1.67a	241.87a	3.02a
DMS	3.46	0.96	1.02	0.95	28.17	0.37
CV%	11.99	9.52	25.93	74.72	18.00	18.86
MG	44.56	15.62	6.11	1.97	241.25	3.09
F	6.26*	0.55Ns	4.81*	1.71Ns	0.01ns	0.68Ns

DMS: Minimum significant difference. CV: Coefficient of variation. MG: Overall mean. f: value of F calculated in the analysis of variance; Nsp = 0.05; * 0.01 = $p < 0.05$; ** $p < 0.01$. The averages in the column followed by the same letter do not differ statistically from each other. The Tukey test was applied at a 5% probability level.

To the number of leaves of the sunflower it was not observed a statistical difference between the grass species, whereas to the dry mass of the aerial part, *Panicum maximum* caused a approximate 16%-reduction; but in the dry mass of the root it was not observed a significant influence in the competition between grass species. In the same way, there was no significant effect between the influences of forage species competitions on stomatal density and functionality.

Comment [s18]: an

Table 3 shows the analysis of variance of the regressions of the sunflower as submitted to different densities of competition among grass species.

Tabela 3: The analysis of variance of the regressions of the sunflower as submitted to different grass competition densities, the following models were tested: linear, quadratic and cubic.

Middle Square								
Forage	FV	GL	PH (cm)	NL	DMAPS (g)	DMRS (g)	SDS (n°S/mm ²)	SFS
<i>U. decumbens</i>	Density	4	137.786	29.756	24.016	15.508	13819.982	2.685
	Residue	16	23.284	3.254	4.032	2.448	1194.041	0.506
	Regression	1	Q*	L**	Q*	Q*	Q**	L*
<i>P. maximum</i>	Density	4	324.542	29.615	19.357	24.085	55316.406	1.570
	Residue	16	64.185	2.326	1.032	2.317	2183.690	0.256
	Regression	1	Q*	Q*	Q*	Q*	L**	Q*

Ns- $p \geq 0.05$; * $0.01 \leq p < 0.05$; ** $p < 0.01$. L: polynomial of 1st degree. Q: polynomial of 2nd degree. PH – plant height; NL – number leaf; DMAPS – Dry mass of the air part of the sunflower; SDS – Stomata density of the sunflower and SFS – Stomata functionality of the sunflower.

The Sunflower in competition with *Urochloa decumbens* presented a constant increase in its height till reached the maximum point of approximately nine plants, after this point a drop occurred in its development, similarly to competition with *Panicum maximum*, also presented a quadratic response, but its peak approximately occurred until six plants, after this density there was a reduction in its growth as Figure 1 shows.




Figure 1: Sunflower plant height as submitted to different densities of

competition among grass species. S= Sunflower; D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

Regarding the number of sunflower leaves, a negative linear response was observed by increasing *Urochloa decumbens* plant density. However, the sunflower presented a quadratic negative response to the competition with *Panicum maximum*, in which there was the minimum point of leaves until the competition of approximately eight plants as Figure 2 shows.




Figure 2: Number of sunflower leaves as submitted to different densities of competition between grass species. S= Sunflower; D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

A quadratic response was observed to dry mass of aerial part of sunflower, in which a minimum of ten and eleven plants were respectively observed for *Urochloa decumbens* and *Panicum maximum*, as Figure 3 shows.




Figure 3: Figure 3: Dry mass of leaves of sunflower as submitted to different densities of grass competitions. S= Sunflower; D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

Similarly, it was possible to observe that the dry mass of the sunflower root presented a quadratic negative response as the density of grassy plants increased, which presented a minimum point of ten plants regardless of grass species, as Figure 4 displays.




Figure 4: Dry mass of root of sunflower (DMRS) as submitted to different densities of grass competitions. S= Sunflower; D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

145

146 According to Figure 5, the stomatal density of the abaxial face of the
 147 sunflower showed a quadratic response under competition with the *Urochloa*
 148 *decumbens* species presenting a minimum reduction up to ten plants, whereas
 149 for the *Panicum maximum* species a negative linear response was observed
 150 whereas there was an increase in competition density.

151





Figure 5: Stomata density of sunflower (SDS) when submitted to different densities of grass competitions. S= Sunflower; D= *Urochloa decumbens* and M = *Panicum maximum*. Andradina, 2018.

152

153 A difference in the size and number of stomata on the abaxial face of the
 154 sunflower is noticeable when there was an increase in competition density with
 155 grass species as observed in Figure 6.

156






Figure 6: Stomata of the abaxial sunflower leaf when submitted to different densities of grass competitions. A - Absence of competition; B - Two plants of *Urochloa decumbens*; C - Four plants of *Urochloa decumbens*; D - Eight plants of *Urochloa decumbens*; E - Sixteen plants of *Urochloa decumbens*; F - Two *Panicum maximum* plants; G - Four plants of *Panicum maximum*; H - Eight plants of *Panicum maximum* and I - Sixteen plants of *Panicum maximum*. PD – Polar diameter; ED – Equatorial diameter.

157

158 The competition with the *Urochloa decumbens* species caused a
159 negative linear response in the stomata functionality of the sunflower, in which a
160 quadratic response was observed that presented the minimum point with eight
161 grassy plants in the competition, as Figure 7 shows.

162




Figure 7: Stomata functionality of sunflower (SFS) as submitted to different densities of grass competitions. S= Sunflower; D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

Grass

A linear response was observed at both grass heights according to Table 4.

Table 4: The analysis of variance of the regressions of the grass in different densities competition, where the models were tested: linear, quadratic and cubic.

Middle Square					
Forage	FV	GL	PHG (cm)	DMAPG (g)	DMRG (g)
<i>U. decumbens</i>	Density	4	226.802	39.326	18.043
	Residue	12	28.083	0.662	2.448
	Regression	1	L*	L*	L*
<i>P. maximum</i>	Density	4	660.002	10.182	21.638
	Residue	12	18.291	0.315	2.317
	Regression	1	L**	L*	L**

Ns- $p \geq 0.05$; $0.01 \leq p < 0.05$; ** $p < 0.01$. L: polynomial of 1st degree. Q: polynomial of 2nd degree. PH – plant height; NL – number leaf; DMAP – Dry mass of the air part; S: Stomata; SD – Stomata density and SF – Stomata functionality.

However their responses were antagonistic, once *Urochloa decumbens* presented a positive response to the increase of the number of grassy plants, while the *Panicum maximum* species showed a negative response as Figure 8 shows.




Figure 8: Plant height of grasses under different densities of competitions. D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

According to Figure 9, as the grass density increased, a linear response in dry mass in the aerial part occurred, regardless of the grass species.




Figure 9: Dry mass of aerial part of grass in different density of competition. D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

Similarly, dry mass of grass root presented a positive response as the density of grassy plants increased, as figure 10 shows.

Comment [s19]: Figure




Figure 10: Dry mass of root of grass in different density of competition. D= *Urochloa decumbens* and M= *Panicum maximum*. Andradina, 2018.

Discussion

The constant growth of Sunflower height did not occur due to the greater availability of nutrients and light, but rather, it triggered survival ability competition through its sowing. For the increase in height when in competition, it represents an adaptation to a future competition with neighboring plants, as it is a result of processes of the hormonal dynamics and cell division more accentuated in the search for light [18].

Thus, plants that are in free competition use a good part of their photoassimilates for root development, but, under competition the plant redirects these photoassimilates to growth of aerial part in order to gain more leaf area to minimize this competition effect [10,19]. Plants that are submitted to competition for light and are exposed to shading, a change in their morphological architecture causes an elongation of stem and leaf petioles [20].

The interaction caused by the competition results in a lesser number of leaves due to the decrease of photoassimilate, entailing a lower dry mass [21,10]. The intensity of weed interference on the crop of economic interest may lead to a decrease in its development, since grasses presented negative and positive linear responses, it is possible to observe that this occurred due to the natural difference between species [13]. *Urochloa decumbens* presents a lower nutritional requirement and more persistent and adaptive action in the

environmental variations that is inserted, making it more tolerant to these variations. However, the *Panicum maximum* species extracts more nutrients from the soil, which impairs the development of the sunflower crop, in case of competition between them.

The competition for water, light and nutrients is a crucial factor for crops, if there is intense growth of the aerial part of the crop, a greater availability of water and nutrients from the soil is required, and the increase of dry matter of the root is directly related to this availability, so that, the competition with grassy plants entailed a reduction in the root dry mass of the sunflower. Other species, such as cotton, when submitted to competition with grasses suffered interference in its development, mainly in dry mass of aerial part and root, once the species competed for light, as this interference decreased their photosynthetic rate, leading to reduction in the accumulation of dry matter [20].

However, it is important to note that necessary a better understanding of the genetic factors involved in the development of this species [22,23]. In the aerial part, as well as in the root, the competition with the sunflower crop did not interfere in the development of the grasses, since both presented a positive linear response regarding the dry mass conteny, this result was already expected, since there was an increase in the number of plants by area.

Density and stomata functionality presented negative responses as the grass competition density increased, since the presence of weeds in the production environment can cause damage to the crop mainly due to the action of allelopathy or even accentuation of water stress, which influences the stomata conductance. In studies carried out by [24] with the peanut crop, it was observed that the stomata behavior of the crop presented a variation when there was a change in its environment mainly by the water restriction. When plants absorb CO₂ they lose water due to the opening of the stomata cleft, this natural loss can be accentuated when the plant is exposed to stress caused by competition between species [25,26,27].

In this way, it is well known that the competition between sunflower and different species of grasses significantly affects its initial development, requiring the crop to increase energy expenditure for its growth. It is necessary to take greater care with invasive plants, taking certain measures for their control, so that no major problems with their productivity will occur[28].

Conclusions

The initial development of the sunflower was impaired under a 49.06 cm² competition per weed and the *Panicum maximum* species caused greater damage to the crop.

References

1. Gomes DP, Bringel JMM, Moraes MFH, Gomes JJA, Leite RMVBC. Qualidade fisiológica e sanitária de sementes de girassol produzidas na região de Timon, Maranhão. Summa Phytopathologica. 2006;32(3):291-292. (in Portuguese) <http://dx.doi.org/10.1590/s0100-54052006000300016>
2. PERSON, L. C. A cultura do girassol como estratégia de competitividade para o agronegócio regional e nacional: importância para a agroenergia e a alimentação. In: VI Workshop Agroenergia Matérias Primas, 2012, Ribeirão Preto. Anais do VI Workshop Agroenergia Matérias Primas, Ribeirão Preto: APTA, 2012.
3. Pitelli RH. Interferência de plantas daninhas em culturas agrícolas. Infome Agropecuário. 1985;11(129):16–27. (in Portuguese)
4. Swanton CJ, Nkoa R, Blackshaw RE. Experimental Methods for Crop–Weed Competition Studies. Weed Science. 2015;63:2-11. <https://doi.org/10.1614/WS-D-13-00062.1>
5. Dass A, Shekhawat K, Choudhary AK, Sepat S, Rathore SS, Mahajan G, Chauhan BS. Weed management in rice using crop competition-a review. Crop Protection. 2017;95:45-52. <http://dx.doi.org/10.1016/j.cropro.2016.08.005>
6. Adegas FS, Oliveira MF, Vieira OV, Prete CEC, Gazziero DLPE, Voll E. Phytosociological survey of weeds in sunflower crop. Planta Daninha. 2010;28(4):705-716. (in Portuguese) <http://dx.doi.org/10.1590/s0100-83582010000400002>
7. Avelini ED. Interferência entre plantas daninhas e cultivadas. In: Kogan M, Lira VJE. Avances en manejo de malezas en la producción agrícola y florestal. Santiago do Chile: PUC/ALAM, 1992. p.41–58.

- 271 8. Borchardt L, Jakelaitis A, Valadão FCA, Venturoso LAC, Santos CL.
272 Períodos de interferência de plantas daninhas na cultura do feijoeiro-
273 comum (*Phaseolus vulgaris* L.). Revista Ciência Agronômica.
274 2011;42(3):725-734. (in Portuguese)
- 275 9. Alves GS, Tartaglia FL, Rosa JC, Lima PC, Cardoso GD, Beltrão NEM.
276 Periods of weed interference in sunflower crop in Rondônia. Revista
277 Brasileira de Engenharia Agrícola e Ambiental. 2013;17(3):275-282. (in
278 Portuguese) <http://dx.doi.org/10.1590/s1415-43662013000300005>
- 279 10. Lamego FP, Reinehr M, Cutti L, Aguiar ACM, Rigon CAG, Pagliarini IB.
280 Morphological alterations in wheat, italian ryegrass and radish seedlings
281 when in competition in at the initial stages of growth. Planta Daninha.
282 2015;33(1):13-22. (in Portuguese) [http://dx.doi.org/10.1590/S0100-](http://dx.doi.org/10.1590/S0100-83582015000100002)
283 [83582015000100002](http://dx.doi.org/10.1590/S0100-83582015000100002)
- 284 11. Colbach N, Collard A, Guyot SHM, Mézière D, Munier-Jolain N.
285 Assessing innovative sowing patterns for integrated weed management
286 with a 3D crop: weed competition model. European Journal of Agronomy.
287 2014;53:74-89. <https://doi.org/10.1016/j.eja.2013.09.019>
- 288 12. Munier-Jolain NM, Collard A, Busset H, Guyot SHM. Investigating and
289 modelling the morphological plasticity of weeds. Field Crops Research.
290 2014;155:90-98. <http://dx.doi.org/10.1016/j.fcr.2013.09.018>
- 291 13. Raji B, Cantarella H, Quaggio JÁ, Furlani AMC. Recomendações de
292 adubação e calagem para o Estado de São Paulo. 2.ed. Campinas: IAC,
293 285p. 1996. (in Portuguese)
- 294 14. CARLQUIST, S. Ecological strategies of xylem evolution. Berkeley:
295 University of California, 1975. 259 p.
- 296 15. Lisboa LAM, Dias GHO, Ribeiro KB, Hiago Augusto Amaral Sacco HAA,
297 Padovan JVR, Rodrigues GB, Silva GG, Cardoso AS, Pereira LB, and
298 CRA. Urochloa Cultivated in Soil with Aluminium: A Development and
299 Stomatal Approach. International Journal of Plant & Soil Science.
300 2017;24(4):1-7. <http://dx.doi.org/10.9734/IJPSS/2018/44180>
- 301 16. Banzatto DA, Kronka SN. Experimentação Agrícola. 4.ed. Funep, 2013.
302 237p. (in Portuguese)

17. Silva FAS, Azevedo CAV. The Assistat Software Version 7.7 and its use in the analysis of experimental data. *African Journal Agriculture Research*. 2016;11(39):3733-3740. <http://dx.doi.org/10.5897/AJAR2016.11522>
18. Merotto Junior A, Vidal RA, Fleck NG, Almeida ML. Weed interference in the initial development of soybean and rice plants through light quality. *Planta Daninha*. 2002;20(1):9-16. (in Portuguese) <http://dx.doi.org/10.1590/s0100-83582002000100002>
19. Figueiredo PAM, Ramos SB, Viana RS, Lisboa LAM, Heinrichs R. Alterações morfoanatômicas foliares da cana-de-açúcar na fase de estabelecimento em condições de matocompetição. *Planta Daninha*. 2013;31(4):777-784. (in Portuguese) <http://dx.doi.org/10.1590/s0100-83582013000400003>
20. Lisboa LAM, Viana RS, Ferrari S, Venâncio VG, Figueiredo PAM. Initial development of cotton when submitted to grass competition. *Colloquium Agrariae*. 2017;13(3):88-97. (In portuguese) <http://dx.doi.org/10.5747/ca.2017.v13.n3.a177>
21. Ballaré CL, Casal JJ. Light signals perceived by crop and weed plants. *Field Crops Research*. 2000;67(2):149-160. [http://dx.doi.org/10.1016/s0378-4290\(00\)00090-3](http://dx.doi.org/10.1016/s0378-4290(00)00090-3)
22. Pitelli RA. Competição e controle de plantas daninhas em áreas agrícolas. *Série Técnica IPEF*. 1987;4:1-24. (in Portuguese)
23. Silva RR, Reis MR, Mendes KF, Aquino LA, Pacheco DD, Ronchi CP. Interference periods of weeds in sunflower crop. *Bragantia*. 2013;72(3):255-261. (in Portuguese) <http://dx.doi.org/10.1590/brag.2013.040>
24. Graciano ESA, Nogueira RJMC, Lima DRM, Pacheco CM, Santos RC. Growth and photosynthetic capacity of the peanut cultivar BR 1 under saline conditions. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2011;15(8):794–800. (in Portuguese) <http://dx.doi.org/10.1590/S1415-43662011000800005>
25. Mc Dermitt DK. Sources of error in the estimation of stomatal conductance and transpiration from porometer data. *Hort Science*. 1990;25(12):1538-48.

- 336 26. Paiva AS, Fernandes EJ, Rodrigues TJD, Turco JEP. Stomatal
337 conductance in leaves of bean plants submitted to different irrigation
338 regimes. *Engenharia Agrícola*. 2005;25(1):161-169. (in Portuguese).
339 <http://dx.doi.org/10.1590/s0100-69162005000100018>
- 340 27. Gholz HL, Ewel KC, Teskey RO. Water and forest productivity. *Forest*
341 *Ecological Management*. 1990;30(1):1-18. [https://doi.org/10.1016/0378-](https://doi.org/10.1016/0378-1127(90)90122-R)
342 [1127\(90\)90122-R](https://doi.org/10.1016/0378-1127(90)90122-R)
- 343 28. Brighenti AM, Castro C, Oliveira Junior RS, Scapim CA, Voll E, Gazziero
344 DLP. Interference periods of weeds in sunflower crop. *Planta Daninha*.
345 2004;22(2):251-257. (in Portuguese) [http://dx.doi.org/10.1590/s0100-](http://dx.doi.org/10.1590/s0100-83582004000200012)
346 [83582004000200012](http://dx.doi.org/10.1590/s0100-83582004000200012)