Original Research Article

Comparison of the Physicochemical Properties of Aubergine (Solanaceae) varieties *Solanum aethiopicum gilo* and *Solanum melogena* grown in Northern Côte d'Ivoire

ABSTRACT

In Northern Côte d'Ivoire, aubergines *Solanum aethiopicum gilo* and *Solanum melogena* are widely consumed but no rather exploited. From both aubergines fruits, the comparative physicochemical properties reveal higher size, weight and protein content for *S. melogena*, with respective values of 15.51±1.73 cm, 161.21±33.82 g, and 1.81±0.06%. With *S. aethiopicum gilo*, greater contents are recorded for reducing carbohydrates (0.18±0.02 %), phytates (27.74±0.86 mg/100g), and oxalates (36.85±5.63 mg/100g). Besides, the medial circumference and the contents in moisture, fat, total carbohydrates, crude fibre, vitamin C, polyphenols, ash and minerals elements, and also the total caloric energy value do not show any significant change between both raw food products. The study strengthens the dietary fibre and natural polyphenols and minerals intake from the consumption of these aubergines although phytates and oxalates are recovered within as antinutrients components.

***Keywords**: Aubergine, Solanum aethiopicum gilo, Solanum melogena, Physico-chemical properties, Korhogo*

1. INTRODUCTION

Aubergine (*Solanum melogena*), chili pepper (*Capsicum* spp.) and tomato (*Solanum lycopersicum*) are among the forty (40) most produced vegetable species in the world **[1]**. According to statistics from the United Nations Food and Agriculture Organization, world production of S. melogena is estimated at 49.4 million tons, tomato production is estimated at 164.5 million tons when chili production is of 8.8 million tons.

However, the aubergine variety *Solanum aethiopicum gilo* is not rather known over the world. This aubergine variety is of the fruit vegetables that are most commonly grown and consumed in tropical Africa and therefore spelled ''African aubergine''. According to Lester and Seck **[2]**, it is ranked third in terms of legumes consumption after tomatoes, onions and okra. From sub-Saharan Africa, these authors estimated annual fruit production for 4,500 tons in Burkina Faso, 8,000 tons in Senegal, and 60,000 tons in Côte d'Ivoire.

The plant species *S. aethiopicum gilo* originates from Africa, while the running aubergine (*S. melogena*) is an Asian originated product. Both of these plants are grown in Côte d'Ivoire, where their fruits and leaves are usually eaten by populations **[3]**. Specifically, in Korhogo, northern Côte d'Ivoire, these vegetables are mostly grown by women but also by local young men seeking to find incomes and fit their livelihoods. This crop is grown in slopes of lower or larger dams in northern Côte d'Ivoire **[4]**.

The aubergines raw products are used for the preparation of sauces and consumed with rice and pounded meals such as local *''foufou''* or *''foutou''*. They are also intrant ingredients for the preparation of ''*Tchepodjen''*, a rice meal originating from Senegal. These aubergines are expected to display greater antioxidant activity and hypotensive and diuretic effects.

However, scanty laboratory research information is known from the aubergines varieties *Solanum melogena* and *Solanum aethiopicum gilo*. Unfortunately, they fit no rather utilization for meals and still rotting in numerous markets once harvested and conveyed for the sale. The current study is attempted for providing data regarding the physicochemical properties of both aubergines and so contributing to their valorization.

UNDER PEER REVIEW

2. MATERIAL AND METHODS

2.1 Material

The investigations are performed on aubergines fruits deriving from *Solanum aethiopicum gilo* and *Solanum melogena* species. Both aubergines are produced in Korhogo, northern Côte d'Ivoire.

2.2 Methods

2.2.1 Sampling

The aubergines samples were purchased from three (3) main markets of Korhogo, namely ''Sinistré'' market, ''Soba'' market, and ''Koko'' market. Per market, 10 kg sample of each aubergine were purchased from three various sellers. Thus, 30 kg of each aubergine were gathered per market, leading to 180 kg (30 kg * 2 varieties * 3 markets) for overall aubergine samples purchased. The samples were then conveyed into laboratory for further analyses.

2.2.2 Physical Characterization of *Solanum aethiopicum gilo* and *Solanum melogena*

Five (5) physical parameters were assessed on the aubergine fruits, namely length, circumference, weight, moisture, and ash.

The length and the circumference of the full fruit were estimated using a meter tape, and allowed the deduction of the general fruit shape. The fruit weight was measured using a 2 digits scale (Sartorius).

The method used for determining the moisture was that suggested by AOAC **[5]**. The moisture was assessed by drying 5 g of aubergine into an oven at 105 °C till constant weight resulted after 24 h. The ash content was measured by incinerating five (5) g of oven-dried aubergine into a muffle furnace at 550 °C for 12 h.

2.2.3 Determination of the chemical trend of the aubergine fruits

2.2.3.1 Acidity

The acidity traits (pH and titratable acidity) were measured using AOAC methods **[5]**. Ten (10) grams of crushed sample were slurried in 100 mL of distilled water. The resulting solution was filtered on Whatman micropore filter paper. The pH was thus directly measured out by immersing the previously calibrated pH meter (HANNA) electrode in the filtrate. However, 10 mL of the filtrate were taken and titrated with a NaOH solution (0.1 N) in the presence of phenolphthalein. The NaOH volume used into the sample for resulting to a persistent pink color allowed the deduction of the titratable acidity given in mEq/100g of dried sample.

2.2.3.2 Total soluble carbohydrates and reducing carbohydrates contents

Ethanosoluble carbobydrates were extracted from 1 g of ground dried aubergine with 20 mL of 80% (v/v) ethanol, 2 mL of 10% (m/v) zinc acetate and 2 mL of 10% (m/v) oxalic acid, according to the method of Agbo *et al*. **[6]**. The extract was centrifuged at speed of 3,000 rpm for 10 min. The ethanol residue was evaporated from the extract upon a hot sand bath.
Then, the extracted total soluble carbohydrates were measured out using the method of Dubois *et al*. **[7]**. The operation consisted in adding 0.9 mL of distilled water, 1 mL of 5 % (m/v) phenol, and 5 mL of 96% sulfuric acid into 100 µL of extract, then measuring the absorbance at 490 nm with a spectrophotometer (PG instruments). For the reducing sugars, 1 mL of extract was processed with 0.5 mL of distilled water and 0.5 mL of 3, 5- dinitrosalycilic acid **[8]** prior to the recording of the absorbance from the final solution at 540 nm with a spectrophotometer (PG instruments).

Calibrations were performed with standard solutions of glucose and sucrose for recovering the final total carbohydrates and reducing carbohydrates contents in the studied samples.

2.2.3.3 Lipids content

UNDER PEER REVIEW

Lipids were quantified from 10 g of ground dried aubergine sample by solvent extraction using 300 mL of n-hexane reagent and a Soxhlet device for 7 h **[9]**. The hexan-oil mixture resulted from the extraction was recovered and separated with a rotavapor apparatus (Heidolph). The difference between the sample weight before and after the experiment allowed the estimation of the lipids content.

2.2.3.4 Proteins content

Crude proteins content was determined as the total nitrogen using the Kjeldhal method **[5]**. Thus, 1 g of aubergine mash was mineralized at 400 °C for 2 h, with adding of concentrated sulfuric acid (H_2SO_4) and potassium sulfate (K_2SO_4) catalyst. The mineralizate was diluted and distilled for 10 min. Thereafter, the distillate collected into a flask containing boric acid and methylen bromocresol reagents ion, was titrated for the total nitrogen using ammonium sulfate ($(NH_4)_2SO_4$). The crude protein content of the aubergine was deduced from the nitrogen level using 6.25 as conversion coefficient.

2.2.3.5 Fibers content

The determination of the crude fibers content consisted in treatment of 2 g of ground aubergine sample with 50 mL of 0.25N sulfuric acid and 50 mL of 0.31 N sodium hydroxide and filtration of the resulting solution upon Whatman paper. The residue was dried for 8 h at 105 °C then incinerated at 550 °C for 3 h into ovens **[10]**. The final residue was weighed as crude fibers and expressed in percentage.

2.2.3.6 Total carbohydrates content and energy value

Total carbohydrates and energy values were determined using calculation formulas **[11]** accounting the moisture, fat, protein, ash contents and the energetic coefficients for macromolecules.

$$\mathbf{TCC\ (\%) = 100\text{-}[P(\%) + W(\%) + F(\%)+A(\%)]}$$

$$\mathbf{CEV\ (kcal/100g) = [(4\times P) + (9\times F) + (4\times C)]}$$

With: TCC, total carbohydrates content; CEV, caloric energy value; P, protein content; M, moisture content; F, fat content; A, ash content; C, total carbohydrates content

2.2.3.7 Polyphenols contents

The phenol compounds were extracted from aubergine with methanol reagent. One gram of dried aubergine sample was homogenized in 10 mL of methanol solution 70% (v/v). The resulting mixture was centrifuged at 1,000 rpm for 10 min. The pellet was recovered and treated likewise. The deriving supernatants were thus gathered into a marked flask and added with distilled water at 50 ml.

The total polyphenols content was measured using Folin-ciocalteu reagent, sodium carbonate solution (20% w/v) and distilled water **[12]**. Essays were measured for their absorbance at 745 nm with a spectrophotometer against standard gallic acid solutions taken as polyphenols control.
The tannins content was deducted from the total polyphenols using vanillin reagent **[13]**. Essays were measured for their absorbance at 500 nm with a spectrophotometer against standard tannic acid solutions taken as tannins control.
Flavonoids content was also determined from the total polyphenols using aluminum chloride (10% w/v), potassium acetate (1 M) and distilled water **[14]**. Essays were measured for their absorbance at 415 nm with a spectrophotometer against standard quercetin solutions taken as flavonoids control.

2.2.3.8 Vitamin C content

The vitamin C was evaluated from the aubergines using 2,6- dichlorophenol-indophenol (DCPIP) reagent **[15]**. Ten (10) grams of ground dried aubergine sample were dissolved into 40 mL of metaphosphoric acid-acetic acid solution (2%, w/v). The resulted mixture was centrifuged at 3,000 rpm for 20 min. Thus, the supernatant was recovered, added with boiled distilled water for 50 mL, and titrated with 2, 6- DCPIP solution (0.5 g/L) previously calibrated with a pure vitamin C solution.

UNDER PEER REVIEW

2.2.3.9 Oxalates content

The oxalate content was determined with the standard AOAC method **[5]**. Two (2) grams of ground dried aubergine sample were homogenized into 200 mL of distilled water and added with 20 mL of 6N hydrochloric acid (HCl). The mixture was heated in boiling water bath for 1 h, cooled, and filtered. Fifty (50) mL of the filtrate were then homogenized into 20 mL of 6 N HCl, and filtered again. The 2nd filtrate was treated with methyl red (0.1%, w/v), concentrated ammonia, heated, and filtered. The 3rd filtrate was boiled, treated with calcium chloride (5%, w/v) for the formation of calcium oxalate crystals, and then filtered once more. The residues deriving from the filtration steps were successively washed with distilled boiling water, dried into an oven; dissolved into 10 mL of diluted sulfuric acid, and titrated with 0.05N potassium permanganate solution.

2.2.3.10 Phytates content

The phytates were measured according to the method processed by Mohammed *et al*. **[16]**. A slight ground aubergine sample (0.5 g) was treated with 25 mL of TCA solution at 3% (w/v) and centrifuged at 3,500 rpm for 15 min. Five (5) mL of the supernatant was removed, treated with 3 mL of ferric chloride 1% (w/v) reagent, heated in a boiling water bath, cooled and also centrifuged at 3,500 rpm for 10 min. The 2nd supernatant was treated with 5 mL of 0.5N hydrochloric acid, 5 mL of 1.5N sodium hydroxide, heated in a boiling water bath and centrifuged once more at 3500 rpm for 10 min. Thus, 1 mL of the final supernatant was added with 4.5 mL of distilled water and 4.5 mL of orthophenantroline reagent and then measured for the absorbance at 470 nm with a spectrophotometer against standard Mohr salt solution treated likewise and taken as phytates ferric control.

2.2.3.11 Determination of mineral elements

The determination of the mineral elements was performed according to the IITA method **[17]**. Finely ground aubergine sample (0.4 g) previously oven dried at 60 ° C was incinerated into a muffle furnace at 550 °C for 3 h. The resulting gray-white ash was cooled, added with 2 mL of half-diluted HCl, placed on a sand bath at 120 °C until full evaporation, and then ovened at 105 °C for a 1 h. The final dry extract was recovered with 2 mL of half-diluted HCl, filtered, and the resulting filtrate added with distilled water, and lanthanum chloride. The mineral elements in the solution were then measured using Atomic Absorption Spectrometry (AAS 20 type VARIAN).

2.2.4 Statistical analyzes

All essays were performed in triplicate. The data collected were statistically analyzed using Statistica software (Statistica 7.1) at 95% significance. A statistical Student T- test was performed for the comparison between both aubergines varieties studied regarding each parameter assessed. For each characteristic, the results were expressed as averages followed by the standard deviations. Also, the general averages have been calculated for the characteristics displaying any significance divergence between the studied products.

3. RESULTS AND DISCUSSION

3.1 Physical Characteristics of aubergines

The aubergine *S. aethiopicum gilo* displays a round shape with flattened ends whereas the aubergine *S. melogena* has an elongated shape. Both aubergines show invarious medium circumference (18.04±0.52 cm) and moisture and ash contents (92.62±0.55% and 0.59±0.25%, respectively). Oppositely, **table 1** shows higher size and weight from *S. melogena* (15.51 cm and 161.21 g) compared to the respective 3.79 cm length and 61.07 g weight of *S. aethiopicum gilo*.

3.2 Major chemical composition of aubergines

UNDER PEER REVIEW

Except for the proteins and reducing carbohydrates contents and the acid values, the main chemical traits do not show any obvious divergence between *S. aethiopicum gilo* and S. melogena (**Table 2**). Indeed, *S. melogena* provides more protein content (1.81%) but lower reducing carbohydrates content (0.11%) compared to *S. aethiopicum gilo* (1.68% and 0.18%, respectively). Both aubergines varieties are fairly acid. However, the comparison shows *S. aethiopicum gilo* as a lower acid traits aubergine (pH of 7.67 and titratable acid value of 0.46 mEq/100g) compared to *S. melogena* (pH of 6.93 and acid value of 0.14 mEq/100g).

Table 1. Some physical parameters of *Solanum aethiopicum gilo* and *Solanum melogena*

Parameters	*S. aethiopicum gilo*	*S. melogena*	T_{-value}	P_{-value}	General average
Shape	Round with flattened ends	Elongated	-	-	
Length (cm)	3.79±0.40[b]	15.51±1.73[a]	130.96	**<0.001**	
Medium circumference (cm)	18.32±0.40[a]	17.77±0.66[a]	1.53	0.284	18.04±0.52
Weight (g)	61.07±3.27[b]	161.21±33.82[a]	26.05	**0.007**	
Moisture (%)	92.45±0.34[a]	92.8±0.84[a]	0.455	0.537	92.62±0.55
Ash (%)	0.41±0.22[a]	0.76±0.23[a]	3.790	0.123	0.59±0.25

Per raw, values with various lower scripts are different at 0.05 statistical P-value

Table 2. Chemical composition of *Solanum aethiopicum gilo* and *Solanum melogena*

Parameters	*S. aethiopicum gilo*	*S. melogena*	T_{-value}	P_{-value}	General average
pH	7.67±0.01[a]	6.93±0.02[b]	16428.00	**<0.001**	
TTA	0.14±0.02[b]	0.47±0.03[a]	$2.921.10^{33}$	**<0.001**	
TPC (%)	1.68±0.04[b]	1.81±0.06[a]	9.274	**0.038**	
FMC (%)	0.12±0.02[a]	0.12±0.01[a]	0.500	0.519	0.12±0.01
TFC (%)	2.75±0.51[a]	1.96±0.49[a]	3.758	0.125	2.35±0.57
TCC (%)	2.10±0.37[a]	2.12±0.51[a]	0.003	0.959	2.11±0.36
RCC (%)	0.18±0.02[a]	0.11±0.03[b]	20.346	**0.011**	
TGC (%)	5.33±0.18[a]	4.50±0.54[a]	6.313	0.066	4.91±0.53
TEV (Kcal/100g)	29.12±0.67[a]	26.26±2.45[a]	3.793	0.123	27.69±2.05

Per raw, values with various lower scripts are different at 0.05 statistical P_{-value}.pH, potential of hydrogen; TTA, total titratable acidity; TPC, total proteins content; FMC, fat matter content; TFC, total fibers content; TCC, total carbohydrates content; RCC, reducing carbohydrates content; TGC, total glucides content; TEV, total caloric energy value;

Regarding the contents in fat matter, total fibre, total carbohydrates, total glucides, as far as the energy value, both studied aubergines provide similar means. For those characteristics, the respective general contents averages are 0.12%, 2.35%, 2.11%, and 4.91%, providing 27.69 Kcal/100 g as energy value (**table 2**).

3.3 Main micronutrients, polyphenols compounds and antinutrients deriving from aubergines

The micronutrients, secondary metabolites (polyphenols) and antinutrients in the aubergines studied are gathered in **table 3**. Both aubergines provide statistically invarious mean values regarding the contents in polyphenols (total polyphenols, tannins, and flavonoids), vitamin C, and mineral elements (phosphorus, potassium, calcium, magnesium, sodium, iron, manganese, zinc, and copper). For polyphenols, the general contents averages from 100 g aubergine dried sample are 51.55 mg total polyphenols, 29.78 mg tannins, and 0.95 mg flavonoids; while the vitamin C content is measured at

UNDER PEER REVIEW

9.46 mg/100 g. From the main mineral elements, macro elements are measured between 8.51 ppm (Sodium) and 0.99 % DM (phosphorous), when oligoelements oscillate from 0.018 ppm (copper) to 4.89 ppm (iron) as shown in **table 3**.

Besides, from the antinutrients compounds, *S. aethiopicum gilo* provides more oxalates content (36.85 mg/100g) and phytates content (27.74 mg/100g) compared to *S. melogena* (25.63 mg/100g and 23.74 mg/100g, respectively).

Table 3. Micronutrients, polyphenols compounds and antinutrients contents in S. aethiopicumgilo and S. melogena

Charateristics		SAG	SM	T_{-value}	P_{-value}	General average
Polyphenols compounds (mg/100g)	**Total polyphenols**	54.74±5.51[a]	48.36±6.89[a]	1.047	0.364	51.55±7.01
	Tannins	35.26±8.81[a]	24.3±2.65[a]	2.837	0.167	29.78±8.51
	Flavonoids	1.18±0.3[a]	0.73±0.39[a]	1.526	0.284	0.95±0.42
Vitamin and minerals	**Vitamin (mg/100g)**	9.08±0.36[a]	9.85±2.04[a]	0.405	0.559	9.46±1.47
	Phosphorus (% DM)	0.92±0.13[a]	1.07±0.3[a]	0.349	0.587	0.99±0.26
	Potassium (% DM)	5.33±0.19[a]	7.74±1.28[a]	6.872	0.059	6.54±1.51
	Calcium (% DM)	0.40±0.039[a]	0.81±0.2[a]	7.167	0.055	0.60±0.25
	Magnesium (% DM)	0.57±0.06[a]	0.74±0.16[a]	1.884	0.242	0.65±0.14
	Sodium (PPM)	7.21±0.4[a]	9.82±1.75[a]	4.234	0.109	8.51±1.82
	Iron (PPM)	4.44±0.3[a]	5.35±0.68[a]	2.912	0.163	4.89±0.7
	Manganese (PPM)	2.30±0.16[a]	2.23±0.19[a]	0.144	0.723	2.26±0.18
	Zinc (PPM)	0.25±0.13[a]	0.31±0.13[a]	0.206	0.673	0.28±0.13
	Copper (PPM)	0.02±0.008[a]	0.02±0.005[a]	0.250	0.643	0.018±0.007
Antinutrients (mg/100g)	**Oxalates**	36.85±5.63[a]	25.63±2.72[b]	9.637	**0.036**	
	Phytates	27.74±0.86[a]	23.74±1.90[b]	11.046	**0.029**	

Per raw, values with various lower scripts are different at 0.05 statistical P_{-value}.
SAG, Solanum aethiopicum gilo; SM, Solanum melogena

Discussion

The study shows higher size of *Solanum melogena* fruit compared to *Solanum aethiopicum* gilo, a rounded aubergine species, whereas their medium circumferences do not vary. Thus, *S. melogena* obviously displays higher weight than *S. aethiopicum gilo*. Both aubergines are unvariously with great moisture value, over 92%. Such moisture contents are close to that revealed from onion variety V1310 (90.44%) by Konate *et al*. **[18]**. The high moisture content of aubergines is disadvantageous for their preservation since the crops could be submitted to rapid post-harvest change and rotting. This phenomenon has been previously observed from ripe palm fruit by Ali *et al*. **[19]**.

S. melogena is more provided with proteins (1.81%) compared to *S. aethiopicum gilo*. But both aubergines contain similar and lower lipids rate (0.12 and 0.12%), that is closed to the 0.14% lipid content in cooked spinach **[20]**. Their total glucides content are also lower (5.33% and 4.50%). Thus, they are not considered as glucides products since the glucides values are more strengthened from the common starchy products such as sweet potato (28.5%) according to the FAO **[21]**. In evidence, the aubergines are legumes crops and are not really used as source of carbohydrates, nor lipids and proteins. So, they are advisable in low-calory dietaries.

The studied aubergines also display similar fibre content around 2.35%, a lower value compared to the 3.92% fibre recorded with the fresh dough processed from the new shoots tuber of *Borassus aethiopum* as reported by Niamke *et al*. **[22]**. However, these aubergines could be significant sources of dietary fibers that are essential for the digestive balance in the intestinal duct and the stomach. Indeed, dietary fibers are factors of healthy body. Many studies have shown opposite correlation between consumption of dietary fibers and the upcoming of colon cancer. Dietary fibers can complex with carcinogenic molecules, thus preventing their contact with the colon and facilitating their

UNDER PEER REVIEW

excretion **[23, 24]**. They account laxative role and help against colorectal cancer. They also usually drop the blood glucose, HDL-cholesterol, LDL-cholesterol, and so contribute in reduction of coronary heart disease **[25]**. Consumption of aubergines may therefore increase gastric volume and constitute a post-ingestive state for feeling more quickly the satiety **[24, 26]**.

Vitamin C is measured in both aubergines below 10 mg/100 g. The consumption of these aubergines added with other vegetables richer in vitamins would be more beneficial. Vitamin C contributes in healthy bones, cartilage, teeth, and gums. It also protects against infections, accelerates healing and promotes the absorption of iron.

Significant total polyphenols contents (around 50 mg/100 g) are recorded from the studied aubergines, with any statistical change per variety. Most important parts of these polyphenols run for tannins (around 30 mg/100 g), while flavonoids are scarcely measured (1 mg/100 g). Polyphenols are credited with numerous health benefits, such as reduction of cardiovascular concerns, inflammatory or neuro-degenerative diseases, cancer prevention, antiplatelet effects, blood pressure regulation, *etc*. **[27]**. But, according to the predominance of tannins, aubergines could not be easily consumed fresh and their heating during cooking submits polyphenols biomolecules to degradation. Nevertheless, the heating process is affordable for succeeding in degradation of antinutrients components (oxalates and phytates) also highly found in the aubergines studied. Otherwise, these antinutrients could counteract the absorption of iron, zinc and calcium **[28]**.

From their mineral composition, the aubergines studied are provided with the same contents in macro elements (Ca, P, Mg, Na, K) and oligo-elements (Fe, Zn, Cu, Mn). It's widely known that the major nutritional concerns from populations in developing countries are about deficiencies regarding proteo-energy nutrients and micronutrients especially minerals. The mineral elements are involved in a wide range of functions in the body, namely mineralization, control of hydric balance, enzymatic and hormonal systems, the muscular system, the nervous and immune systems **[20, 18]**. Thus, the essential nutrients are recommended in food intake according to Martin **[29]** and Konate *et al*. **[18]**. As overall legumes, aubergines are great source of mineral elements as strengthened by the current works. These raw crops are available in sufficient quantities in rural areas, and their use in diversified diet could be highly recommended for local populations.

4. CONCLUSION

The aubergines *Solanum aethiopicum gilo* and *Solanum melogena* contain macronutrients (carbohydrates, proteins, lipids) and several micronutrients (mineral elements, polyphenols, and vitamin) that highlight their significant nutritional potential. In addition, they records significant dietary fibers that regulate the appetite and the digestion. Both aubergines have closer nutritional potential. But, they should be processed for the degradation of the antinutrients compounds prior to their safe consumption.

REFERENCES

1. Lebeau A, 2010. Resistance of tomato, eggplant and chilli to Ralstonia solanacearum: interaction between resistance genitors and bacterial diversity, characterization and mapping of genetic factors involved in eggplant. Thesis. Faculty of Science and Technology. University of Reunion.

2. Lester RN & Seck A, 2004. Solanum aethiopicum L. In: Grubben G. J. H. and O. A. Denton (Eds.). Plant Resources of Tropical Africa 2. Vegetables. PROTA Foundation, Backhuys Publishers, Wageningen, The Netherlands: pp 530-536.

3. Djedji AH & Fondio L, 2013. Cultivate Eggplant in Ivory Coast. National Center for Agronomic Research, 4p. https: // www. Agrici.net/pdf/ftec_aubergine.pdf

4. Tanguy LG, 2004. Agricultural and Pastoral Development in Northern Côte d'Ivoire: Coexistence Problem. Cahier d'Outre-Mer. pp. 259-288.

5. AOAC, 1990. Official methods of analysis of the AOAC, 15th Ed, Methods 932.06, 925.09, 985.29, 923.03. Association of official analytical chemists. Arlington, VA, USA.

6. Agbo NG, Uebersax M & Hosfield G, 1986. An efficient extraction technique of sugars from dry edible beans (*Phaseolus vulgaris*) and estimation in HPLC, Université Nationale de Côte d'Ivoire. Annals series C (sciences) Tome XXI: 169-184.

UNDER PEER REVIEW

7. Dubois M, Gilles KA, Hamilton JK, Rebers PA & Smith F, 1956. Colorimetric method for determination of sugar and related substances. *Anal. Chem*. 28: 350-356.

8. Bernfeld P, 1955. Amylase bêta and alpha (Assay Method), in methods in enzymology, Ed., Academic press, New-york, USA, pp 149-154.

9. AFNOR, 1986. Collection of French standards. Quality control of dairy products. AFNOR, Paris - defense.

10. Wolff JP, 1968. Fat Analysis Manual; Azoulayed., Paris (France), 519p.

11. FAO, 2002. Report of the International Rice commission. Vingtième session, 23-26 juillet 2002, Bangkok, Thaïlande.

12. Singleton VL, Orthofer R &Lamuala RRM, 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of FOLIN-Ciocalteu reagent. *Methods Enzymol*. 299:152-178.

13. Bainbridge Z, Tomlins K, Willings K & Westby A, 1996. Methods for assessing quality characteristics of non-grain starch-staple. Part 4 Advanced methods. National ressources institute, University of Greenwish, UK ISBN 0-85954- 400, 43-79.

14. Meda A, Lamien CE, Romito M, Millogo J &Nacoulma OG, 2005. Determination of the total phenolic, flavonoid and proline contents in Burkina fasan honey, as well as their radical scavenging activity. *Food Chem*. 91: 571-577.

15. AOAC, 1984. Official methods of analysis of the association of official Analytical Chemists (14thedn) Washington. DC.

16. Mohammed AI, Ponhamperuma AJP & Youssef SH, 1986. New chromophore for phytic Acid Determination. *Cereal Chem*. 63(6): 475-478.

17. IITA, 1981. Analyse des prélèvements pédologiques et végétaux. Manuel N°1, Oyo-Road, Nigeria, 66p.

18. Konate M, Parkouda C, Tarpaga V, Guira F, Rouamba A & Hagretou SL, 2017. Evaluation of the nutritional potential and the ability to preserve eleven onion (Allium cepa L.) bulb varieties introduced in Burkina Faso. Int. J. Biol. Chem. Sci. 11 (5): 2005-2015.

19. Ali A., Alhadji.,Tchiegang C. &Saïdou C., 2010. Physico- chemical properties of palmyra palm (*Borassus aethiopum* Mart.) fruits from Northern Cameroon. Afri. J. Foods Sci. 4: 115-119.

20. Ciqual, 2013. Nutritional composition of foods table. https // ciqual. Anses.fr. viewed on 26/08/2018 at 1 pm.

21. FAO, 1982. Etude FAO, Alimentation nutrition 47/2. Utilisation des plantes tropicales : tubercules et racines. 135p.

22. Niamke AM, Saki SJ, Koffi KM, Sea TB, Bidie ADP, Djaman AJ &Biego GH, 2013. Studies on physicochemical properties of fresh paste of the young growths of *Borassus aethiopum*. *Int. J. Plant Anim. and Env. Sci*, 3(4): 197-202.

23. Jansen MC, Buenode-MHB, Buzina R, Fidenza F, Menotti A, Blackburn H, Nissinen AM, Kok FJ &Kromhout D, 1999. Dietary fiber and plant foods in relation to colectoral cancer mortality: the seven countries study. *Int. J. cancer*. 81: 174-179.

24. Chene C, 2003. Journal de l'ADRIANOR Agro-jonction n°33. Septembre-octobre 2003. Les fibres alimentaires pp.1-8

25. Jalili T, Wildamn RFC, Medeiros DM, 2000. Nutraceutical roles of dietary fiber. *J. Nutri. Funct. Med. Foods*. 2: 19-34.

26. Al Dobaib N, 2009. Effect of diets on growth, digestibility, carcass and meat quality characteristics of four rabbits breeds. *Sandi J. Biol. Sci*. 17: 83-93.

27. Purchase S, 2013. Polyphenols in the diet: Extraction, antioxidant power and interaction with metal ions. Thesis. A. Mira University. Bepala. Faculty of Sciences of Nature and Life. Department of Food Sciences (Algeria). University of Avignon and Pays de Vancluse. Doctoral School 536-Avignon (France). 211P.

28. Amiot MJ, 2015. Fresh fruits and vegetables. In Equationnutrition. Aprifel.p1

UNDER PEER REVIEW

29. Martin A, 2001. Recommended dietary intakes for the French population (3rd ed). Tec and Doc Lavoisier, CNERNA-CNRS: Paris; 658p.

UNDER PEER REVIEW