

Evaluation of the insect pest population dynamics in common bean cultivars in relation to the foliar fertilization with potassium silicate.

ABSTRACT

This study is aimed at a grouped evaluation of the occurrence of insect pests from the entomological orders diptera, hemiptera, lepidoptera and coleoptera in two different genotypes of common bean, cultivated in two sowing seasons, according to the foliar fertilization with potassium silicate. The experiment was conducted under field conditions during the Rainy Season Crop and the Dry Season Crop (from August to December 2014 and from February to June 2015, respectively), in a typical eutroferic Red Latosol, in the rural area of Assis Chateaubriand, State of Paraná, Brazil. The experiment used a completely randomized 2 x 5 factorial design with four replications. The first factor refers to the bean genotypes (IPR Campos Gerais and IPR Tuiuiú), and the second factor refers to the doses of potassium silicate (0.0, 250, 500, 750 and 1000 ml ha⁻¹). Potassium silicate applications were divided into biweekly sprays, from phenological stage V₃ to R₈, using the same quantity established for each treatment at different times of application. Insect pest samplings were carried out weekly during the phenological stages V₃ through R₈ for both experiments. Samplings were performed at random points of each plot to provide a better representation. Data collected from each common bean pest found in the crops evaluated were grouped according to their entomological orders and submitted to statistical analysis through use of Sisvar 5.1 software at the 0.05 significance level (P≤0.05) by the F test. The significant results from the analysis of variance were submitted to regression analysis at the 0.05 significance level (P≤0.05) as well. Foliar fertilization with potassium silicate caused a population decrease of the entomological orders evaluated for the common bean genotypes studied. IPR Tuiuiú genotype presented a smaller quantity of hemiptera insects compared to IPR Campos Gerais during the reproductive stage of the Rainy Season Crop (2014). During the vegetative stage of the Dry Season Crop (2015), IPR Campos Gerais presented a smaller quantity of diptera insects compared to IPR Tuiuiú. Interactions between treatments and the genotypes evaluated occurred, decreasing the population of hemiptera and lepidoptera insects during the vegetative and reproductive stages of the Dry Season Crops (2015). IPR Campos Gerais Genotype presented a smaller quantity of lepidoptera and hemiptera insects than IPR Tuiuiú.

Keywords: *Crysodeicis includens*; *Diabrotica speciosa*; *Liriomyza* sp.; Hemiptera.

1. INTRODUCTION

The common bean (*Phaseolus vulgaris* L.) is very susceptible to insect pest attacks, which affect its productivity. All the common bean plant structures have been shown to be susceptible due to the diversity of pest species. Many of these species have been reported as bean pests [1], but few of them are known to be of economic importance.

However, several arthropods and molluscs species can cause significant reductions in crop productivity. Such damages may vary according to the pest species, crop growth stage, bean cultivar and sowing season, with potential losses up to 100% [2]. Among the insect pests found in bean crops in Brazil, Silverleaf Whitefly (*Bemisia tabaci*), Bean Leafhopper (*Empoasca* sp.), Cucurbit Beetle (*Diabrotica speciosa*), Southern Armyworm (*Spodoptera eridania*) and Tobacco Budworm (*Heliothis virescens*), Broad Mite (*Polyphagotarsonemus latus*), Southern Green Stink Bug (*Nezara Viridula*) and Neotropical Brown Stink Bug (*Euschistus heros*) are those responsible for the largest productivity losses. In the producing regions, Leaf Miner (*Liriomyza* sp.), slugs (*Vaginula lamsdorffii*), *Chrysomelidae* and *Caliothrips phaseoli* larvae have been highlighted as important pests of beans [3].

The importance of common bean in Brazil makes necessary development of new crop management methods to provide farmers with increased productivity and lower costs. Thus, an alternative is the use of foliar fertilization with some micronutrients, such as silicon (Si).

The use of foliar fertilization with macro- and micronutrients aiming at increase of productivity or induction of resistance has become a common practice by farmers in Brazil's bean crops [4].

Silicon [5] does not meet the criteria of essentiality for plants. However, [6, 7] this element is characterized as beneficial for plants since it makes them more resistant to pest attacks and diseases, improves photosynthetic capacity, increases the number of leaves, stem diameter, plant size and, consequently, the productivity. Thus, fertilization with potassium silicate may result in a lesser occurrence of insect pests in the bean crop.

Studies related to management that employs silicon in the agriculture are still incipient and inconclusive [8], especially those seeking to make clear the relation between plant nutrition and micronutrient and injuries caused by insect pests in agronomic aspects of the vegetables.

This study is aimed at a grouped evaluation of the occurrence of insect pests from the entomological orders diptera, hemiptera, lepidoptera and coleoptera in two different genotypes of common bean, cultivated in two sowing seasons, according to the foliar fertilization with potassium silicate.

2. MATERIAL AND METHODS

The experiment was conducted under field conditions during the Rainy Season Crop and the Dry Season Crop (from August to December 2014 and from February to June 2015, respectively), in a typical Eutroferic RED LATOSOL of the rural area of Assis Chateaubriand, located in the Western State of Paraná, Brazil. The area is located at coordinates 24°17'27.40'' S latitude and 53°35'03.99'' W longitude, at an altitude of 321 m.

According to Köppen (1984), the region climate is type Cfa (subtropical), with an average temperature under 18°C (mesothermic) in the coldest month and an average temperature over 22°C in the hottest month. This climate is characterized by hot summers, infrequent frosts, and by the propensity to concentrate rainfall in summer months, but without a defined dry season. Meteorological data observed during the experiments conduction are in (figures 1 and 2).





Figure 1: Maximum temperature, rainfall, and relative air humidity during the experiment conduction. Bean crop phenological stages evaluated: VE (germination/emergence); V2 (primary leaves totally opened); V3 (first trifoliate leaf); V4 (third trifoliate leaf); R5 (floral buds); R7 (first pods appear); R8

65 (first full pods); R9 (change of color of pods – physiological maturation). Rainy Season Crop (2014). Assis
 66 Chateaubriand, Paraná, Brazil.
 67
 68



69
 70
 71 **Figure 2: Maximum temperature, rainfall, and relative air humidity during the experiment**
 72 **conduction. Bean crop phenological stages evaluated: VE (germination/emergence); V2 (primary**
 73 **leaves totally opened); V3 (first trifoliolate leaf); V4 (third trifoliolate leaf); R5 (floral buds); R7 (first pods appear); R8**
 74 **(first full pods); R9 (change of color of pods – physiological maturation). Dry Season Crop (2015). Assis**
 75 **Chateaubriand, Paraná, Brazil.**
 76

77 The experiment used a completely randomized 2 x 5 factorial design with four replications.
 78 The first factor refers to the common bean (*Phaseolus vulgaris* L.) cultivars (IPR Campos Gerais and
 79 IPR Tuiuiú), and the second factor refers to the doses of potassium silicate (0.0, 250, 500, 750 and
 80 1000 ml ha⁻¹) of a commercial product (Supa Sílica – Agrichem®) containing 0.9 p.v. SiO₂ (90 grams
 81 per liter of water) and 18% K₂O in its formulation. The product was diluted in humic acid. Plots were
 82 10 meters long by 7.74 m wide, totaling 18 planting rows and 77.4 m² area. Spacing used was 0.43 m
 83 between planting rows, at a density of 12 seeds per linear meter.

84 A commercial adjuvant (LI 700 – Sangose®) was added to the spray mixture preparation with
 85 potassium silicate to improve retention of the treatments on bean leaves. This adjuvant is composed
 86 of a blend of Phosphatidylcholine (Soya lecithin) and Propionic Acid, which boosts foliar absorption of
 87 nutrients.

88 Seeds were treated with fungicide (Carbendazim – dose of 300 ml/100 kg of seeds) and
 89 insecticide (Imidacloprid + tiodicarb, dose of 0.6 L/100 kg of seeds), as recommended by the
 90 manufacturer, so that the seedlings would be protected until the effective period of the products is
 91 finished (approximately 15 days) and the applications of the treatments evaluated begin.

92 Sowings were made on August 23, 2014 (Rainy Season Crop), and on February 28, 2015
 93 (Dry Season Crop). Harvests were made on December 13, 2014, and on June 11, 2015, respectively.

94 When necessary during the crop growth, weed control was carried out through manual
 95 weeding (V2 phenological stage of the Rainy Season Crop – 2014) or application of the herbicide
 96 Fomesafen + fluazifop-P-butyl at a dose of 1,6 L ha⁻¹, as recommended by the manufacturer (V7
 97 phenological stage of the Rainy Season Crop and V4 phenological stage of the Dry Season Crop –
 98 2015).

For this study, a composite soil sample (500 grams) was collected, based on 20 sub-samples, to a depth of 0 – 20 cm. The sample was sent to laboratory for chemical (COODETEC) and physical (UNITHAL) determination of macro/micronutrients, and for soil soluble Silicon determination (Federal University of Uberlândia - UFU), presenting the following results (table 1):

Table 1. Chemical analysis of soil performed (0 -20 cm) in depth, indicating the characteristics of soil fertility of the experimental area.

P	pH	pH	H + Al	Al ³⁺	Mg ²⁺	Ca ²⁺	K ⁺
mg dm ⁻³	CaCl ₂)	(H ₂ O)	cmol _c dm ⁻³				
6.30	4.80	5.6	3.18	0.00	1.25	4.05	0.16
Mn	Fe	Cu	Zn	Si			
mg dm ⁻³							
146.49	55.72	7.62	3.28	19.6			
V	Clay	Coarse sand	Fine sand	Gravel	Silt		
%							
63.19	73.5	2.3	3.2	0.0	21.0		
Textural class							
Clayey soil							

Base fertilization was performed according to the soil analysis, using 16-16-16 (N-P-K) fertilizer at a dose of 300 Kg ha⁻¹ in the sowing time (August 23, 2014, for Rainy Season Crop and February 28, 2015, for Dry Season Crop). Subsequently, at the V₄ phenological stage, nitrogen top dressing was applied with urea (45% N) at a dose of 64 kg ha⁻¹, as observed in the (figures 1 and 2).

Potassium silicate applications were divided into biweekly applications from phenological stage V₃ to R₈ (figures 1 and 2). The same quantity of potassium silicate established for each treatment was used at different application times. For this purpose, an electric knapsack sprayer was used with fixed working pressure at 45 psi, aided by a spray bar with 4 flat fan nozzles spaced 50 cm apart, with spray volume of 186 L ha⁻¹. A digital pH meter was calibrated with pH 4 and 7 buffers, and then used to measure the spray mixture pH of each treatment. The pH values of the products used (potassium silicate and spray adjuvant) were provided by the manufacturers.

Insect pest samplings, in turn, were carried out weekly between the phenological growth stages V₃ and R₈ for both experiments, as seen in the (figures 1 and 2). Each Insect pest sample was divided into three sub-samples per plot. Then, insects found in each of these sub-samples were counted to determine the total in one sample from each plot. The same criteria of insect pest sampling was performed in both crops conducted (Rainy Season Crop, 2014, and Dry Season Crop, 2015). After the sampling time in each crop finished, insects were quantified and grouped according to the entomological orders to make possible the determination of pest occurrence. The average number of insects counted in each treatment with potassium silicate was set according to their entomological orders, comparing these results with the phenological stages of both crops (rainy season crop, 2014, and dry season crop, 2015). Zig-zag samplings were performed in the plots to provide a better representation [9].

For insects belonging to the order **diptera**, only the leaf sapper (*Lyriomyza sp.*) was sampled between the growth stages V₃ and V₄. Only larvae present in the galleries formed in the foliar limb of the plants were counted.

As for the insects from the other orders evaluated (Lepidoptera, Hemiptera and **coleoptera**), samplings were carried out between the phenological stages V₃ - R₈. The sampled insects corresponding to the order Hemiptera were the Southern Green Stink Bug – *Nezara viridula* (Linnaeus, 1758); the Neotropical Brown Stink Bug – *Euschistus heros* (Fabricius, 1798), and the Bean Leafhopper – *Empoasca sp.* (Göethe, 1875). As for **coleoptera**, the insects sampled were the Cucurbit Beetle – *Diabrotica speciosa* (Germar, 1824); the *Cerotoma arcuate* (Olivier, 1791); and the *Lagriites Solier* (Sphindidae, 1839). Regarding the order Lepidoptera, caterpillars sampled were the Soybean Looper – *Crysodeixis includes* (Walker, 1858); the Velvetbean Caterpillar – *Anticarsia gemmatilis* (Hübner, 1818); the Southern Armyworm – *Spodoptera eridania* (Cramer, 1782); and the Tobacco Budworm – *Heliothis virescens* (Fabricius, 1781).

To aid the sampling of insect pests belonging to the orders **coleoptera** (Cucurbit Beetle, *Cerotoma arcuate* and *Lagriites Solier*), Hemiptera (Bedbugs) and Lepidoptera (Caterpillars), a white beat cloth measuring 1 m x 1 m was used. At each sampling, the cloth was inserted between the planting rows, covering the plants of the next row. Afterwards, the plants were shaken on the beat cloth so that the insects contained on them were counted. Samplings were performed at random points of each plot to provide a better representation.

Sampling times were divided according to the crop growth stages for the two seasons. In the experiment conducted during the **Rainy Season Crop** (2014), three sampling times were carried out during the vegetative stages and nine during the reproductive stages. For **Dry Season Crop** (2015), four sampling times were carried out during the vegetative stages and five during the reproductive stages of bean growth. The quantity of sampling times differed between cultivations due to the growth period of each crop. Sampling times were related to the crops cycle and the time they were in the field. Use of data transformation was not required.

Data collected from each bean insect pest found were grouped according to the entomological orders for both crops evaluated. Afterwards, statistical analysis was conducted using Sisvar 5.1 software at the 0.05 significance level ($P \leq 0.05$) by the F test. The significant results from the analysis of variance were submitted to regression analysis at the 0.05 significance level ($P \leq 0.05$) as well.

3. RESULTS AND DISCUSSION

Insects found in samplings carried out in the **Rainy Season Crop – 2014** (figure 1) and in the **Dry Season Crop – 2015** (figure2) were counted according to the vegetative and reproductive phenological stages of the crop and according to their entomological classification. From these results, analysis of variance (ANOVA) was performed. The results of the analysis of variance can be observed in (tables 2 and 3).

Table 2. Summary of the analysis of variance (ANOVA). Total of insects counted in samplings carried out during the **Rainy Season Crop – 2014** (figure 1), grouped according to the entomological orders **Diptera (DIP)**, **Hemiptera (HEM)**, **Coleoptera (COL)**, and **Lepidoptera (LEP)**, in the vegetative and reproductive stages of the common bean phenological growth.

Period: vegetative stage of the common bean - rainy season crop (2014)				
Q.M.				
Source of variation	GL	DIP	HEM	COL
Cultivar	1	0.0090 ^{Ns}	0.1323 ^{Ns}	0.9923 ^{Ns}

Block	3	0.8280	0.5883	2.3657
Doses	4	0.4219 ^{Ns}	0.3085 ^{Ns}	1.2594 ^{Ns}
Cultivar x Doses	4	0.2009 ^{Ns}	0.3548 ^{Ns}	0.9591 ^{Ns}
Error	27	0.1769	0.3425	1.0882
C.V. (%)		70.10	89.01	60.91
Overall average		0.6000	0.6575	1.7125

Period: reproductive stage of the common bean - rainy season crop (2014)

Q.M.				
Source of variation	GL	LEP	HEM	COL
Cultivar	1	0.0303 ^{Ns}	10.4040	0.7840 ^{Ns}
Block	3	0.1616	8.8767	1.36267
Doses	4	0.3828 ^{Ns}	0.5984 ^{Ns}	3.3216 ^{Ns}
Cultivar x Doses	4	0.4278 ^{Ns}	0.3846 ^{Ns}	5.9871 ^{Ns}
Error	27	0.3401	1.3633	5.1621
C.V. (%)		77.50	38.16	30.01
Overall average		0.7525	3.0600	7.5700

Q.M (Mean square); GL (Degree of Freedom); C.V. (Coefficient of variation); Ns - not significant; *Significant at (P≤0.05) of probability according to ANOVA test.

Table 3. Summary of the analysis of variance (ANOVA). Total of insects counted in samplings carried out during the Dry Season Crop – 2015 (figure 2), grouped according to the entomological orders Diptera (DIP), Hemiptera (HEM), Coleoptera (COL), and Lepidoptera (LEP), in the vegetative and reproductive stages of the common bean phenological growth.

Period: vegetative stage of the common bean - dry season crop (2015)					
Q.M.					
Source of variation	GL	DIP	HEM	COL	LEP
Cultivar	1	46.6560	0.1288 ^{Ns}	0.0723 ^{Ns}	57.3603 ^{Ns}
Block	3	135.6653	1.1595	0.6423	20.5743
Doses	4	3.3873 ^{Ns}	1.7834	2.5584 ^{Ns}	18.2294 ^{Ns}

Cultivar x Doses	4	11,5460 ^{NS}	0,3299 ^{NS}	0,7891 ^{NS}	53.4434
Error	27	10,5251	0,2076	0,9985	17,0904
C.V. (%)		21,73	53,81	49,41	73,33
Overall average		14,9300	0,8468	2,0225	5,6375

Period: reproductive stage of the common bean - dry season crop (2015)

Q.M.				
Source of variation	GL	LEP	HEM	COL
Cultivar	1	445.5296	4.9000	6768.5567
Block	3	4,8116	7,7723	873,0629
Doses	4	31,0205 ^{NS}	0,8397 ^{NS}	123,5023 ^{NS}
Cultivar x Doses	4	26,1835 ^{NS}	2.4380	970.9799
Error	27	15,5485	0,6752	152,1777
C.V. (%)		34,93	33,31	26,37
Overall average		11,2875	2,4667	46,7832

Q.M (Mean square); GL (Degree of Freedom); C.V. (Coefficient of variation); NS - not significant; *Significant at (P≤0.05) of probability according to ANOVA test.

Tabela 4. Average of the samples per plot. Population of insect pests in the vegetative and reproductive stages of the common bean phenological growth for the Rainy Season Crop – 2014 (figure 1) and the Dry Season Crop – 2015 (figure 2). Assis Chateaubriand, Paraná, Brazil.

Period: reproductive stage of the common bean - rainy season crop (2014)

Entomological Order		Hemiptera	
Cultivars		IPR Campos Gerais	IPR Tuiuiú
Nº of insects		3,57 b	2,55 a
C.V. (%)		38,16	

Period: vegetative stage of the common bean - dry season crop (2015)

Entomological Order		Diptera	
Cultivars		IPR Campos Gerais	IPR Tuiuiú

N° of insects	13,85 a	16,01 b
C.V. (%)	21,73	
Period: reproductive stage of the common bean - dry season crop (2015)		
Entomological Order	N° of insects	
	Lepidoptera	Coleoptera
Cultivars		
IPR Campos Gerais	7,95 a	59,79 b
IPR Tuiuiú	14,62 b	33,78 a
C.V. (%)	34,93	26,37

C.V. (Coefficient of variation). Means followed by the same letter are not significantly different by Tukey's test at (P≤0.05) probability.

As seen in (tables 2 and 4), regarding the experiment conducted in the Rainy Season Crop (2014), a significant difference ($P \leq 0.05$) was observed by ANOVA and Tukey tests between the population means of insect pests for the order hemiptera (table 2), in the cultivars evaluated during the reproductive stage of the crop. IPR Tuiuiú cultivar presented a population of 2.55 hemiptera insects per sampling, while IPR Campos Gerais cultivar presented a population of 3.57 hemiptera insects (table 4).

Significant differences ($P \leq 0.05$) were also observed by the Tukey test between the common bean cultivars in the vegetative stage, during the dry season of 2015, as seen in table 3. IPR Campos Gerais cultivar presented a population of insects from the order diptera of 13.85, while IPR Tuiuiú cultivar presented 16.01 insects per sampling (table 4).

For the dry season crop, the Tukey test pointed out a significant difference ($P \leq 0.05$) between the means of the cultivars evaluated in their reproductive stage for the orders lepidoptera, hemiptera and coleoptera. The average quantities of insects per sampling point in the IPR Campos Gerais cultivar were 7.95 (lepidoptera), 2.12 (hemiptera), and 59.79 (coleoptera). In the IPR Tuiuiú cultivar, the average quantities of insects per sampling point for the cited orders were 14.62, 2.82, and 33.78, respectively. It is important to emphasize that the evaluated cultivars belong to different genotypes of common bean. The IPR Campos Gerais cultivar belongs to the Pinto bean group, whereas the IPR Tuiuiú belongs to the Black turtle bean group. Thus, different behaviors are observed between them since they present unique genetic characteristics of each material.

The bean cultivars studied have an indeterminate growth habit type II, and its inflorescences arise from axillary buds. Apical bud continues to grow even in the reproductive stage, forming a branch that does not exceed a few centimeters; total plant height reaches approximately 70 cm. The side branches are a bit short and the cultivars present a flowering season range from 15 to 20 days. Pods mature evenly and, in general, plants have a life cycle from 80 to 90 days, with about 3,897 Kg ha⁻¹ of yield potential for IPR Campos Gerais cultivar, and 3,950 Kg ha⁻¹ for IPR Tuiuiú cultivar [10].

For both experiments, the results can probably be assigned to the genetic basis of the cultivars studied, since different common bean genotypes are concerned. Results obtained in the experiments conducted during rainy season (2014) and dry season (2015) crops made possible to observe contestations among the genotypes, suggesting a difference in the preference of the insect pests for the evaluated materials. Plants resistance to insects is determined by chemical, physical and morphological factors that probably act separately or jointly. Such factors provide plant with different degrees of resistance, especially through toxins, digestibility reducers, trichomes, hardness of the leaf epidermis, and nutritional improprieties as present in common bean genotypes [11].

Another study evaluated the behavior of 19 bean genotypes [12], focusing the infestation of *Bemisia tabaci* biotype B in rainy, dry and winter season crops. It concluded that genotypes IAC Una, Pérola, Gen 96A45-3-51-52-1, Gen 96A98-15-32-1, FT Nobre, IAC Tybatã, IAC Alvorada, LP 02-130, LP 01-38, LP 98-122, IAC Diplomata and Gen 96A3P1-1-1 are less ovoposited by *B. tabaci* biotype B in the experiment conducted during rainy season crop. That study also concluded that the highest incidence of nymphs occurred in the experiment conducted during dry season crop. The present

study observed the same dynamics regarding the attraction of the insect pests analyzed to the bean cultivars evaluated, as seen in the tables presented above.

The highest insect pest population indices in the common bean crop, apart from atypical climatic changes or environmental balance alterations, have been observed in the warmer months [13]. Such pattern of occurrence negatively affects the sowing of the dry season crop since the problems with pests at that season intensify from the beginning of the crop development. This can be seen in this study, where statistically significant differences were observed ($P \leq 0.05$) by Tukey test regarding the insect pest orders hemiptera and lepidoptera evaluated for the cultivars concerned (table 4).

During the conduction of this study, as observed in the (figures 1 and 2), meteorological conditions unfavorable to the development of the bean occurred, with temperatures above 30°C and high relative humidity of the air above 75% during the entire crop growth stage. It is also worth mentioning that rainfall in this period was irregular, presenting an uneven water distribution during the physiological growth stages of the common bean cultivars. Environments under conditions similar to those observed in this study are considered favorable to the reproduction of insect pests, thus justifying the high population of the insects from the entomological orders evaluated.

There was a significant effect ($P \leq 0.05$) of probability, by the Tukey test, regarding potassium silicate doses evaluated for the order hemiptera during the vegetative stage of the Dry Season Crop, (2015). It is important to note that insect pests from the order hemiptera usually do not cause damage to bean culture when it is in vegetative stages of phenological growth. Damages are commonly observed during the reproductive growth stages, when plants present formation of flowers and pods and grains arise. In this case, insects from this order begin to cause damage to the crop, providing abortion of flowers and pods, besides causing uneven formation of grains. The main bean insect pest from the order hemiptera is Bedbug [14].




Figure 3. Number of Hemiptera in relation to treatments with doses of potassium silicate. Vegetative stage of the common bean (*Phaseolus vulgaris* L.) – dry season crop (2015).

As seen in (figure 3), significant results ($p \leq 0.05$) by the Tukey test were found due to the foliar application of potassium silicate, which in contact with the surface of the plant tissue formed a silica compound that is not absorbed by plants, called polysilicic acid.

By working with silicon in solution, as its concentration increases, long-chain polymers are created from monosilicic acid (H_4SiO_4), forming polysilicic acid which easily polymerizes in acid environment [15]. Thus, the most likely is that this compound provides plants with an external protective barrier on the leaves to the pest attacks, causing differentiation regarding the population of insects according to the treatments used. In this situation, formation of potassium silicate-polymer external to the plant tissue occurs, which hinders insect pests from feeding on plants.

Interaction between cultivars x doses was also analyzed for the occurrence of the orders lepidoptera (vegetative stage of the crop phenological growth), hemiptera and coleoptera (reproductive stage of the crop phenological growth) during the dry season crop of 2015 (figures 4, 5 and 6).




Figure 4. Lepidoptera population in the vegetative stage of common bean (*Phaseolus vulgaris* L.) by applying doses of potassium silicate – dry season crop (2015).




Figure 5. Hemiptera population in the reproductive stage of common bean (*Phaseolus vulgaris* L.) by applying doses of potassium silicate – dry season crop (2015).




Figure 6. Coleoptera population in the reproductive stage of common bean (*Phaseolus vulgaris* L.) by applying doses of potassium silicate – dry season crop (2015).

Such interaction may have resulted in an external layer of silica on leaves, working as a protective layer that reduced the insect attacks. This justifies the genetic characteristics (color of leaves, plant height, dry matter, cellulose content, among others) found in the common bean evaluated.

When analyzing the climate (meteorological behavior) during the course of the study (2014 and 2015) and the phenological development of materials in the cultivation environment of both crops (figures 1 and 2), differences observed in each crop demonstrated that adverse environmental conditions (especially rainfall and temperature) may cause changes in the insect pests behavior and population dynamics at different stages of the phenological growth.

No significant effect was observed regarding potassium silicate application for some insect pest orders evaluated (tables 2 and 4). It probably happened because the experiment was conducted in a clayey soil (73.5 % clay). The soluble Si content is higher in that type of soil. Red Latosol soils are very weathered, with higher quantity of clay [16]. The location where the experiment was conducted presents 19.6 mg dm³ of soil soluble Silicon, a value considered to be very high. This is the reason why Si would hardly demonstrate its effects on the crop, mainly by the fact that common bean is a silicon-non-accumulating plant.

When Si is found in the liquid phase of the soil as a monocyclic acid, it is absorbed by plant roots through passive transport, which occurs when soil nutrient moves to the root surface in favor of the concentration gradient, requiring no expenditure of metabolic energy by the plant. Thus, nutrient moves from the highest concentration area (rhizosphere) to the lowest concentration area (root) [17].

Plants can absorb mineral more easily through root system, since it has a greater quantity of membrane transport proteins by which molecules and ions can diffuse [18]. Consequently, plants absorb Silicon by the roots with less energy expenditure, not justifying its foliar absorption - which would require the plant to spend more metabolic energy.

It is important to emphasize that Latosol soils contain a large quantity of kaolinite, which still undergoes weathering action and consequently releases soluble Si to soil [19].

4. CONCLUSION

Foliar fertilization with potassium silicate caused a population decrease of the entomological orders evaluated for the common bean genotypes studied. During the reproductive stage of the Rainy Season Crop, 2014, there was a significant difference ($P \leq 0.05$) between the bean genotypes for the order hemiptera. In the vegetative stage of the Dry Season Crop, 2015, there was a significant difference ($P \leq 0.05$) between the bean genotypes for the order diptera. Potassium silicate also caused a population decrease of the entomological orders hemiptera and lepidoptera in the vegetative stage

of the Dry Season Crop, 2015, for cultivar x doses interaction. In addition, there was a significant difference ($P \leq 0.05$) in the reproductive stage of the Dry Season Crop, 2015, for the entomological orders lepidoptera, hemiptera, and coleoptera between the genotypes evaluated.

REFERENCES

[1] King ABS, Sanders JL. Las plagas invertebradas de cultivos anuales alimenticios en América Central: una guía para su reconocimiento y control. Turrialba: CATIE, London: TDRI; 1984. Spanish

[2;3] Quintela ED, Barbosa FR. Manejo de pragas. In: Carneiro JES, Paula Junior T, Borém A editors. Feijão: do plantio à colheita. Viçosa: UFV; 2015. Portuguese

[4] Alves AL. Adubação foliar com silicato na ocorrência de pragas e características agrônômicas do feijoeiro comum. 2016. 69 f. Dissertação (Mestrado em Agronomia) - Universidade Estadual do Oeste do Paraná, Marechal Cândido Rondon, 2016. Accessed 9 September 2018. Available: <http://tede.unioeste.br/handle/tede/1327>. Portuguese

[5] Arnon DL, Stout PR. Molybdenum as an essential element for higher plants. Plant Physiology, Rockville, 1939;14:3599-02, 1939. Available in: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC437771/pdf/plntphys00301-0192.pdf>. Accessed 19 August 2018.

[6] Marschner H. Mineral nutrition of higher plants. 2th ed. London: Academic Press; 1995.

[7] Malavolta E. Manual of mineral nutrition of plants. São Paulo: Editora Ceres; 2006. Portuguese

[8] Gomes FB, Moraes JC, Neri DKP. Adubação com silício como fator de resistência a insetos-praga e promotor de produtividade em cultura de batata em sistema orgânico. Ciência e agrotecnologia, Lavras, 2009;33:1. Accessed 15 September 2018. Available: <http://dx.doi.org/10.1590/S1413-70542009000100002>

[9] QUINTELA, E. Manejo integrado de pragas do feijoeiro. In: Curricular Técnica, 46. Santo Antonio de Goiás: Embrapa Arroz e Feijão, 2001. Accessed 15 September 2018. Available in: <https://www.embrapa.br/documents/1344498/2767895/manejo-integrado-de-pragas-do-feijoeiro.pdf/c8bb5013-3bf8-4579-a9ea-64570cb70e90>.

[10] INSTITUTO AGRONÔMICO DO PARANÁ (IAPAR). Main characteristics of the bean cultivars with seeds available in the market. Accessed 1 June 2015. Available in: <http://www.iapar.br/modules/conteudo/conteudo.php?conteudo=1363>.

[11] Lara FM. Princípios de resistência de plantas a insetos. 2. ed. São Paulo: Icon; 1991. Portuguese

[12] Jesus FG, Boiça Junior AL; Carbonell SAM, Stein CP, Pitta RM, Chiorato AF. Infestation of Bemisia tabaci biotype B and Caliothrips phaseoli in genotypes of Feijoeiro, Bragantia, São Paulo, v. Brazil. 2010;69(3)637-48. Accessed 17 July 2018. Available: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0006-87052010000300016

[13] Comissão Técnica Sul-brasileira de Feijão- CTSBF. Informações técnicas para o cultivo de feijão na Região Sul brasileira. Florianópolis: Epagri; 2012. Accessed 13 November 2015. Available: http://www.epagri.sc.gov.br/wpcontent/uploads/2013/10/informacoes_tecnicas_cultivo_feijao.pdf. Portuguese

[14] Quintela ED. Manejo integrado dos insetos e outros invertebrados pragas do feijoeiro. Informe agropecuário, Belo Horizonte 2004; 223:25. Accessed 15 September 2018. Available: <https://pt.slideshare.net/carpen2/informe-agropecuário-feijo>. Portuguese

[15] Iler RK. The chemistry of silica. New York: Wiley Interscience; 1979.

381 [16] Camargo MS, Pereira HS, Korndorfe GE, Reis GE. Soil reaction and absorption of silicon by
382 rice. Scientia Agricola, Piracicaba, 2007;64:2. Accessed 18 august 2018. 2007. Available:
383 <http://dx.doi.org/10.1590/S0103-90162007000200011>.
384
385 [17];[18]Taiz L, Zeiger E. Fisiologia Vegetal. 5. ed. Porto Alegre: ARTMED; 2013. Portuguese
386
387 [19]LIMA, H. N. Genesis, chemistry, mineralogy and micromorphology of soils of the Western
388 Amazon. Thesis (Doctorate) – Universidade Federal de Viçosa, Viçosa, 176 fls, 2001. Accessed 18
389 March 2018. Availablein:
390 <http://www.locus.ufv.br/bitstream/handle/123456789/7934/texto%20completo.pdf?sequence=1&isAllo>
391 [wed%20=%20y](http://www.locus.ufv.br/bitstream/handle/123456789/7934/texto%20completo.pdf?sequence=1&isAllo).
392