

Influence of arbuscular mycorrhizal fungi and phosphorus doses in the production of *Parkia nitida* (Miquel) in seedling nursery in the south of Amazonas

Aims: Little is known about the Arbuscular mycorrhizal Fungi (AMF) inoculation effects on the initial development of *Parkia* sp. and phosphorus nutritional needs. This study aimed to evaluate the effect of the AMF inoculation in different phosphorus doses on initial *Parkia nitida* growth of. **Study design:** The study was carried out in a completely randomized design in a 5x2 factorial scheme (5 doses of phosphorus in the absence and presence of AMF inoculation). Inoculation was done by applying 5 ml of inoculum soil containing a mix of spores (70 spores/ml) of the species *Acaulospora longula*, *Acaulospora morrowiae*, *Dentiscutata heterogama* and *Paraglomus occultum*. **Place and duration of study:** Experiment occurred at the Amazonas Federal Institute of Education, Science and Technology - Campus Lábrea (7°15'02.9"S 64°47'05.5"W) between January 2017 and December 2017. **Methodology:** The variables analyzed were the Survival Rate (SR), height (h), lap diameter (d), Seedling Quality Index (h/d), leaflet area (LFA), leaflet number (LFN), shoot dry mass (SDM), dry mass of the root (DMR), DMPA/DMR ratio, total dry mass (TDM) and Dickson Quality Index (DQI). **Results:** The variables h and h/d were significant ($P=.05$) for the interaction between factors only for the treatment without inoculation. It was not possible to determine the appropriate phosphorus dose for the *Parkia nitida* seedlings production. However, aerial part of the seedlings positively responded to the phosphorus doses increase. **Conclusion:** Inoculation did not show significant results in the experimental conditions as well as in the phosphorus interaction.

Keywords: Amazonian Wood; Degraded areas; Dormancy breaking; Faveira-pé-de-arara.

1. INTRODUCTION

One of the greatest difficulties for the Amazonian tree species use in agroforestry systems, afforestation, restoration and recovery of degraded areas, is to obtain seedlings on commercial scale presenting good quality. *Parkia nitida*, popularly known as "Faveira-branca", is a neotropical tree of great economic and ecological importance [1], a leguminous tree, can be considered a specie with great potential to achieve several of these goals. *Parkia nitida* (Fabaceae Mimosoideae), The species that occurs in Brazil is frequent in the Amazon and Central Region of the country, in "terra firme" forests and high varzeas of clay soils [2]. It is a large tree that can reach up to 40 m in height, being widely distributed from southern Panama to the central region of the Amazon according to Moraes, Ferraz & Procópio (2015) [1]. Due to the great economic and ecological interest, *Parkia nitida* is recommended for reforestation and recovery of degraded areas [3].

25 For degraded areas, leguminous tree species are prominent because they present rapid growth and,
26 in association with nitrogen-fixing bacteria, can supply a large amount of this nutrient [4]. In regions
27 such as the Amazon South where the difficulty in acquiring fertilizers and correctives limits regional
28 development, having the possibility of producing vigorous seedlings and with local resources can
29 change such reality. The classes of soils of prominence in the Amazon are classified as Latosols,
30 Argisols, Plinthosols and Spodosols [5]. Soils are generally acidic and nutrient poor, mainly
31 phosphorous (P). Acidity correction is performed by liming, but logistical difficulties and cost make this
32 soil management practically unfeasible in the region. Phosphorus is an integral component of
33 important plant cell compounds, including phosphate-intermediate sugars from respiration and
34 photosynthesis, as well as the phospholipids that make up the cell membrane and nucleic acids [6].
35 Phosphorus is a critical element for food production systems, manufacturing industries and general
36 economic growth, generating long-term supply insecurity concerns for regional and national
37 economies [7].

38 Arbuscular mycorrhizal fungi (AMF) belong to Glomeromycota phylum and are associated with the
39 majority of vascular plants according to **Moreira & Siqueira (2006)** [4]. Accordint to Celebi et al (2010)
40 [8] AMF are ecologically significant because they contribute to relationships in and on the roots of the
41 plant in a symbiotic association. When associated with plants roots, they increase their exploitation
42 area, influencing their capacity to absorb water and nutrients, especially those with low mobility such
43 as phosphorus [9], promoting greater resistance to biotic and abiotic stress, therefore, the plant
44 provides AMF with soluble carbon sources and AMD provides the plant with an increased capacity to
45 absorb more water and nutrients from the soil. Therefore, according to Kapulnik et al (2010) [10] AMF
46 increases plant resistance to stress, enhancing productivity and sustainability of crops, especially
47 under prolonged stress conditions, like drought.

48 The quality of seedlings produced is a major factor for the success of forest stands [11]. The benefits
49 of symbiosis for seedlings depend on the specific combination between fungal isolate and cultivated
50 plant [12]. According to **Lacerda et al (2011)**, it is widely accepted that seedlings mycorrhization favors
51 its nursery initial growth and its field adaptation and establishment [13], especially the perennial, like
52 the fruit trees [14]. Mycorrhizal inoculation is generally more efficient than the use of high fertility
53 substrate to promote plant growth and reduce stress after transplant and may also act as biological
54 control.

55 As a result, the use of mycorrhizal fungi is an alternative for the optimization of fruit tree seedlings,
56 since it allows seedling formation time reduction, increasing nursery productivity, rotation in the
57 occupation of infrastructure and the efficiency of specialized staff [15]. However, little is known about
58 *Parkia nitida* nutritional requirements and ecological relationships, such as their ability to form
59 symbioses with fungi from Glomeromycota phylum [13].

60 In the Amazonian environment P has low availability, since both the forest and the soil are drains. The
61 tropical forest has 54.5 kg ha⁻¹ of P immobilized in its biomass, with cycling values of the order of 17
62 kg ha⁻¹ [16]; this balance is also derived from interactions and biological activities and these are
63 essential for the maintenance of the climactic situation of the vegetation [17]. One of the biological
64 factors directly linked to this success are the Arbuscular Mycorrhizal Fungi (AMF). This is in line with
65 the research by Cebeli et al (2010) [8] where AMF application increased leaf and maize ear ratio
66 generally and a decrease in stem ratio. Therefore, strategies aimed at maximizing fertilization are
67 recommended with the use of arbuscular mycorrhizal fungi. Typically, higher concentrations of P in
68 soil inhibit fungal colonization in the roots and lower concentrations, instead, favor colonization
69 according to Smith et al (2011) [18]. Thus, the highest values of colonization by fungi are found in the
70 roots submitted to soils of low concentration of P in the applied nutrient solution, in comparison to the
71 treatment with high level of P.

72 Considering the logistical difficulties of southern Amazonas, the study sought to present a new,
73 efficient and inexpensive way of producing seedlings in nursery, specifically *Parkia nitida*, which
74 represents, besides a cheap solution, also has great potential for the recovery of degraded areas.
75 Therefore, this study aimed to evaluate the effect of inoculation with Arbuscular Mycorrhizal Fungi in
76 different doses of phosphorus on the initial growth of *Parkia nitida* in southern Amazonas.

77 78 **2. MATERIAL AND METHODS**

79
80 The study was conducted in southern Amazonas, Lábrea municipality, which has a total area of 6,000
81 ha [19]. Experimental conduction occurred at the Amazonas Federal Institute of Education, Science
82 and Technology - Campus Lábrea (7 ° 15'02.9 "S 64 ° 47'05.5" W).
83 Seedlings substrate was obtained through the mixture of soil, sand and semi-composite sawdust in
84 the ratio of 5:3:2 (v/v) trait adapted from Santos et. al (2006) [20], previously sieved with a 4 mm mesh

85 to homogenize. Sand originated from the Purus River and the sawdust was obtained from a local
86 logging.

87 The soil used was classified according to the Brazilian Soil Classification System [21] as Flavic Type 2
88 Neosol (average texture = 35% clay, 50% silt and 15% Sand) with the following analytical results: pH
89 4.2; K: 17.10 mg.dm⁻³; P: 1.07 mg.dm⁻³; Ca: 0.25 cmol.dm⁻³; Mg: 0.09 cmol.dm⁻³; Al: 3.20 cmol.dm⁻³;
90 H+Al: 18.73 cmol.dm⁻³; SB: 0.38 cmolc.dm⁻³; t: 3.58 cmolc.dm⁻³; T: 19.11 cmolc.dm⁻³; V: 2.01 %, m:
91 89.39 %; O.M: 2.58 dag.kg⁻¹; P-rem: 8.31 mg.L⁻¹; Zn: 0.10 mg.dm⁻³; Fe: 227.22 mg.dm⁻³; Mn: 0.91
92 mg.dm⁻³; Cu: 0.23 mg.dm⁻³; B: 0.12 mg.dm⁻³; S: 224 mg.dm⁻³ and the extractor was Mehlich 1 (P – Na
93 – K – Fe – Zn – Mn – Cu).

94 The correction was performed considering raising saturation to 50% using dolomitic limestone with
95 relative total neutralization of 92.54% following literature recommendations [8]. After this procedure,
96 substrate was homogenized. In the fertilization, granulated super-phosphate (P₂O₅) was used in 5
97 concentrations as follows: 0, 20, 40, 80 and 160 mg of P₂O₅ L⁻¹ substrate, according to literature [22].
98 Arbuscular mycorrhizal fungi spores samples were obtained in soils collected in a Brazil-nut tree
99 *Bertholletia excelsa* dominated forest fragment, located at the Transamazônia Highway (7 ° 28'15.3
100 "S 64 ° 39'26.3" W). Arbuscular mycorrhizal fungi spores were extracted through wet sieving method
101 [23]. Spores were conditioned in sterilized soil in 3 kg culture pots, aiming to its multiplication for an 8
102 month period [24].

103 Seed dormancy breaking was adapted from Brito et al (2017) [25]. Mechanical scarification was
104 performed using abrasive rotary grinding wheel coupled to a bench-mounted hand drill with a
105 transverse insertion in the seed. This procedure results in a small removal of part of the tegument
106 laterally related to the thread until reaching the primary endosperm. Subsequently, they were soaked
107 in water for 24 hours for the activation of germination enzymes [26;27].

108 The sowing occurred in two 200 and three 128 cell trays, filled with sand less than four millimeters in
109 size and covered with black shading screen with 50% luminosity at a height of 10 cm in relation to the
110 tray surface. Irrigation took place over 10 days at intervals of 10 hours each day, after which the
111 seedlings were immersed in water to facilitate the trays removal, avoiding roots damage. These were
112 selected and pruned, leaving only five centimeters in roots length.

113 Germination of *Parkia nitida* is epigeal type and phanerocotyledonar [28]. The substrate used for this
114 stage was adequate, as observed for arboreal legume [29], the shading maintained adequate

115 moisture and solar incidence. A total of 494 seedlings were obtained, representing 63% of the seeds
116 used. There was no significant difference between the trays types. Germination rate for 10 days was
117 close to values obtained by the regression model observed by Cruz (2001) [28] using a similar
118 method of scarification for dormancy breaking. Seedlings present desirable phenotypic
119 characteristics. Propagation of nursery seedlings exceeds the direct germination rates previously
120 observed [30] even for high rates.

121 A total of 300 plastic bags (18x25 cm) were prepared with 1 L of substrate for seedlings. Sixty
122 seedlings plastic bags were fertilized for each dose of P_2O_5 , half of which was inoculated with
123 Arbuscular Mycorrhizal Fungi and the other half remained inoculum-free.

124 Transplanting was done to 6 cm depth pits and the roots of the seedlings were pruned to 5cm for
125 standardization. Inoculation was carried out in the pit, trait adapted from the research by Meir et al
126 (2010) [31] through the application of 5 ml inoculum soil (colonized roots, pieces of hyphae, spores
127 and soil), containing a mixture of *Acaulospora longula*, *Acaulospora morrowiae*, *Dentiscutata*
128 *heterogama* e *Paraglomus occultum* spores (70 spores.ml⁻¹).

129 The substrate used to produce the seedlings showed to be permeable and slightly hard. The
130 permeability was due to the introduction of the sand trace and the hardness was related to the clay
131 type of the soil used. Normally, the substrates used in the production of seedlings (coffee and fruit)
132 have 30% of manure as an increase to organic matter. Proportion adaptation [32] of 20% of organic
133 matter was used, in this case from sawdust that did not undergo chemical and/or physical
134 characterization, which makes inferences about the degree of natural degradation impossible and it
135 can acidify the substrate besides having tannins that can affect the symbiosis. However, the
136 production of seedlings in a conventional way largely uses sawdust and, in the Amazon region, closer
137 to the cities, it is a co-product with great availability, which does not occur with animal manures.

138 The effect of AMF in interaction with the dose of P_2O_5 was evaluated by the inoculation of half of the
139 seedlings, while the other half was not inoculated. In order to evaluate the different concentrations of
140 phosphorus in the mycorrhizal process, the work was carried out based on previously experiment by
141 Lacerda et. al (2001) [11], with 5 treatments in relation to the P, as follows: T1 – control, without
142 addition of P_2O_5 ; T2 - addition of 20 mg.L⁻¹ P_2O_5 content; T3 - addition of 40 mg.L⁻¹ P_2O_5 content; T4 -
143 addition of 80 mg.L⁻¹ P_2O_5 content, and T5 - addition of 160 mg.L⁻¹ P_2O_5 content.

Experimental design was done in randomized blocks, with factorial scheme 5x2, in the presence and absence of AMF, with five variations in the phosphorus dose. Each plot was constituted by one seedling.

Data were collected by evaluating the height (h), leaf area (LA), shoot dry mass (SDM), dry mass of the root (DMR) and total dry mass (TDM) [33].

At the end of the experiment, aerial part and the root system were separated to determine the dry matter biomass (n=60). Samples were packed in paper bags and then dried at 70°C/48h (forced air circulation drying oven). To obtain the SDM and DMR, samples were taken from the oven, placed in desiccator to reach room temperature and then weighed in an analytical balance (0.001 g accuracy). TDM seedling was obtained by the sum of the root and aerial part dry matter weights.

The following variables were observed to analyze the seedlings development: i) length of the aerial part or height (h): graduated ruler, being taken as standard the terminal bud - apical meristem; ii) Collar diameter (d): Measurement above 1 cm of the node formed just above the soil surface of the vessel, with the aid of a digital caliper; iii) relation of height and diameter of the collar (h/d): ratio between the height of the seedling and its diameter of the collar allowed to obtain a quality index of seedlings; iv) shoot dry mass part and dry mass of the root ratio (SDM/DMR): quality index of seedlings; v) Survival rate (SR): survival percentage; vi) Dickson quality index (IQD) [34]: determined as a function of aerial part (h), collar diameter (d), shoot dry mass (SDM) and dry mass of the root (DMR).


Data from the last collection were used (153 days after transplanting). Observations of the last collection were submitted to the individual and joint analysis of variance (factorial 5x2), in which the effects of systems were evaluated by the F-test and the effects of doses by the Tukey's test. At the same time, the significance of the model ($P < 0.05$) and the coefficient of determination (R^2). Data was submitted to statistical tests with the aid of Sisvar software.

3. RESULTS AND DISCUSSION

In the production of *Parkia* seedlings commercially, soil sterilization was not performed, which facilitates root colonization by AMF, which may interfere with the inoculation results.

P doses had a significant effect in the absence of AMF (Fig. 1), the peak dose was 98.67 mg L⁻¹ of P, and this enabled a larger increase in h.

175



176 **Fig. 1 Height (h) of *Parkia nitida* seedlings as a function of arbuscular mycorrhizal fungi (FMA)**
 177 **and phosphate fertilization, 153 days after transplanting**


178 The curve behavior observed (Fig. 2) resembles the one described previously by Shimizu [et. al \(2011\)](#)
 179 [27] in the treatment without fungus that involved phosphate fertilization, by the curvilinear behavior,
 180 the elevation of the dose of phosphorus above the maximum point does not provide significant
 181 increases for the variables h, which meets the maximum law [in line with Vasconcellos et. al \(2001\)](#)
 182 [35].

183 At 153 days after transplanting, the mean height of the seedlings was 23.6 cm and the mean collar
 184 diameter was 6.5 mm, the height for forest species should be superior to 50 cm [36]. Observing this
 185 data, it is possible that the maintenance of the seedlings for a longer period in the greenhouse allows
 186 positive increases in its quality.

187 The seedling quality index that relates height per diameter showed an average of 3.6 (Fig. 2), showing
 188 that there was a higher growth rate in height in relation to the diameter.

189 In order to take the seedlings to field planting, a balanced development is necessary, as observed in
 190 [Campos et. al \(2002\)](#) [37]. This index only underwent significant interaction of the following sources of
 191 variation: doses of phosphorus in the absence of fungus.

192




193 **Fig. 2 Quality index of height per diameter (h/d) of *Parkia nitida* seedlings as a function of**
 194 **arbuscular mycorrhizal fungi (AMF) and phosphate fertilization, 153 days after transplanting**

195 The variables h and h/d presented significance for the interaction of factors (dose of phosphorus per
 196 fungus). However, this occurred only for treatments in the absence of AMF, evidencing that further
 197 studies with other fungal species are necessary to determine the ability or not of mycorrhizae of the
 198 *Parkia nitida* plant and its effects on the development of the species.


199 The leaflet area (LA) was significantly higher at higher P rates. A linear increase with a coefficient of
 200 determination of 89.07% was observed (Fig.3). This behavior of increment in the leaflet area by
 201 phosphorus dose is in line with **Rodrigues et al (2003)** [38], since phosphorus is linked to the
 202 photosynthetic capacity of the plants, so it may be positively influencing.

203



204 **Fig. 3 Influence of phosphate fertilization on the total leaflet area of *Parkia nitida* seedlings at**
 205 **153 days after transplanting.**


206 Another verified index was SDM/DMR, which averaged 3.16, higher than the 2.0, that is established
 207 as reference to this index consistent with the finding of research by Gomes et al. (2002) [39],
 208 however, studies directed to *Parkia nitida* are still necessary to determine the specific value. It is
 209 possible that the roots do not have adequate photoassimilated resources for their development.
 210 The development of the leaflets in the area by increment in the dose of P reflected in the SDM (Fig.
 211 4). The LA is closely related to SDM and TDM, since the gain was in both area and mass.
 212
 213



214 **Fig. 4 Influence of the phosphate fertilization on the shoot dry mass part of the *Parkia nitida***
 215 **seedlings at 153 days after transplanting.**

216 The total dry mass (TDM) (Fig. 5), being a composition of variables, was influenced by the dry mass
 217 of the root (DMR). The TDM had a lower coefficient of determination than that of the SDM showing
 218 this interference. On average, the DMR was 1.57 g, corresponding to less than half of the SDM. It is
 219 verified in the literature that the DMR is directly linked to the availability of P. In the experimental
 220 conditions, the result of this study indicated that the plant priorities the development of the aerial part
 221 and it is possible that higher doses of phosphorus improve the indexes of quality like the DQI, which
 222 presented average of 0.96.

223



224 **Fig. 5 Influence of phosphate fertilization on total dry mass of *Parkia nitida* seedlings at 153**
 225 **days after transplanting.**

226
 227 The behaviour of *Parkia nitida* related to doses of P is increasing and linear for the variables LA, SDM
 228 and TDM, which evidences the necessity of application of P for better results in the seedlings
 229 production.

230 *Parkia nitida* presents desirable characteristics for propagation in nursery, with emphasis on its
 231 physiological resistance. The seedling survival rate (SR) was 100%, demonstrating that root pruning
 232 at transplant did not interfere with survival according to Azevedo et al. (2010) [40].

233 The interactions in the treatments that obtained significance were those that did not undergo
 234 inoculation by inoculum soil. According to the research by Kapulnik et al (2010) [10], the ability to
 235 respond noticeably to AMF inoculation depends not only on plant genotype but also on the identity of
 236 the AMF inoculated. Seedlings used in this work come from seeds and, therefore, may have varied
 237 genotypes. It is possible that the genetic variability of *Parkia nitida* seeds is high. This is a desirable
 238 feature for species destined for floristic recovery and recomposition. For degraded areas, the lower
 239 the genotype/environment interaction, the greater the possibility of success.

240 For commercial production of “Faveira-branca” seedlings it is recommended the phosphate
 241 fertilization for better development of the plant.

242 AMF form a symbiotic mutualistic association with the roots of most tropical plants and extend the
 243 area of exploitation in the soil favoring the absorption of water and nutrients, mainly phosphorus.
 244 Although this association is well studied, little is known about the effects of FMA inoculation on the
 245 initial development of *Parkia* genus plants and mainly on their nutritional need for phosphorus. In
 246 regions such as the South of the Amazon where the difficulty in acquiring fertilizers and correctives

limits regional development, having the possibility of producing vigorous seedlings and with local resources can change this reality. Therefore, this manuscript hopes to contribute to new research that will help reduce these limitations and also, contribute to studies related to the use of FMA specifically in the development of *Parkia nitida*.

4. CONCLUSION

The species *Parkia nitida* (Miquel) presented a small effect of inoculation with arbuscular mycorrhizal fungi on initial development in the nursery.

The seedlings of *Parkia nitida* showed linear growth, in the biomass and in the leaflet area, with increasing doses of phosphorus, reaching peak at 98.67 mg L⁻¹ of P.

ACKNOWLEDGEMENTS

The authors are grateful to the Federal Institute of Education, Science and Technology of Amazonas for funding part of the research and for making the infrastructure available. The first author thanks Federal University of Lavras for the opportunity to attend the master's degree.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

AUTHORS' CONTRIBUTIONS

Author Hugo Cesar Tadeu designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors Marco Aurélio Carbone Carneiro and Márcio Rodrigues Miranda guided the research and assisted in writing the final manuscript. Authors Leandro Coutinho Alho, Pedro Italiano de Araújo Neto e Álefe Lopes Viana assisted in literature research, field activities and laboratory tests. All authors read and approved the final manuscript.

REFERENCES

1. Moraes GJVP, Ferraz IDK, Procópio LC. Imaturidade fisiológica e condicionamento hídrico de sementes de *Parkia nitida* Miq com dormência física. *Ciência Florestal*. 2015;25(4): 1053-1059. Portuguese. DOI: 10.5902/1980509820667.
2. Carrero GC, Pereira RS, Jacaúna MA, Vieira Junior MJ. Árvores do Sul do Amazonas: guia de espécies de interesse econômico e ecológico. Manaus: IDESAM; 2017.
3. Costa KCP, Ferraz JBS, Bastos RP, Reis TS, Ferreira MJ, Guimarães GP. Estoques de biomassa e nutrientes em três espécies de *Parkia* em plantios jovens sobre área degradada

284 na amazônia central. Floresta. 2014;44(4): 637-646. Portuguese. DOI:
285 10.5380/rf.v44i4.34135

286 4. Moreira FMS, Siqueira JO. *Microbiology and soil biochemistry*. Lavras: UFLA; 2006.

287 5. Vale Junior JF, Souza MIL, Nascimento PPRR, Cruz DLS, Solos da Amazônia: etnopedologia
288 e desenvolvimento sustentável. Revista Agroambiente. 2011;5(2): 158-165. Portuguese. DOI:
289 10.18227/1982-8470ragro.v5i2.562

290 6. Pereira IS, Nava G, Picolotto L, Vignolo GK, Gonçalves MA, Antunes LEC. Exigência
291 nutricional e adubação da amoreira preta. Revista de Ciências Agrárias. 2015; 58(1): 96-104.
292 Portuguese. DOI: 10.4322/rca.1755

293 7. Withers PJA, Elser JJ, Hilton J, Ohtake H, Schipper WJ, Van Dijk KC. Greening the global
294 phosphorus cycle: how green chemistry can help achieve planetary P sustainability. Green
295 Chemistry. 2015;17(4): 2087–2099. DOI: 10.1039/C4GC02445A

296 8. Ribeiro AC, Guimarães PTG, Alvarez VH. Recommendations for the use of correctives and
297 fertilizers in Minas Gerais: 5th approach. Viçosa: UFV; 1999.

298 9. Celebi SZ, Demir S, Celebi R, Durak ED, Yılmaz IH. The effect of Arbuscular Mycorrhizal
299 Fungi (AMF) applications on the silage maize (*Zea mays* L.) yield in different irrigation
300 regimes. European Journal of Soil Biology. 2010; 46(5): 302-305. DOI:
301 <https://doi.org/10.1016/j.ejsobi.2010.06.002>

302 10. Kapulnik Y, Tsrer L, Zipori I, Hazanovsky M, Wininger S, Dag A. Effect of AMF application on
303 growth, productivity and susceptibility to Verticillium wilt of olives grown under desert
304 conditions. Symbiosis. 2010; 52 (2): 103-111. DOI: [https://doi.org/10.1007/s13199-010-0085-](https://doi.org/10.1007/s13199-010-0085-z)
305 [z](https://doi.org/10.1007/s13199-010-0085-z)

306 11. Alves AS, Lima VLA, Medeiros SS. Desenvolvimento inicial de mudas de *Parkia platycephala*
307 *Benth.* em diferentes tamanhos de recipientes. Revista Verde. 2012;7(2):206-2011.
308 Portuguese. DOI: 10.18378/rvads.v7i2.1262

309 12. Anzanello R, Souza PVD, Casamali B. Fungos micorrízicos arbusculares (FMA) em porta-
310 enxertos micropropagados de videira. Bragantia. 2011;70(2):409-415. Portuguese. DOI:
311 10.1590/S0006-87052011000200023

312 13. Lacerda KAP, Silva MMS, Carneiro MAC, Reis EF, Saggin Júnior OJ. Fungos micorrízicos
313 arbusculares e adubação fosfatada no crescimento inicial de seis espécies arbóreas do
314 Cerrado. Cerne. 2011;17(3):377-386. Portuguese. DOI:10.1590/S0104-77602011000300012

315 14. Zemke JM, Pereira F, Lovato PE, Silva AL. Avaliação de substratos para inoculação
316 micorrízica e aclimatização de dois porta-enxertos de videira micropropagados. Pesquisa
317 Agropecuária Brasileira. (2003);38(11):1309-1315. Portuguese. DOI: 10.1590/S0100-
318 204X2003001100009

319 15. Silveira APD, Silva LR, Azevedo IC, Oliveira E, Meletti LMM. Desempenho de fungos
320 micorrízicos arbusculares na produção de mudas de maracujazeiro-amarelo, em diferentes
321 substratos. Bragantia. 2003;62(1):89-99. Portuguese. DOI: 10.1590/S0006-
322 87052003000100012

16. Clevelario Junior J. Distribution of carbon and mineral elements in a low montane tropical rainforest ecosystem. Viçosa: UFV; 1996.
17. Novais RF, Alvarez VH, Barros NF, Fontes RL, Cantarutti RB, Neves JCL. Soil fertility. Viçosa: Sociedade Brasileira de Ciência do Solo; 2007
18. Smith SE, Jakobsen I, Grønlund M, Smith FA. Roles of arbuscular mycorrhizas in plant phosphorus nutrition: interactions between pathways of phosphorus uptake in arbuscular mycorrhizal roots have important implications for understanding and manipulating plant phosphorus acquisition. *Plant Physiology*. 2011;156(3): 1050–1057. DOI: 10.1104/pp.111.174581
19. Thaines F, Braz ME, Mattos PP, Thaines APR. Equações para estimativa de volume de madeira para a região da bacia do Rio Ituxi, Lábrea, AM. *Pesquisa Florestal Brasileira*. 2010;30(64), 283-289. Portuguese. DOI: 10.4336/2010
20. Santos HS, Scapim CA, Maciel SL, Vida JB, Schwan-Estrada KRF, Brandão Filho JUT. Patogenicidade de *Meloidogyne javanica* em alface em função do tamanho de células de bandeja e idade de transplante das mudas. *Acta Scientiarum. Agronomy*. 2006;28(2), 253–259. Portuguese. DOI:10.4025/actasciagron.v28i2.1115
21. Empresa Brasileira De Pesquisa Agropecuária. Brazilian system of soil classification. Rio de Janeiro: Embrapa; 2013
22. Trajano MAT, Kasuya MCM, Tótola MR, Borges A, Novais RF. Suprimento de fósforo e formação de micorrizas em mudas de eucalipto em sistema de raízes divididas. *Rev. Árvore*. 2001; 25:193-2001. Portuguese.
23. Gerdemann JW, Nicolson TH. Spores of mycorrhizal *Endogone* extracted from soil by wet sieving and decanting. *Trans. Brit. Mycol. Soc.* 1963; 46(2): 235-244. DOI: [https://doi.org/10.1016/S0007-1536\(63\)80079-0](https://doi.org/10.1016/S0007-1536(63)80079-0)
24. Douds Junior DD, Nagahashi G, Pfeffer PE, Kayser WM, Reider C. On-farm production and utilization of arbuscular mycorrhizal fungus inoculum. *Canadian Journal of Plant Science*. 2005;85(1): 15-21. DOI: 10.4141/P03-168
25. Brito VN, Tellechea FRF, Heitor LC, Freitas MSM, Martins MA. Fungos micorrízicos arbusculares e adubação fosfatada na produção de mudas de Pariká. *Ciência Florestal*. 2017; 27(2): 485-497. Portuguese DOI: 10.5902/1980509827730
26. Dantas BF, Correia JS, Marinho LB, Aragão CA. Alterações bioquímicas durante a embebição de sementes de catingueira (*Caesalpinia pyramidalis* Tul.). *Revista Brasileira de Sementes*. 2008;30(1): 221-227. Portuguese. DOI:10.1590/S0101-31222008000100028.
27. Shimizu ESC, Pinheiro HA, Costa MA, Santos Filho BG. Aspectos fisiológicos da germinação e da qualidade de plântulas de *Schizolobium amazonicum* em resposta à escarificação das sementes em lixa e água quente. *Revista Árvore*. 2011;35(4): 791–800. Portuguese. DOI: 10.1590/S0100-67622011000500004
28. Cruz ED, Carvalho JEU, Leão NVM. Métodos para superação da dormência e biometria de frutos e sementes de *Parkia nitida* Miquel. (Leguminosae - Mimosoideae). *Acta Amazonica*.

2001;31(2): 167. Portuguese DOI:10.1590/1809-43922001312177.

29. Rocha CRM, Costa DS, Novembre ADLC, Cruz ED. Morfobiometria e germinação de sementes de *Parkia multijuga Benth.* Pesquisas Agrárias e Ambientais. 2014; 2(1): 42–47. Portuguese. DOI: 10.14583/2318-7670.v02n01a08

30. Camargo JLC, Ferraz IDK, Imakawa AM. Rehabilitation of degraded areas of Central Amazonia using direct sowing of forest tree seeds. Restoration Ecology. 2002;10(4): 636–644. DOI:10.1046/j.1526-100X.2002.01044.x

31. Meir D, Pivonia S, Levita R, Dori I, Ganot L, Meir S et al. Spanish Journal of Agricultural Research. 2010; 8 (S1): S5-S10. DOI: 10.5424/sjar/201008S1-1221

32. Trindade AV, Saggin Junior OJ, Silveira APD. Arbuscular mycorrhizae in the production of seedlings of fruit and coffee plants. In: Siqueira JO, Souza FA, Cardoso EJB, Siu M T. Mycorrhizae: 30 years of research in Brazil. Lavras: UFLA; 2010.

33. Cavalcante UMT, Maia LC, Nogueira RJMC, Santos VF. Respostas fisiológicas em mudas de maracujazeiro amarelo (*Passiflora edulis* Sims. f. *flavicarpa* Deg.) inoculadas com fungos micorrízicos arbusculares e submetidas a estresse hídrico. Acta Botanica Brasilica. 2001;15(3): 379-390. Portuguese. DOI:10.1590/S0102-33062001000300008.

34. Dickson A, Leaf A. L, Hosner J. F. Quality appraisal of white spruce and white pine seedling stock in nurseries. The Forestry Chronicle. 1960;36(1): 10-13. DOI: 10.5558/tfc36010-1

35. Vasconcellos CA, Alves VMC, Pereira Filho IA, Pitta GVE. Nutrition and fertilization of the maize in order to obtain baby corn. Sete Lagoas: Embrapa Milho e Sorgo; 2001

36. Oliveira MC, Ogata RS, Andrade GA, Santos DS, Souza RM, Guimaraes TG et al. Nursery manual and seedling production: tree species native to the Cerrado. Brasília: Universidade de Brasília; 2012

37. Campos MAA, Uchida T. Influência do sombreamento no crescimento de mudas de três espécies amazônicas. Pesquisa Agropecuária Brasileira. 2002;37(3):281-288. Portuguese. DOI:10.1590/S0100-204X2002000300008.

38. Rodrigues AF, Carvalho JG, Melo César P. Efeito do fósforo e do zinco sobre o crescimento de mudas do cupuaçuzeiro (*Theobroma grandiflorum Schum.*). Cerne. 2003;9(2), 221-230. Portuguese.

39. Gomes JM, Couto L, Leite HG, Xavier A, Garcia SLR, Parâmetros morfológicos na avaliação de qualidade de mudas de Eucalyptus grandis. Revista Árvore. 2002;26(6): 655-664. Portuguese. DOI:10.1590/S0100-67622002000600002

40. Azevedo IMG, Alencar RM, Barbosa AP, Almeida NO. Estudo do crescimento e qualidade de mudas de marupá (*Simarouba amara* Aubl.) em viveiro. Acta Amazonica. 2010;40(1): 157-164. Portuguese. DOI: 10.1590/S0044-59672010000100020