

Nutrient solution for production and quality of strawberry grown in substrate

ABSTRACT

Aims: This study determined a nutritive solution and evaluated the performance in the development, production and quality of strawberry cultivated in substrate.

Study design: The treatments were commercial and recommended nutritional solutions for strawberry using the methods of Castelane and Araújo (C.A.), Furlani and Fernandes Junior (F.F.J.) and the proposed solution with seven replicates.

Place and Duration of Study: The experiment was carried out in the experimental area of the Federal Technological University of Paraná, Brazil, in the period between May and December 2014.

Methodology: Agronomic variables such as yield, number of fruits, nutrient content, physiological indicators, physical and chemical characteristics of fruits were analyzed.

Results: The proposed nutrient solution resulted in larger masses of fresh and dry matter ($225.4 \text{ g plant}^{-1}$ and $27.5 \text{ g plant}^{-1}$), number of fruits (40.1) and fresh fruit mass (g plant^{-1}), in relation to the other evaluated solutions. The proposed solution resulted better physical and chemical characteristics such as soluble solids, reducing and total sugars, anthocyanins, flavonoids, phenolic compounds and ascorbic acid and the strawberry fruits presented an attractive color and met the quality standards for the consumer. The highest levels of nitrogen (33.7 g kg^{-1}), phosphorus (9.3 g kg^{-1}), and potassium (28.2 g kg^{-1}) in the leaf tissue were found in the proposed solution and contributed to productivity and fruit quality gains of a strawberry.

Conclusion: These results provide a nutrient base and can be adapted to other cultivars in different locations.

Keywords: *Fragaria x ananassa* Duch, nutrients, colour, physical and chemical characteristics

1. INTRODUCTION

The strawberry (*Fragaria x ananassa* Duch) is one of the fruits most appreciated by consumers in different regions of the world, highlighting its color, aroma, flavor and versatility in cooking and gastronomy. For this reason, strawberries are in great demand both in the natura and industrial processings [1].

Strawberry fruits with better physicochemical characteristics guarantee acceptance by the consumer market and increase yield in processing and industrialization. In this sense, the nutritional solution concentration, together with use of processing techniques have been important factors taken into account to improve the productivity and physicochemical properties of the fruit [2].

Strawberry cultivation in substrate is a production technique used in several regions of Brazil and around the world, allowing to obtain high production and greater ergonomics in crop management [3]. The main problem faced by producers in this production system is with regard to composition and management of the concentration of nutrient solution.

In the literature [4], reported that concentrations of nutrient solution with values of electrical conductivity (EC) between 1.4 and 1.8 dS m⁻¹ and up to 2.0 dS m⁻¹ [5] are proposed to obtain quality and productivity of strawberry fruits, but the great difficulty with most nutrient solutions is to adjust the amount of nutrients for substrate cultivation.

In this sense, the need arises for studies with nutritive solutions with determination of ionic balance of nutrients and their relationship with yield and quality of strawberry fruits in substrate cultivation. In this study we determined a nutrient solution for strawberry and evaluated the agronomic characteristics and fruit quality. The results provide nutrient content information extracted by the plants with the proposed solution, production data and physiological indicators of fruit quality, which contribute to meet the demands of the consumer market and make the production system more sustainable.

2. MATERIAL AND METHODS

2.1 Plant Material and Growing Conditions.

The experiment was carried out in the experimental area of the Federal Technological University of Paraná, Brazil (25°42'52 "S, 53°03'94" W, 530 m altitude), in the period between May and December 2014, covered with a 150-micron plastic film.

The seedlings of the cultivar Camino Real were purchased from a suitable nurseryman of varietal quality, from Maxxi Mudas®, from Patagonia, Argentina. These were transplanted in plastic pots with a capacity of 8 L in dimensions 24 × 23 cm, placed in lines, on the soil of the protected environment, filled with sand of medium granulometry, being transplanted one plant per pot, distributed with a density of eight plants per square meter.

The replenishment of nutrients was carried out daily by means of a drip irrigation system, with drippers of the brand netafim®, with a spacing of 0.20 m and a flow of 3.2 L hour⁻¹, with a dripper per vessel, thus maintaining the sand in the field capacity. The total fertigated volume was 535.7 mm and the total irrigation time was 47.5 hours for all treatments.

The meteorological data (temperature, relative air humidity and solar radiation) were obtained every 15 minutes using Akso® brand AK 172 dataloggers installed in meteorological shelters, located in the center of the protected environment.

The fertilizers used to compose the evaluated nutrient solutions were potassium nitrate (KNO₃), calcium nitrate Ca(NO₃)₂, monoammonium phosphate (NH₄H₂PO₄) and magnesium sulfate (MgSO₄). For micronutrients the amount of 25 g per 1000 L of water of the commercial product Conmicros Standard® was used in all the nutrient solutions, which presented the concentrations of B (2.0%), CuEDTA (2.0%), FeEDTA (7.9%), MnEDTA (2.0%), Mo (0.4%) and ZnEDTA (0.8%).

The nutrient solutions after addition of the nutrients presented the following values of electrical conductivity and pH 2.0 and 6.0 mS cm⁻¹ for the commercial solution (F.F.J.) [6],

1.7 and 6.2 mS cm⁻¹ for the commercial solution (C.A.) [7], and 1.8 and 5.8 mS cm⁻¹ for the proposed solution.

In relation to the management of nutrient solutions, the fertigations were done daily, and at each application of the fertirrigation, a new solution for each treatment was prepared. Also, twice-a-week irrigations were carried out only with water to avoid salinization of the substrate. Electrical conductivity and pH were measured with conductivity and portable HI 98130 Hanna® brand portable pH meters each time the solution was prepared. pH values between 5 and 6, and electrical conductivity greater than 1.5 mS cm⁻¹ were maintained during the experiment [8].

2.2. Treatments and Experimental Design

The treatments were commercial and recommended nutritional solutions for strawberry using the methods (C.A.) [7], (F.F.J.) [6] and proposed solution, with seven replicates. The amounts of nutrients used for each solution are shown in Table 1. The calculation of the proposed solution was based on ionic nutrient balance [9].

Table 1. Quantities of nutrients used in the preparation of nutrient solutions for strawberry cultivation on substrate.

Nutrients (mg L ⁻¹)	Commercial solution C.A.	Commercial solution F.F.J.	Proposed solution
N-NO ₃ ⁻	124.6	116.2	166.9
N-NH ₄ ⁺	5.6	5.6	31.92
P - H ₂ PO ₄ ⁻	46.5	49.6	78.12
K ⁺	195.0	234.0	182.13
Ca ⁺⁺	124.0	104.0	68.0
Mg ⁺⁺	24.60	36.0	13.7
S-SO ₄ ⁻	43.20	48.0	16.6

2.3. Evaluated Parameters

The content of macronutrients and micronutrients in leaf tissue was determined and four leaves per plant were completely expanded in the flowering period [10].

During the full flowering period (120 days after transplanting [DAT]) and at 190 DAT, measurements of the relative index of total chlorophyll in the abaxial and adaxial parts of the last two expanded leaves of each plant were performed at 11:00 AM using the chlorophyllometer model Clorofilog Falker® brand. The fresh matter mass of fruits and

106 number of fruits per plant was determined by adding all the harvests during the evaluated
107 period (fifteen harvests).

108 The average mass of fruits was obtained by dividing the fresh matter mass of fruits by the
109 number of fruits per plant. The fruits were harvested when they presented more than 75% of
110 the epidermis with pink coloration [11].

111 The fruit colour was determined in 10 fruits randomly selected from each nutrient solution,
112 using a digital colorimeter (Minolta model, Cr 200 b), where the values of luminosity ("L")
113 were determined, ranging from light to dark. The value 100 corresponds to white colour and
114 value 0 (zero), the black colour, and component "c" which expresses chroma degree of the
115 fruits, where, by the proposed classification, more colourful fruits present smaller values and
116 less colourful fruits present higher values [12].

117 The soluble solids (SS) content was obtained by direct reading in Hanna® bench
118 refractometer model HI 96801, using homogenized pulp and filtered at room temperature,
119 obtaining the values in degrees (Brix). The determination of the titratable acidity (T.A.) was
120 by titration with 0.1N NaOH until it reaches pH 8.1. The ratio (SS/TA) was determined by
121 dividing the soluble solids content by the titratable acidity.

122 Total sugar concentrations were determined by the method described by Dubois et al. [13]
123 those of reducing sugars were obtained by the method described by Miller [14].

124 The quantification of total phenolic compounds (mg gallic acid 100 g pulp⁻¹) was carried out
125 according to spectrophotometric method of Follin-Ciocauteau, proposed by Woisky and
126 Salatino [15]. The ascorbic acid content (Vitamin C) was determined by standard titration
127 method of AOAC modified by Benassi and Antunes [16]. Vitamin C content was calculated
128 based on titration values of standard solution of ascorbic acid and the results expressed in
129 mg of 10 g of ascorbic acid 100 g pulp⁻¹.

130 In the quantification of anthocyanins and flavonoids, the procedure described by Mazaro et
131 al. [17] was used. All the physicochemical analyses were determined in a single crop, at 150
132 DAT, which corresponded to the peak of production. A composite sample of 100 fruits per
133 treatment was used for all the analyses, taking seven subsamples of approximately 50 g
134 each.

135 At 190 DAT, which corresponded to end of the experiment, the mass of fresh matter in a
136 precision scale (0.001 g) of all the plants of experiment was determined. After the plants
137 were placed to dry in a forced circulation air oven at 65°C until reaching constant mass to
138 determine the mass of dry matter.

139 140 **2.4. Statistical Analysis**

141
142 The data of experiment were submitted for analysis of variance (Test F), when the F test was
143 significant the means were compared by Tukey's test ($P=0.05$), using "SAS Studio" [18].
144

145 146 **3. RESULTS AND DISCUSSION**

147
148 During the conduction of experiment, the average temperature, relative humidity and
149 average daily radiation was 19.2°C, 75% and 949.7 kJ m⁻². The temperature conditions
150 during the experiment were found to be within the ranges suitable for the crop.

151 Temperatures that range between 18°C and 24°C are considered adequate for development
152 of the crop [19].
153

154 The electrical conductivity (EC) in the solutions tested ranged from 1.5 to 2.1 dS m⁻¹, with an
155 average of 1.8 dS m⁻¹. The mean conductivity is at the upper limit of recommended range of
156 1.4 to 1.8 dS m⁻¹ [6]. The pH variations of the solutions were between 5.0 and 7.0, with an
157 average of 6.2. pH ranges between 5.5 and 6.5 are most indicated for the culture [6].
158

159 It was observed that the EC in the evaluated solutions improved fruit quality by increasing
160 the solids content and sugars. It was found that evaluated solutions were within the
161 recommended pH range for strawberry.
162

163 Nutrient solutions significantly influenced nutrient content in leaf tissue. The highest levels of
164 nitrogen (33.7 g kg⁻¹), phosphorus (9.3 g kg⁻¹) and potassium (28.2 g kg⁻¹) in the leaf tissue
165 were found in the proposed solution (Table 2). The other macronutrients did not differ
166 significantly. For micronutrients, there were significant differences for the boron content in
167 F.F.J. solution and higher iron and manganese contents in the proposed solution.
168

169 **Table 2. Nutrient content in leaf tissue of fertigated strawberry with different nutrient**
170 **solutions.**

Nutrients	N	P	K	Ca	Mg	S	B	Cu	Fe	Mn	Zn
	g kg ⁻¹	g kg ⁻¹	g kg ⁻¹	g kg ⁻¹	g kg ⁻¹	g kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹	mg kg ⁻¹
Sol. C.A.	31.1 b*	7.8 b	21.8 b	11.1 ^{ns}	6.9 ^{ns}	1.1 ^{ns}	100.5 b	5.3 ^{ns}	82.6 b	194.0 b	30.0 ^{ns}
Sol. F.F.J.	30.4 b	8.0 b	23.4 b	10.9	6.8	1.2	103.0 a	5.7	81.7 b	216.5 b	29.5
P. solution	33.7 a	9.3 a	28.2 a	10.5	7.0	1.2	98.4 b	5.5	86.5 a	222.0 a	31.6
Mean	31.7	8.4	24.5	10.8	6.9	1.2	100.6	5.5	83.6	210.8	30.4
C.V. (%)	3.9	9.2	3.3	3.1	5.6	4.3	1.2	2.3	3.5	2.6	2.2

171 *Means followed by the same letter in the column do not differ significantly by Tukey test, at

172 **P=0.05**; ns: **no significant**; C.V.: Coefficient of variance

173
174 The macronutrients, in descending order, nitrogen (N), potassium (K), calcium (Ca),
175 phosphorus (P), magnesium (Mg), and sulphur (S) were the nutrients extracted in greater
176 quantity by the strawberry. The following ranges are recommended: N, **15-25 g kg⁻¹**; P, 2-4 g
177 kg⁻¹; K, 20-40 g kg⁻¹; Ca, 10-25 g kg⁻¹; Mg, 6-10 g kg⁻¹; and S, 1-5 g kg⁻¹. For boron (B), iron
178 (Fe), manganese (Mn), copper (Cu) and zinc (Zn) are 35-100, 50-300, 30-300, 5-20, and 20-
179 50 mg kg⁻¹, respectively [20]. The contents found in the foliar tissue for the studied solutions
180 are superior to those suitable for N and P, within the recommended range for K, Ca, Mg, B,
181 Fe, Mn, Zn, and Cu. No visual symptoms of nutritional deficiency were observed in the
182 strawberry plants during the experiment.
183

184 The proposed solution resulted in highest relative indices of total chlorophyll in the flowering
 185 phase and at end of the crop cycle (Table 3). There was also a decrease in the relative
 186 index of total chlorophyll in final phase of the cycle in all evaluated solutions.

187
 188 The mass of the fresh and dry matter presented significant differences, being largest
 189 accumulation of fresh ($225.4 \text{ g plant}^{-1}$) and dry ($27.5 \text{ g plant}^{-1}$) mass obtained in the
 190 proposed solution (Table 3). The fresh mass of the proposed solution was 6.74% higher
 191 than the Castelane and Araújo commercial solution.

192
 193
 194 **Table 3. Relative index (I.R.) of total chlorophyll phases full flowering and the end,**
 195 **masses of fresh and dry matter of shoot (M.F. and M.S.) of fertigated strawberry**
 196 **plants with different nutrient solutions.**

Solutions	Full Flowering I. R. of total chlorophyll	End of cycle I. R. of total chlorophyll	M.F. (g planta^{-1})	M.S. (g planta^{-1})
Sol. C. A.	54.4 b*	50.2 b	210.2 b	21.0 b
Sol. F. F.J.	57.1 b	52.9 b	208.7 b	20.7 b
P. solution	62.2 a	59.4 a	225.4 a	27.5 a
Mean	57.9	54.17	215.0	23.1
C.V. (%)	9.3	8.4	20.4	22.0

197 *Means followed by the same letter in the column do not differ significantly by Tukey test, at
 198 $P=0.05$; C.V.: Coefficient of variance

199
 200 The highest relative chlorophyll index in the proposed solution is justified by higher nitrogen
 201 content present in the leaf tissue. The content of chlorophyll in the leaf is used to predict
 202 nutritional **level of N in plants**, due to the fact that the amount of this pigment correlates
 203 positively **with N** content in the plant [21]. This relationship is attributed mainly to the fact that
 204 50% to 70% of total N of the leaves is integral with enzymes, which are associated with
 205 chloroplasts [22].

206 The decrease of relative index of chlorophyll in the final phase of crop cycle can be
 207 explained by the advancing age of the leaf, because in this phase, there is a decline of
 208 photosynthetic capacity. The photosynthetic efficiency is linked to the amount of chlorophyll
 209 and consequently, to the growth phase of the plant [23].

210
 211
 212 The results of mass of the fresh and dry matter obtained with the proposed solution may be
 213 related to higher nutrient intake, especially the nitrogen present in the foliar tissue of the
 214 proposed solution (Table 2). Plant development, productivity and strawberry fruit quality are
 215 strongly influenced by nitrogen fertilization [24].

216 Moreover, the increase of K in the plant causes an increase in the production of
 217 photoassimilates and consequently, a greater mobilization of leaf N in the synthesis of
 218 macromolecules, which in turn are used in vegetative growth and fruit production [25].
 219

The number of fruits per plant, average fruit mass, and fresh fruit mass were influenced by evaluated treatments, obtaining best results in the proposed solution (Table 4). There were gains of 6.1% and 7.94% in the fresh fruit mass in relation to commercial solutions F.F.J. and C.A, which can be attributed to ionic balance of the proposed solution, which met the nutritional demand of strawberry with nutrient amounts without excesses or deficiencies, contributing to fruit quality and sustainable management of fertilizers in agriculture.

For the luminosity (L) of the epidermis, the nutrient solutions evaluated did not present significant influence (Table 4). The values of luminosity of the evaluated solutions were below the value 29.24 and according to [12] indicate dark colour. The dark colour of strawberry fruits in the evaluated solutions is a desirable characteristic for both industry and consumers, because dark red fruits are more attractive in the eyes of consumers.

For the colour component or chroma value of the epidermis, the proposed solution presented darker and more colourful fruits. The "C" component expresses the colour of fruits, where values less than 24.92 have more colour of the epidermis, values between 24.92 and 36.08 have intermediate colour, and values above 36.08 have less colourful fruits [12]. It is of great importance that external aspect of the fruit in commercialization is mainly in natura, the proposed solution resulted in fruits being more attractive for commercialization.

Table 4. Number of fruits plant⁻¹ (N.F.P.), mean fruit mass (M.F.M), fresh fruit mass (F.F.M.), luminosity of the epidermis, colour of the epidermis (Chroma) of fertigated strawberry with nutritive solutions.

Solution	N.F.P.	M.F.M. (g)	F.F.M. (g plant ⁻¹)	Luminosity	Chroma
Sol. C.A.	30.3 b*	12.5 b	690.8 b	28.7 ^{ns}	35.1 a
Sol. F.F.J.	32.5 b	13.3 b	704.6 b	27.8	34.93 a
P. solution	40.1 a	15.7 a	750.4 a	26.0	32.10 b
Mean	34.3	13.83	715.3	27.8	34.0
C.V. (%)	22.4	13.7	27.8	3.81	3.87

*Means followed by the same letter in the column do not differ significantly by Tukey test, at

P=0.05; ns: no significant; C.V.: Coefficient of variance

The superiority in the number of fruits and fresh fruit mass in proposed solution can be attributed to ionic balance of the solution, which favored the absorption of some ions, such as potassium (28.2 g kg⁻¹) (Table 2), which improved productivity and fruit quality [26]. The increase in mean mass of fruits, influenced by potassium present in the proposed solution, can be attributed to the important role that this nutrient plays in the translocation of photoassimilates from leaves to the fruits and the role it exerts in cell extension [25].

The number of fruits found in the proposed solution was 33.9% higher than that observed by Vignolo et al. [27], where 26.5 fruits per plant with the cultivar Camino Real in cultivation

were carried out in the soil. The fresh fruit mass transformed into yield results in 60 t ha⁻¹, which is higher than the yield obtained from 9.07 and 10.55 t ha⁻¹, with the same cultivar in conventional and organic systems, respectively [28].

The proposed solution resulted in more colourful fruits, with a higher content of soluble solids and of total and reducing sugars, possibly due to the higher content of potassium. Potassium is one of the nutrients most used by strawberry, considered the “element of quality” in plant nutrition, to improve physical-chemical characteristics and to increase production [10].

The SS/TA were not influenced by the evaluated nutrient solutions, the average value being 0.90 g.100 g pulp⁻¹ and 9.33, respectively (Table 5). It was verified that the proposed nutrient solution resulted in a higher content of soluble solids and concentration of total and reducing sugars, anthocyanins, flavonoids, phenolic compounds and ascorbic acid (Table 5).

266

Table 5. Soluble solids (S.S.) (°Brix), total sugars (T.S.) (mg.g⁻¹ fresh fruit mass), reducing sugars (R.S.) (mg.g⁻¹ fresh fruit mass), flavonoids (F.) (mg.100 g⁻¹ fresh fruit mass), anthocyanins (A.) (mg.100 g⁻¹ fresh fruit mass), phenolic compounds (P.C.) (mg of galic acid 100 g⁻¹ fresh fruit mass) and ascorbic acid (A.A.) (mg.100 g⁻¹ pulp) of fertigated strawberry fruits with different nutrient solutions.

Solution	S.S.	T.S.	R.S.	F.	A.	P.C.	A.A.
Sol. C. A.	8.0 b*	8.7 b	1.1 b	3.3 b	34.0 b	75.1 b	45.1 b
Sol. F. F. J.	8.2 b	9.0 b	1.3 b	3.1 b	35.0 b	76.4 b	47.0 b
P. solution	8.9 a	10.1 a	1.9 a	4.0 a	40.4 a	80.6 a	52.5 a
Mean	8.4	9.3	1.4	3.5	36.1	77.4	48.2
C.V. (%)	8.1	7.7	19.3	17.9	20.2	10.3	12.7

*Means followed by the same letter in the column do not differ significantly by Tukey test, at

P=0.05; C.V.: Coefficient of variance.

The soluble solids content in the proposed solution was 25.3% higher than the results obtained by [26] of 6.65 °Brix. The minimum values of soluble solids should be higher than 7.0 °Brix, guaranteeing acceptable taste [11], all nutritional solutions presented values above 7.0 °Brix, considered acceptable for consumers.

In the relationship between sugar content and acidity (SS/TA) there was no statistically significant difference between the evaluated solutions, with a mean value of 9.33. This value meets the minimum relationship patterns for strawberry fruits of 8.75 [11]. The strawberry fruits of the cultivar Camino Real presented an adequate SS/TA ratio, with degree of maturation and fruit quality. The SS/TA ratio is an important parameter to determine fruit

283 maturation, and fruit taste evaluation, as well as an indicator of fruit palatability, being
 284 directly linked to the preference and acceptance of the fruits by the consumer [29].

285 Commercially, the colour of the fruits can be influenced by the anthocyanins, which
 286 contributes greatly to quality evaluation, since the consumers correlate between the colour
 287 and total quality of specific products [30]. The anthocyanin content in the proposed solution
 288 was higher than that reported in the literature (20.93 mg 100 g⁻¹ fresh fruit mass) by [31] with
 289 the same cultivar, on different commercial substrates. According to [32] the anthocyanin
 290 levels may present variations related to climatic factors, seasonality, degree of maturation,
 291 nutrition and type of cultivar.

292 The results of present study indicate that fruits of the proposed solution presented an
 293 attractive colour and fruits with higher concentrations of anthocyanins, allowing greater
 294 benefits to the consumer due to the antioxidant effect.

295 As the anthocyanins content may be a criterion of choice at the time of feeding, due to the
 296 health benefits [31], the consumer will be eating higher anthocyanin content when
 297 consuming strawberries of the Camino real proposed solution. For humans, the intake of
 298 foods rich in anthocyanins, such as red fruits, is related to health benefits, as these
 299 components have high antioxidant and antitumor activity, as well as acting as an anti-
 300 inflammatory and preventing the formation of edemas [33].

301
 302 The phenolic compounds and ascorbic acid contents presented significant differences for
 303 the evaluated solutions. The proposed solution resulted in an increase in phenolic
 304 compounds and ascorbic acid. In the literature it is reported that potassium fertilization
 305 exerts a beneficial effect on vitamin C levels [34].

306 Furthermore, the phenolic contents found in this study are lower than those verified by [35]
 307 with the same cultivar (174.3 mg 100 g⁻¹ pulp). Phenolic compounds are significantly
 308 influenced by the genetic factors of the cultivar [36]. In addition, the "open" culture system
 309 provides a higher content of phenolic compounds than the protected environment system
 310 [37].

311 Another factor that possibly influenced the content of phenolic compounds was the
 312 temperature. It is known that the synthesis of phenolic substances is favored by the milder
 313 temperatures, especially the nocturnal ones and also the temperature variation from day to
 314 night, affects pigment deposition [38]. The average temperature of 19.2 °C favored the
 315 deposition of phenolic compounds, anthocyanins and flavonoids in fruits.

316 Potassium exerts influence on phenolic content, as it is related to photosynthesis and to
 317 biosynthesis of starch and proteins. With the increase of K doses in the plant, the production
 318 of photosynthates increases, which may increase the targeting of excess carbon fixed to the
 319 pathway of shikimic acid, which is the pathway for the formation of phenolic compounds,
 320 which may increase the concentration of phenolics in the plant [39].

321 The ascorbic acid (Vitamin C) in strawberry may vary according to the cultivar, stage of
 322 ripening and fertilization. It is one of the most important nutritional components in fruits and
 323 human food and its content can be used as an index of food quality [11]. In addition to
 324 mineral nutrition, the intensity of solar radiation (949.7 kJ m⁻²) associated with the time of
 325 year (summer) contributed to the increase in ascorbic acid content. The intensity and
 326 duration of fruit exposure to sunrays during growth influence the amount of ascorbic acid
 327 formed is synthesized from sugars supplied by photosynthesis, which increases with the
 328 highest incidence of radiation [38].

329
330 The phenolic compounds and ascorbic acid contents presented significant differences for
331 the evaluated solutions. The proposed solution resulted in an increase in phenolic
332 compounds and ascorbic acid. In the literature it is reported that potassium fertilization
333 exerts a beneficial effect on vitamin C levels [34].

334 Furthermore, the phenolic contents found in this study are lower than those verified by [35]
335 with the same cultivar (174.3 mg 100 g⁻¹ pulp). Phenolic compounds are significantly
336 influenced by the genetic factors of the cultivar [36]. In addition, the "open" culture system
337 provides a higher content of phenolic compounds than the protected environment system
338 [37].

339 Another factor that possibly influenced the content of phenolic compounds was the
340 temperature. It is known that the synthesis of phenolic substances is favored by the milder
341 temperatures, especially the nocturnal ones and also the temperature variation from day to
342 night, affects pigment deposition [38]. The average temperature of 19.2 °C favored the
343 deposition of phenolic compounds, anthocyanins and flavonoids in fruits.

344 Potassium exerts influence on phenolic content, as it is related to photosynthesis and to
345 biosynthesis of starch and proteins. With the increase of K doses in the plant, the production
346 of photosynthates increases, which may increase the targeting of excess carbon fixed to the
347 pathway of shikimic acid, which is the pathway for the formation of phenolic compounds,
348 which may increase the concentration of phenolics in the plant [39].

349 The ascorbic acid (Vitamin C) in strawberry may vary according to the cultivar, stage of
350 ripening and fertilization. It is one of the most important nutritional components in fruits and
351 human food and its content can be used as an index of food quality [11]. In addition to
352 mineral nutrition, the intensity of solar radiation (949.7 kJ m⁻²) associated with the time of
353 year (summer) contributed to the increase in ascorbic acid content. The intensity and
354 duration of fruit exposure to sunrays during growth influence the amount of ascorbic acid
355 formed is synthesized from sugars supplied by photosynthesis, which increases with the
356 highest incidence of radiation [38].

357

358

359 **4. CONCLUSION**

360

361 In this study, the proposed nutrient solution contributed to productivity gains, fruit quality and
362 comes as an option of adequate nutrient content for the strawberry, with ionic balance,
363 without excess nutrients. These results provide a nutrient base and can be adapted to other
364 cultivars in different locations.

365

366

367 **ACKNOWLEDGEMENTS**

368

369 The authors thank the company Maxxi mudas ® for the donation of the seedlings to carry
370 out the research.

371

372

373 **COMPETING INTERESTS**

374

375 Authors have declared that no competing interests exist.

376

377 **AUTHORS' CONTRIBUTIONS**

378

379 This work was carried out in collaboration between all authors. Authors DP and AS designed
380 the study and wrote the first draft of the manuscript. Authors ICZ and DP performed the
381 experiments. Authors ICZ; DP and FK participated in fieldwork and laboratory analysis.
382 Authors ICZ and FK managed the analyses of the study. Author AS managed the literature
383 searches. All authors read and approved the final manuscript.

384

385

386 **REFERENCES**

387

388 1. Antunes LEC, Peres NA. Strawberry production in Brazil and South America.
389 International Journal of Fruit Science. 2013;13:156-161. Portuguese.

390 2. Almeida MEM, Martin ZJ, Makiyama PAA. Strawberry Industrialization. Agricultural
391 Report. 1999;20:84-88. Portuguese.

392 3. Moraes CAG, Furlani PR. Cultivation of fruit vegetables in hydroponics in a
393 protected environment. Agricultural Report. 1999;20:105-113. Portuguese.

394 4. Paranjpe AV, Cantliffe DJ, Lamb EM, Stoffela PJ. Winter strawberry production in
395 greenhouses using soilless substrates: an alternative to methyl bromide soil
396 fumigation. Proceedings of the Florida State for Horticultural Science. 2003;116:98-
397 105.

398 5. Sarooshi RA, Cresswell GC. Effects of hydroponic solution composition, electrical
399 conductivity and plant spacing on yield and quality of strawberries. Aust J Exp Agric.
400 1994;34:529-535.

401 6. Furlani PR, Fernandes JF. Hydroponic strawberry cultivation in protected
402 environment. In: National Strawberry Symposium. Pelotas: Embrapa Clima
403 Temperado. 2004;p.102-115. Portuguese.

404 7. Castellane PD, Araujo JAC. Cultivation without soil: hydroponics. Jaboticabal:
405 FUNEP; 1994. Portuguese.

406 8. Gonçalves MA, Cocco C, Picolotto L, Vignolo GK, Antunes LEC. Short-Day
407 Strawberry Cultivars. Technical Bulletin. 2015;322:1-8. Portuguese.

408 9. Andriolo JL. Physiology of protected cultures. Santa Maria: Publisher UFSM; 1999.
409 Portuguese.

410 10. Malavolta E, Vitti GC, Oliveira SA. Evaluation of the nutritional status of plants:
411 principles and applications. 2nd ed. Piracicaba: Brazilian Association for Research
412 on Potassium and Phosphate; 1997. Portuguese.

413 11. Chitarra MIF, Chitarra AB. Post harvest of fruits and vegetables: physiology and
414 handling. Lavras: UFLA; 2005. Portuguese.

- 415 12. Conti JH, Minami K, Tavares FCA. Production and quality of strawberry fruits in trials
416 conducted in Atibaia and Piracicaba. *Hortic Bras.* 2002;20:10-17.
- 417 13. Dubois M, Gilles KA, Hamilton J.K, Rebers PA, Smith F. 1956. Colorimetric method
418 for determination of sugar and related substances. *Analitical Biochemistry.*
419 1956;28:350-356.
- 420 14. Miller GL. 1959. Use of dinitrosalicylic and reagent for determination of reducing
421 sugar. *Analitical Chemistry.*1959;31:426-428.
- 422 15. Woisky RG, Salatino A. Propolis analysis: some parameters and procedures for
423 chemical quality control. *J Apic Res.* 1998;37:99-105.
- 424 16. Benassi MT, Antunes AJ. A comparison of metaphosphoric and oxalic acids as
425 solutions of extracts for the determination of vitamin C in vegetables. *Archives of*
426 *Biology and Technology.* 1998;31:507-513.
- 427 17. Mazaro SM, Fogolari H, Wagner A, Citadin I, Santos I. Potential of extracts based
428 on *Calendula officinalis* L. on the induction of phytoalexin synthesis and the
429 fungistatic effect on *Botrytis cinerea*, *in vitro* substrates. *Rev Bras Pl.* 2013;15:208-
430 213.
- 431 18. Sas Institute. Sas Studio. Acessed 15 August 2014. Available:
432 http://www.sas.com/en_us/software/university-edition.html//.
- 433 19. Ronque VER. Strawberry culture: review and practice. Curitiba: EMATER; 1998.
434 206p. Portuguese.
- 435 20. Tedesco MJ, Gianello C, Anghinoni I, Bissani CA, Camargo FAO, Wiethölter S.
436 Fertilization and liming manual for the RS and SC states. Porto Alegre: Brazilian
437 Society of Soil Science; 2004. Portuguese.
- 438 21. Smeal D, Zhang H. Chlorophyll meter evaluation for nitrogen management in corn.
439 *Commun Soil Sci Plant Anal.* 1994;25:1495-1503.
- 440 22. Stoking CR, Ongun A. The intracellular distribution of some metallic elements in
441 leaves. *Am J Bot.* 1962;49:284-289.
- 442 23. Conceição HEO, Oliva MA, Lopes NF, Rocha OGN. Resistance to drought in rubber
443 trees. *Pesqui Agropecu Bras.* 1985;20:1041-1050.
- 444 24. Otto RF, Morakami RK, Reghin MY, Caires EF. Cultivars of strawberry of neutral
445 day: production as a function of doses of nitrogen during the summer. *Hortic Bras.*
446 2009;27:217-221.
- 447 25. Anttonen MJ, Hoppuls KI, Nestby MJ, Karjalainen RO. Influence of fertilization,
448 mulch color, early forcing, fruit order, planting date, shading, growing environment,
449 and genotype on the contents of selected phenolics in strawberry (*Fragaria x*
450 *ananassa* Duch.) fruits. *J Agric Food Chem.* 2006;54:2614-2620.
- 451 26. Andriolo JL, Jänisch DI, Schmitt OJ, Dal Picio M, Cardoso FL, Erpen L. Potassium
452 and calcium doses in plant growth, yield and quality of strawberry fruits in soilless
453 cultivation. *Cienc Rural.* 2010;40:267-272.

- 454 27. Vignolo GK, Araújo VF, Kunde RJ, Silveira CAP, Antunes LEC. Production of
455 strawberries from alternative fertilizers in pre-planting. *Cienc Rural*. 2011;41:1755-
456 1761.
- 457 28. Camargo LKP, Resende JTV, Galvão AG, Camargo CK, Baier JE. Productive
458 performance and average mass of strawberry fruits obtained from different cropping
459 systems. *Ambience*. 2010;6:281-288. Portuguese.
- 460 29. Cecatto AP, Calvete EO, Nienow AA, Costa RC, Mendonça HFC, Pazzinato AC.
461 Culture systems in the production and quality of strawberry cultivars. *Acta Sci*.
462 2013;35:471- 478.
- 463 30. Kays SJ. Preharvest factors affecting appearance. *Postharvest Biol Technol*.
464 1999;15:233-247.
- 465 31. Calvete EO, Nienow AA, Wesp CL, Cestonaro L, Mariani F, Fioreze I, Cecchetti D,
466 Castilhos T. Hydroponic production of strawberry in vertical column system under
467 protected cultivation. *Rev Bras Frutic*. 2007;29:524-529. Portuguese.
- 468 32. Clifford MN. Anthocyanins—nature, occurrence and dietary burden.
469 *J Sci Food Agric*. 2000;80:1063-1072.
- 470 33. Aaby K, Wrolstad, RE, Ekeberg D, Skrede G. Polyphenol composition and
471 antioxidant activity in strawberry purees: impact of achenes level and storage. *J*
472 *Agric Food Chem*. 2007;55:5156-5166.
- 473 34. Lee SK, Kader AA. Preharvest and postharvest factors influencing vitamin C content
474 of horticultural crops. *Postharvest Biol Technol*. 2000;20:207-220.
- 475 35. Pineli LLO, Moretti CL, Santos MS, Campos AB, Brasileiro AV, Cordova AC,
476 Chiarello MD. Antioxidants and other chemical and physical characteristics of two
477 strawberry cultivars at different ripeness stages. *J Food Compost Anal*. 2011;24:
478 11–16.
- 479 36. Atkinson CJ, Dodds PAA, Ford YY, Le Mièrre J, Taylor JM, Blake PS, Paul N.
480 Effects of cultivar, fruit number and reflected photosynthetically active radiation on
481 *Fragaria x ananassa* productivity and fruit ellagic acid and ascorbic acid
482 concentrations. *Ann Bot*. 2006;97:429-441.
- 483 37. Hernanz D, Recamales AF, Meléndez-Martínez AJ, Gonzalez-Miret ML, Heredia FJ.
484 Assessment of the differences in the phenolic composition of five strawberry
485 cultivars (*Fragaria x ananassa* Duch.) grown in two different soilless systems. *J*
486 *Agric Food Chem*. 2007;55:1846-1852.
- 487 38. Abu-Zahra TR, Al-Ismaïl K, Shatat F. Effect of organic and conventional systems on
488 fruit quality of strawberry (*Fragaria x ananassa* Duch) grown under plastic house
489 conditions in the Jordan Valley. *Acta Hortic*. 2007;741:159-171.
- 490 39. Crozier A, Jaganath IB, Clifford MN. Phenols, polyphenols and tannins: An overview.
491 In: Crozier A, Clifford MN, Ashihara H. (eds.). *Plant secondary metabolites:*
492 *occurrence, structure and role in the human diet*. Oxford: Blackwell Publishing.
493 2006;p.1-24.
494