

Agronomic efficiency of organic fertilized in the production of the intercropping of coriander and mint in the northeastern Brazil

ABSTRACT

Oleric plants are grown in a system intercropping by family farmers in organic production areas in the region northeastern Brazil. The experiment was conducted in an experimental area belonging to a semi-arid rural federal university. The study was carried out in the period from August 2014 to November 2014, in order to evaluate agronomic efficiency of organic fertilized in the production of the intercropping of coriander and mint in the northeastern Brazil. The study area was designed in a randomized block area, arranged in a 2 x 4 factorial scheme set up, with three replications. The first factor set up consisted of single crop and intercropping of coriander and mint, and the second was hairy woodrose (*Merremia aegyptia* L.) added to cattle manure in the amounts (0.0, 2.0, 4.0 and 6.0 kg/2.0 m²). The evaluated characteristics for the coriander crop these were plant height, number of stems, green mass, number of bunches and dry mass. The evaluated characteristics for the mint crop these were biomass height, green mass, number of bunches, dry mass, oil content and yield oil. The agronomic efficiency of mint and coriander in intercropping was obtained with the addition of 6.0 kg/2.0 m² of organic fertilized incorporated into the soil. The equivalent area ratio was agronomically efficient, with equivalent area ratio (EAR) of 1.66.

Keywords: Medicinal plant; spice; agroecological production; family farmers.

1. INTRODUCTION

Coriander (*Coriandrum sativum*, Apiaceae) originates from the territories of the Mediterranean Basin. In Brazil, this herb was introduced at the beginning of colonization [1], produced by family farmers, the most produced vegetable in the region of Mossoró-RN [2, 3]. Another species widely cultivated and marketed in the region is mint (*Mentha piperita*), used in juices and salads, and is highly appreciated by consumers.

These species have the potential to be produced in a consortium, since they have different phenological cycles, coriander, 30 to 35 days after sowing and mint, 60 days after transplant, causing competition for water and nutrients at different times in the cultivation, guaranteeing agronomic performance.

The crop consortium is an important component of sustainable farming systems, which include organic farming. It is an agricultural practice in small production units in tropical regions, especially family-based ones [4]. This system consists in the simultaneous cultivation of two or more species in the same space within the phenological cycle of the

crops, being able to be sown simultaneously or different sowing times, guaranteeing a higher income to the farmers [5].

Within this context, the use of plant material of species available in production areas is of great importance to family farmers. Among the species with potential for phytomass production, jitirana, a spontaneous species from the northeastern semi-arid region of predominance in the rainy season, stands out for dry matter production of 4000 kg/ha with nitrogen concentration of 22.0 g/kg. It can be incorporated into the soil or left in the cover, becomes a viable option to be used as an organic fertilizer [6].

The mixture of organic fertilizers of vegetal and animal origin potentiates the production making the system more efficient, reducing the costs in the acquisition of agricultural inputs. The use of alternative sources as organic fertilizer in the production of vegetables has been studied by several researchers, beet [7,8]; carrot [9]; coriander [2] and lettuce [10].

Due to the importance of using the combination of fertilizer sources for the nutritional supply of the cultivation system, the objective was to evaluate agronomic efficiency of organic fertilized in the production of the intercropping of coriander and mint in the northeastern Brazil.

2. MATERIALS AND METHODS

2.1 Characterization of the experimental area

The experiment was conducted in the research area of the Rafael Fernandes Experimental Farm of the Federal Rural Semi-Arid University (UFERSA), in the period from August 2014 to November 2014, in soil of the experimental area was classified as Eutrophic Red Yellow Argissolo, caatinga hyperxerophilic phase and flat relief [11,12]. The area is located in the Alagoinha district, 20 km from the Mossoró, Northeastern Brazil. The farm comprises of some 400 hectares [11].

According to the classification of Köppen, the local climate is BSw_h ', dry and very hot, the dry season being normally from June to January, and a rainy season being from February to May. The average annual rainfall is 673.9 mm and the average relative humidity is 68.9% [13].

Before the installation of the field experiment, soil samples were collected to a 0-20 cm layer and then sent to be processed and analyzed in the UFERSA Water, Soil and Plant

Analysis Laboratory, providing the following results: pH (water 1:2,5) = 6.65; exchangeable cations Ca = 0.60 cmol_c/dm³; Mg = 0.75 cmol_c/dm³; K = 40.2 mg/dm³; Na = 9.5 mg/dm³; P (Mehlich) = 8.6 mg/dm³; organic matter = 1.20 g/kg; Coarse sand = 610 g/kg; fine sand = 290 g/kg; silte = 20 g/kg; clay = 70 g/kg, soil density = 1.42 g/cm³.

2.2 Experimental design and treatments

The study area was designed in a randomized block area, arranged in a 2 x 4 factorial scheme set up, with three replications. The first factor set up consisted of single crop and intercropping of coriander and mint, and the second was hairy woodrose (*Merremia aegyptia* L.) added to cattle manure in the amounts (0.0, 2.0, 4.0 and 6.0 kg/2.0 m²). The organic fertilizer remained incubated in the soil for a period of 30 days before planting [2].

The preparation of the soil consisted in the removal of the vegetal material and in the survey of the beds using rotocanteirador. Before sowing the coriander and transplanting the mint, the soil was kept moist, contributing to the mineralization process [14].

For the coriander crop, the plot was 1.4 m x 1.4 m, with a total area of 1.96 m², containing 1560 coriander plants with 0.1 m x 0.05 m spacing with five plants per pit [3]. In intercropping, the coriander was placed between the mint seeding rows. The coriander crop variety used was Verdão cultivar.

The cultivar "*Mentha piperita*" was used for the mint crop. The study plots were 1.4 m x 1.4 m, with a total area of 1.96/m², with 40 plants in each, spaced 0.35 m x 0.2 m respectively. The yielding area, was therefore 1.26/m², with 18 plants.

2.3 Chemical composition of organic fertilizers

The hairy woodrose (*Merremia aegyptia* L.) was collected in june of 2014, in an area of semi-arid shrub vegetation in the flowering period. Soon after being harvested the material was crushed and dried in the sun, being packed in raffia bags to be used as an organic fertilizer. The material was sent to the soil fertility and nutrition laboratory at the Federal University of the Semi-arid Agricultural Sciences Center, with a chemical concentration of 22.5 g/kg of nitrogen (N); 890 g/kg of organic matter (OM); 412 g/kg of organic carbon (CO); 18/1 carbon/nitrogen ratio (C/N); 0.75 g/kg of phosphorus (P); 4.87 g/kg potassium (K⁺); 1.20 g/kg sodium (Na⁺); 0.32 g/kg of calcium (Ca²⁺) and 0.08 g/kg of magnesium (Mg²⁺).

The cattle manure used was stabilized and sourced from the heifers of the UFERSA cattle herd, which are raised in an intensive system, fed with corn-based concentrate, soy bean and wheat bran, and having as bulk Canarian grass (*Echinocloa polystochya*), with chemical concentration of: 16.5 g/kg of nitrogen (N), 890 g/kg organic matter (OM); 401 g/kg of organic carbon (OC); 25/1 of carbon/nitrogen ratio (C/N) 0.68 g/kg of phosphorus (P); 5.87 g/kg of potassium (K⁺); 0.70 g/kg of sodium (Na⁺); 0.23 g/kg of calcium (Ca²⁺) and 0.05 g/kg of magnesium (Mg²⁺), carried out in the soil fertility and plant nutrition laboratory of the agrarian science center of the Federal Rural University of the semi-arid.

2.4 Harvest and evaluation of the intercropping

The harvest was carried for culture of coriander, at thirty-five days after sowing and for mint crop to sixty days after transplanting. After harvesting, plants were transported to the Post-Harvest of Vegetables Laboratory of the Agrarian Science Center of the Federal Rural University of the semi-arid.

For the coriander crop, the following characteristics were evaluated: For the mint crop, the following characteristics were evaluated: *plant height* (was measured in the field, in centimeters using millimeter ruler, twenty plants per plot), *green mass production* (It was obtained through the cut of the aerial part, within the useful area, weighed in a precision scale of 1.0 g, expressed in kg/100 m²); *number of bunches* (determined by the division of production in kg/100 m² per 50g, corresponding to the weight of the sauce in the supermarket of Mossoró, Northeastern Brazil); *dry mass production* (was obtained from a forced-air heating oven at 65 °C, to constant mass and expressed in kg/100 m²).

For the mint crop, the following characteristics were evaluated: *biomass height* (was measured in the field, in centimeters using millimeter ruler, twenty plants per plot), *green mass production* (It was obtained through the cut of the aerial part, above 5 cm, within the useful area, weighed in a precision scale of 1.0 g, expressed in kg/m²); *number of bunches* (determined by the division of production in kg/m² per 100g, corresponding to the weight of the sauce in the supermarket of Mossoró, Northeastern Brazil); *dry mass production* (was obtained from a forced-air heating oven at 65 °C, to constant mass and expressed in kg/m²), *Oil content* (It was defined by the ratio between the mass in grams of essential oil by the mass of dry leaves inserted into the distillation flask x 1000, being expressed in g/kg) and *Oil production* (It was determined by the oil content in g/kg x the production of dry mass in kg/m², being expressed in g/m²).

For the determination of oil content and yield, dry leaves were used, that the water content in the leaves, after drying, allows the vapor stream generated in the extractor to remove the volatile substances stored in cells, when compared to the green material [15]. Due to the high moisture content in the plants, there is a strong tendency to agglutinate preventing the steam from penetrating more easily of form uniform in plant tissues [16].

In determining the essential oil content and yield, dry mass samples were subjected to hydrodistillation in a modified Clevenger apparatus for 1.5 h, using 600 mL of distilled water in 1 L distillation flask [17].

It was also determined the efficiency of the intercropping through the ETS, defined as the relative area of land under isolated planting conditions required to provide the productivities reached in the consortium [18]. Defined by the following expression: $ETS = (Y_{ccm}/Y_{sc}) + (Y_{mcc}/Y_{msc})$, where the following is true: Y_{ccm} = productivity of coriander consorted with mint; Y_{sc} = productivity of coriander in single crop; Y_{mcc} = productivity of mint consorted with coriander; Y_{msc} = productivity of mint in single crop.

2.5 Statistical Analysis

An analysis of variance was performed on each feature of mint, through the application ESTAT [19]. Those of a quantitative nature (mixture of hairy woodrose with cattle manure), the procedure for adjusting the response curve was performed using the Table Curve [20] application. For the qualitative characteristics (single cropping and intercropping of coriander and mint), the Tukey test ($p < 0.05$).

3.0 RESULTS AND DISCUSSION

3.1 Cultivation of mint

Significant interaction was observed for characteristic yield of oil (Table 1). However, there was an isolated effect at the level of probability ($p < 0.01$) in the different amounts of organic fertilizer for all characteristics studied (Table 1). In relation to the cultivation systems (single crop, and intercropping), there was a statistical difference in the level of probability ($p < 0.01$), for the characteristics of green mass, number of sauces, dry mass, percentage and yield of oil (Table 1), probably due to interspecific competition for water, nutrients and physical space, demonstrating the exploitation capacity of the cultivated area.

Table 1. F values for biomass height, expressed in cm (BH), green mass, expressed in kg/2.0 m² (GM), number of bunches, expressed in units/2.0 m² (NB), dry mass, expressed as g/2.0 m² (DM), percentage of oil, expressed as% (PO) and oil yield, expressed in g/2.0 m² (OY) of mint as a function of organic fertilization

Causes of variation	GL	BH	MV	NM	MS	PO	RE
Amounts of fertilizers (A)	3	9.33 ^{**}	12.82 ^{**}	12.59 ^{**}	9.19 ^{**}	2.71 ^{ns}	6.86 [*]
Cultivation system (B)	1	8.17 ^{**}	13.13 ^{**}	12.90 ^{**}	8.07 ^{**}	1.51 ^{ns}	7.55 [*]
A X B	3	1.86 ^{ns}	2.00 ^{ns}	1.66 ^{ns}	1.50 ^{ns}	0.93 ^{ns}	0.90 ^{ns}
Treatments	7	----	----	----	----	----	----
Blocks	2	4.64 [*]	5.29 [*]	4.10 [*]	6.65 [*]	0.78 ^{ns}	5.80 [*]
Residue	14	----	----	----	----	----	----
CV (%)	----	13.41	11.70	12.3	10.70	8.88	6.95

^{**} = P < 0.01; ^{*} = P < 0.05; ^{ns} = not significant

There was an increasing effect due to the different amounts of organic fertilizer at plant height, with a value of 29 cm, in the amount of 6.0 kg/2.0 m² (Figure 1). In relation to the cultivation systems (single crop and intercropping), there was a statistical difference, with a mean value of 32.3 and 28.2 cm, respectively (Table 2). Behavior lower, with values of 24.7 and 23.7 cm in single crop and intercropped cultures was observed by [21], in the cultivation of mint. [22] evaluating the production of mint under doses of jitirana and harvesting times, with biomass height of 34.6 cm at 90 days after transplanting, value exceeding this research.




Figure 1. Mint biomass height as a function of amounts of organic fertilizer

For green mass and number of bunches, the dose of 6.0 kg/2.0 m² of organic fertilizer, contributed the largest increase, with an average of 3.94 kg/2.0 m² and 39.4 units de bunches/2.0 m², respectively (Figures 2A and 2B). The single crop was statistically higher to the intercrop, with mean values of 4.1 and 3.2 kg/2.0 m² for green mass, corresponding to 41.0 and 32.0 mint sauces, respectively (Table 2).

The number of mint sauces is of paramount importance to family farmers because it represents the form of commercialization in supermarkets and agro-ecological fairs. Completed searches found green mass of 322.5 g/m² in the cultivation of peppermint intercropped with lettuce, being less than said research [23].




Figure 2. Green mass (A) and number of bunches (B) of mint as a function of amounts of organic fertilizer

Table 2. Evaluation of the characteristics biomass height, expressed in cm plant (BH), green mass, expressed in kg/2.0 m² (GM) and number of bunches, expressed in units/2.0 m² (NB) of mint as a function of organic fertilization

cultivation system	BH	GM	NB
Single	32.3 a	4.1 a	41.0 a
Intercropping	28.2 a	3.2 b	32.0 b

*Means followed by similar lowercase letters in the column differ statistically by Tukey test at 5% probability

There was an increase of 270.6 g/2.0 m², in the dry mass between the amounts of 0 kg/2.0 m² and 6.0 kg/2.0 m², adjusting a linear equation of 1 degree, with a maximum value of 396.1 g/2.0 m² (Figura 3). The single crop was statistically superior to the intercropping with values of 412.5 and 374.2 g/2.0 m² (Table 3). The dry mass is the best way to evaluate plant growth, considering that this characteristic is not influenced by soil moisture [24].

No statistical difference was observed for the oil content, with average value of 2.1 g/kg. Similar behavior was observed in the cultivation system (single crop and intercropped) with mean values of 2.1 and 2.0, respectively (Table 3). According to [25], the essential oil content constitutes a genetic characteristic, not changing as a function of the amount of biomass produced.




Figure 3. Dry mass of mint as a function of amounts of organic fertilizer.

Unfolding the amounts of organic fertilizer within the cropping system (single crop and intercropping), with average values of 8.69 and 4.58 g/2.0 m² (Figure 4). For the crop system (single crop and intercropping) there were unfolding within of the amounts, being the single crop superior statistically to the intercropping in the amounts of 0.0; 2.0; 4.0 and 6.0 kg/2.0 m² (Table 3). The increase in the yield of essential oil, possibly occurred to the increase in the production of green mass and dry mass of the plants, as a function of the amounts of organic fertilized incorporated to the soil. Results lower was observed by [26], with a value equivalente of 4.0 g/2.0 m². [22] evaluating the production of mint under doses of jitirana and harvesting times, with oil yield of 3.7 g/2.0 m² at 90 days after transplanting, lower than said research.




Figure 4. Oil yield of mint as a function of amounts of organic fertilizer

Table 3. Unfolding of the cropping system (single crop and intercropping) within of the amounts of organic fertilized.

Cropping system	Amounts of organic fertilized (kg/2.0 m ²)			
	0.0	2.0	4.0	6.0
Single crop	1.6 a	4.52 a	8.20 a	8.69 a
Intercropping	0.78 b	3.15 b	4.70 b	4.75 b

*Means followed by different lowercase letters in the column differ statistically by Tukey test at 5% probability

3.2 Cultivation of coriander

There was significant interaction for green mass, number of sauces and dry mass of coriander. For the factor amounts fertilizer organic and in the cultivation system (single crop and intercropping), a statistical difference was observed at the level of $p < 0.01$ (Table 4).

Table 4. F values for plant height, expressed in cm plant (HP), number of stems, expressed in terms of average (NS), green mass of coriander, expressed in kg/2.0 m² (GM), number of bunches, expressed in units/2.0 m² (NB) and dry mass, expressed in g/2.0 m² (DM) of coriander as a function of organic fertilization

Causes of variation	GL	HP	NS	GM	NB	DM
Amounts of fertilizers (A)	3	28.42 ^{**}	15.60 ^{**}	17.22 ^{**}	16.59 ^{**}	12.46 ^{**}
Cultivation system (B)	1	0.12 ^{ns}	1.10 ^{ns}	9.90 ^{**}	8.30 ^{**}	18.24 ^{**}
A X B	3	0.35 ^{ns}	0.54 ^{ns}	2.66 [*]	5.16 [*]	7.34 ^{**}
Treatments	7					
Blocks	2	5.6 [*]	6.8 [*]	4.10 [*]	5.08 [*]	4.70 [*]
Residue	14	----	----	----	----	----
CV (%)	----	20.4	16.9	17.4	17.9	13.8

** = $P < 0.01$; * = $P < 0.05$; ^{ns} = não significativo

For plant height, it was observed a point of maximum height, with value of 19.2 cm/plant in the amount of 6.0 kg/2.0 m² of organic fertilizer (Figure 5). In the cultivation system (single crop and intercropping) were statistically similar, with mean values of 19.0 and 18.6 cm/plant, respectively (Table 5), lower values were observed by [27] and [3], with mean values of 20.5 and 22.0 cm/plant of coriander, respectively. Height of plant is an important characteristic in the commercialization of coriander, being the standard of commercialization above 15 cm.




Figure 5. Height of coriander plant as a function of organic fertilization

The amount of organic fertilizer of 4.0 kg/2.0 m² contributed to a number of stems of 6.5 stems/plant (Figure 6). In the cultivation system (single crop and intercropping), there was similarity between the observed data with values of 6.3 and 5.8 stems/plant, respectively (Table 5). Behavior superior to that found in the research was observed by [27] in coriander and [2] in the coriander culture, with values of 6.0 and 8.5 stems/plant, respectively.




Figure 6. Stems/plant of coriander as a function of organic fertilization

Table 5. Evaluation of the characteristics plant height, expressed in cm plant (HP), number of stems, expressed in terms of average (NS) of coriander as a function of organic fertilization

cultivation system	HP	NS
Single	19.0 a	6.3 a
Intercropping	18.6 a	5.8 a

*Means followed by similar lowercase letters in the column differ statistically by Tukey test at 5% probability

When unfolding the interaction in the organic fertilizer quantities within the growing system (single crop and intercropping) was observed values of 1200 and 790 g/m² for green

mass (Figure 6A and 6B) and 500 and 350 units /2.0 m² for number of sauces of coriander (Figure 7A and 7B). When unfolding the cultivation system (single crop and intercropping) in the different amounts of organic fertilizer, observed a statistically significant difference for the green mass (Table 5) and number of sauces (Table 6).

The inferiority of the single crop system in relation to the consortium is probably due to the competition for water, light and nutrients, in addition, the number of plants was inferior to the consortium. The number of sauces is the form of commercialization of the coriander in the gondolas of supermarkets and the agroecological fair, being of great importance for the family farmers in the region of Mossoró, RN, Brazil.

[28], studying evaluating optimized amount of hairy woodrose (*Merremia aegyptia* L.) in the productivity of coriander cultivars, verifying yield of 98.6 kg/100 m², equivalent to 1972 g/2.0 m² and 39.4 coriander sauces, being higher than said research. [29], studying the roostertree (*Calotropis procera*) under different amounts and periods of incorporation on yield of coriander with an average yield of 4404 kg/ha, equivalent to 880 g/2.0 m², corresponding to 17,6 bunches, being lower the present research.




Figure 6. Green mass of coriander as a function of single crop (A) and of cultivating intercropping (B) under amounts of organic fertilizer

Table 6. Unfolding the interaction cultivation system (single crop and intercropping) in the different amounts of organic fertilizer in the green mass of coriander, expressed in kg/2.0 m²

cultivation system	Amounts of organic fertilizer (kg/2.0 m ²)			
	0	2.0	4.0	6.0
Single	150 a	610 a	1300 a	1200 a
Intercropping	90 b	390 b	390 b	830 b

*Means followed by different lowercase letters in the column differ statistically by Tukey test at 5% probability




Figure 7. Bunches of coriander as a function of single crop (A) and of cultivating intercropping (B) under amounts of organic fertilizer

Table 6. Unfolding the interaction cultivation system (single crop and intercropping) in the different amounts of organic fertilizer in the green mass of coriander, expressed in kg/2.0 m²

cultivation system	Amounts of organic fertilizer (kg/2.0 m ²)			
	0	2.0	4.0	6.0
Single	4.0 a	13.0 a	26.0 a	23.0 a
Intercropping	3.0 b	8.0 b	13.0 b	16.0 b

*Means followed by different lowercase letters in the column differ statistically by Tukey test at 5% probability

Similar behavior was observed for the dry mass of coriander, with values of 146.7 and 79.9 g/2.0 m², within the cultivation system (single crop and intercropping), respectively, in the amount of 4.0 kg/2.0 m² of organic fertilizer (Figure 8A & 8B). In relation to the cultivation system (single crop and intercropping) within the amounts of organic fertilizer, the single crop was statistically superior to the intercropping, with values of 150 and 88 g/m², respectively. The dry mass is a characteristic that accurately measures plant growth, since it is not influenced by soil moisture conditions. Lower behavior was obtained by [2,29,30] evaluating spontaneous species in substitution of chemical fertilization, with values below that research.




Figure 8. Bunches of coriander as a function of single crop (A) and of cultivating intercropping (B) under amounts of organic fertilizer

Agronomically, two crops are suitable to be intercropping, if the production has an equivalent area ratio greater than 1, indicating superiority to the single crop.

In this context, the consortium of condiment vegetables presented superiority to the single crop, with an equivalent area ratio of 1.66, in the amount of 4.0 kg/2.0 m² of organic fertilizer (Figure 9). A single crop area of 66% would be required to have a similar production to the intercropping. For family farmers in the region Mossoro, RN, Northeast Brazil, who cultivate small areas, the use of two or more cultures in the same physical space, contributes to the best use of the area in the environmental and economic aspect.




Figure 9. Equivalent area ratio under amounts of organic fertilizer

4. Conclusion

The agronomic efficiency of mint and coriander in intercropping was obtained with the addition of 6.0 kg/2.0 m² of organic fertilized incorporated into the soil. The equivalent area ratio was agronomically efficient, with equivalent area ratio (EAR) of 1.66.

5. Acknowledgements

Special thank the Jitirana (*Merremia aegyptia* L.) Research Group committed to the study of spontaneous species of the Caatinga Biome as fertilizer source and the Department of Agronomic and Forestry Sciences the Federal Rural Semi-Arid University (UFERSA), for the support in the conduction of the project.

References

1. Melo RA, Menezes D, Resende LV, Wanderley Junior LJG, Santos VF, Mesquita JCP, Magalhães AG. Genetic variability in progenies of coriander half-brothers. *Horticultura Brasileira*. 2009; 27 (3): 324-329. Portuguese.
Available:<http://dx.doi.org/10.1590/S0102-05362009000300011>
2. Linhares PCF, Pereira MFS, Assis JP, Bezerra AKH. Amounts and times of decomposition of scarlet starglory on agronomic performance of cilantro. *Ciênc Rural*. 2012; 42 (2): 243-48. Portuguese. Available:<http://dx.doi.org/10.1590/S010384782012000200010>
3. Linhares PCF, Oliveira JD, Pereira MFS, Fernandes JPP, Dantas RP. Spacing for the coriander culture fertilized with carnauba straw in Mossoró-RN conditions. *Revista Verde de Agroecologia e Desenvolvimento Sustentável*, 2014. 9 (3): 01–06. Portuguese.
Available:<http://www.gvaa.com.br/revista/index.php/RVADS/article/view/2781>
4. Montezano EM, Peil RMN. Consortium system for the production of vegetables. *Revista Brasileira de Agrociência*. 2006; 12 (1): 129-132. Portuguese.
Available:<http://dx.doi.org/10.18539/cast.v12i2.4502>
5. Carrilho AJ. Production and bromatological analysis of cabbage and radish in consortium. Brasília 2013. 29 páginas. Monografia. Monografia de Graduação. Universidade de Brasília/Faculdade de Agronomia e Medicina Veterinária. Portuguese.
Available:http://www.bdm.unb.br/bitstream/10483/7564/6/2013_AngelinaJunqueiraCarrilho.pdf
6. Linhares PCF. Green manure as soil conditioner. *Revista Campo e negócios*, 2013; 11 (127): 22-23. Portuguese.
Available:<http://www.adubarofuturo.com.br/adubacao-verde-como-condicionadora-do-solo/>

7. Batista MAV, Bezerra Neto F, Ambrosio MMQ, Guimarães LMS, Saraiva JPB, Silva ML. Soil microbiological attributes and productivity of radish influenced by the use of spontaneous species. *Horticultura Brasileira*. 2013; 31(4):587-94. Portuguese. Available:<http://dx.doi.org/10.1590/S010205362013000400013>

8. Batista MAV, Bezerra Neto F, Silva ML, Ambrosio MMQ, Cunha JLXL. Soil-plant attributes and beet production influenced by fertilization with species of Brazilian Caatinga. *Horticultura Brasileira* 2016;34(1):31-38. Portuguese. Available:<http://dx.doi.org/10.1590/S0102053620160000100005>

9. Silva ML, Bezerra Neto F, Linhares PCF, Bezerra AKH. Production of carrot fertilized with roostertree (*Calotropis procera* (Ait.) R. Br.). *Ciências Agronômica*. 2013; 44(4):732-40. Portuguese. Available:<http://dx.doi.org/10.1590/S180666902013000400009>

10. Bezerra Neto F, Góes SB, Sá JR, Linhares PCF, Góes JB, Moreira JN. Agronomic performance of lettuce at different amounts and decomposition periods of fresh scarlet starglory. *Revista Brasileira Ciências Agrárias*. 2011; 6(2): 236-42. Portuguese. Available:<http://dx.doi.org/10.5039/agraria.v6i2a977>

11. Rêgo LGS, Martins CM, Silva EF, Silva JJA, Lima RNS. Pedogenesis and classification of soils of an experimental farm in Mossoró, Rio Grande do Norte, Brazil) *Revista Caatinga*, 2016; 29 (4): 1036-1042. Portuguese. Available:<https://dx.doi.org/10.1590/1983-21252016v29n430rc>

12. Embrapa. National Soil Research Center. 2006. Brazilian soil classification system. Rio de Janeiro: EMBRAPA Soils. 306p. Portuguese Available: <https://www.embrapa.br/solos/sibcs/classificacao-de-solos>

13. Carmo Filho, F, Oliveira, OF, Mossoró: um município do semiárido nordestino, caracterização climática e aspecto florístico. Mossoró: ESAM, 1995. 62 p. (Coleção Mossoroense, Série B). Available:http://www.scielo.br/scielo.php?script=sci_nlinks&pid=S0102...

14. Meurer EJ. Factors influencing plant growth and development. In: Novaes, RF, Alvarez VVH, Barros NF, Fontes RLF, Cantarutti RB, Neves JCL. (eds.) *Fertilidade do solo*. Viçosa: SBCS, 2007; p.65-90. Portuguese. Available:http://www.scielo.br/scielo.php?script=sci_nlinks&ref=000096&pid=S0100...lng...

15. Martins PM. Influence of temperature and drying air velocity on the content and chemical composition of lemongrass essential oil (*Cymbopogum citratus* (D.C.) STAPF). 2000; 77p. Dissertação (Mestrado- Área de Concentração em Plantas Medicinais) – Universidade Federal de Viçosa. Portuguese. Available:<http://www.ppmac.org/sites/default/files/tese-paulams.pdf>

- 422 16. Guenther E. The essential oils. 6 ed. Huntington, N.Y.: R.E. Krieger, 1972; p.63.
423 Available:[http://www.nature-helps.com/.../oilsinfo/.../The%20volatile%20oils%201948-](http://www.nature-helps.com/.../oilsinfo/.../The%20volatile%20oils%201948-Guenther%20..)
424 [Guenther%20..](http://www.nature-helps.com/.../oilsinfo/.../The%20volatile%20oils%201948-Guenther%20..)
425
- 426 17. Simões CMO, Shenkel EP, Gosmann G, Mello JCP, Mentz LA, Petrovick PR.
427 Pharmacognosy: from the plant to the medicine. (Farmacognosia: da planta ao medicamento).
428 5.ed. Porto Alegre/Florianópolis: Editora da UFRGS/Editora da UFSC: 2003; 615-656.
429 Portuguese.
430 Available:http://www.scielo.br/scielo.php?script=sci_nlinks&ref=000107&pid=S1413...lng...
431
- 432 18. Willey RW, Osiru DSO. Studies on mixture of maize and beans (*Phaseolus vulgaris*)
433 with particular reference to plant population. Journal of Agricultural Science, 1972; 77:
434 531540. Available: <http://dx.doi.org/10.1017/S0021859600025909>
435
- 436 19. Kronka SN, Banzato DA, Estat, system for statistical analysis (Estat: sistema para análise
437 estatística. Versão). 2. 3.ed. 1995; Jaboticabal: Funep, 243 p. Portuguese
438 Available:<https://www.bdpa.cnptia.embrapa.br/consulta/busca?b=ad...%22KRONKA...>
439
440
- 441 20. Jandel Scientific, Table Curve: Curve Fitting Software: *Jandel Scientific*, 1991; 280 p.
442 Portuguese.
443 Available:https://www.scielo.br/scielo.php?script=sci_nlinks&ref=000090&pid=S1806...lng..
444 .
445
- 446 21. Carvalho LM, Nunes MUC, Oliveira IR, Leal MLS. Productivity of tomato in single and
447 intercropping with aromatic and medicinal species. Horticultura Brasileira, 2009; 27 (4): 458-
448 464. Portuguese.
449 Available:<http://dx.doi.org/10.1590/S0102-05362009000400010>.
450
- 451 22. Almeida AMB. Agroeconomic viability of mint (*Mentha piperita* L.) in two successive
452 crops under jitirana doses and harvesting times. 2018. 82f. (Masters dissertation) Federal
453 University of Campina Grande. 2018. Portuguese.
454 Available:bdtd.ibict.br/vufind/Record/UFCG_e66648909a4f95848ce5a79dd5f19545
455
- 456 23. Guerra AMN, Ferreira JBA, Lima TC, Costa ACM, Tavares PRF. Intercropping
457 cultivation of lettuce with medicinal plants in the Amazonian conditions. Revista Agrarian,
458 2015; 8 (30): 369-375. Portuguese.
459 Available:<https://www.ojs.ufgd.edu.br/index.php/agrarian/article/view/2729>
460
- 461 24. Taiz L, Zeiger E. Plant Physiology, 3. ed. Porto Alegre: Artmed, 2003, 954 p.
462 Available:<https://www.bdpa.cnptia.embrapa.br/consulta/busca?b=ad...vazio...%22TAIZ>.
- 463 25. Chagas JH, Pinto JEBP, Bertolucci SKV, Santos FM, Botrel PP, Pinto LBB. Production
464 of Japanese mint due to organic fertilization at planting and cover.
465 Horticultura Brasileira, 2011; 29 (3): 412-417. Portuguese.
466 Available:<http://dx.doi.org/10.1590/S0102-05362011000300026>.

26. Sousa JS. Jitirana, silk flower and mattock as a source of green manure in coriander productivity. Dissertação (Master in Agronomy: Agroindustrial Systems). Federal University of Campina Grande, 2014; 44f. Portuguese.
Available:<https://www.periodicos.ccta.ufcg.edu.br/index.php/PPSA/article/view/70>
27. Linhares PCF, Assis JP, Sousa RP, Sá JR, Pereira MFS, Ramalho WB, Silva RIG, Silva RA, Pereira KLV. Optimized amount of hairy woodrose (*Merremia aegyptia* L.) in the productivity of coriander cultivars. Bulgarian journal of Agricultural Science, 2018; 24 (4): 654-659.
Available:https://www.researchgate.net/.../327939671_Optimized_amoun..
28. Linhares PCF, Maracajá PB, Pereira MFS, Assis JP, Sousa RP. Roostertree (*Calotropis procera*) under different amounts and periods of incorporation on yield of coriander. Revista Verde de Agroecologia e Desenvolvimento Sustentável, 2014; 9 (3): 07-12.
Available:<https://www.oaji.net/articles/2015/2238-1445534912.pdf>
29. Linhares PCF. Spontaneous vegetation as green manure on the agroeconomic performance of leaf vegetables. Thesis (PhD in Agronomy: Phytotechny), Federal Rural Semi-Arid University (UFERSA), 2009; 109f. Portuguese.
Available:<https://www.https://www.agrolink.com.br/.../vegetação%20espontânea%20como%20a...>
30. Tavella LB, Galvão RO, Ferreira LF, Araújo Neto SE, Negrieros RS. Organic farming of coriander in no-tillage system fertilized with compost using dead and living mulching. Revista Ciência Agronômica, 2010; 41 (4): 614-618. Available: <http://dx.doi.org/10.1590/S1806-66902010000400014>