

# **Effect of Irrigation Depths on the Growth of Papaya Seedlings**

## **ABSTRACT**

Water is a limiting factor for the production of Papaya tree (*Carica papaya* L.) seedlings. Its lack leads to a reduction in the absorption of nutrients and its excess provides an environment for the development of pathogens, contributing to the leaching of nutrients, besides generating wastes. The present study aimed to evaluate the effects of irrigation depths on the growth of seedlings of 'Rubi INCAPER 511' papaya. The experiment was carried out at the Federal Institute of Espirito Santo, Brazil. The experimental design was completely randomized. Six treatments were used: 4, 6, 8, 10, 12 and 14 mm d<sup>-1</sup> with 24 seedlings per treatment, totaling 144 in the experimental field. 60 days after sowing, the following morphological characteristics were evaluated: plant height, stem diameter, number of leaves, leaf area, SPAD index, shoot dry mass, dry mass of the root system and Dickson quality index. The depths between 12.5 and 14 mm d<sup>-1</sup> presented higher Dickson quality index, stem diameter, plant height and leaf number, with the 12.5 mm d<sup>-1</sup> depth being recommended for economic reasons in the production of seedlings of papaya 'Rubi INCAPER 511'.

**Keywords:** *Carica papaya* L.; seedling quality; nursery.

## **1. INTRODUCTION**

The papaya (*Carica papaya* L.) plays an important role in tropical fruit production, and its cultivation is spread worldwide [1]. India, Brazil, Mexico and Indonesia appear as the main products accounting for about 68% of world production in 2016 [2]. In Brazil, the state with the highest production is Bahia with 753,417 tons followed by Espirito Santo with production of 251,365 tons, the two states accounting for 70% of all Brazilian production in 2016 [3].

The formation of seedlings in the cultivation of papaya is an extremely important stage for the plants to express the maximum productive potential. Among the costs for the implantation of the crop, the irrigation system is highlighted because it is very onerous. Thus, the investment in alternative technologies that are sustainable, reduce production costs and improves the producer's income becomes extremely necessary [4,5].

The amount of water available is of great importance for the papaya crop, and the lack or excess interferes with the growth of the plant, besides reducing the quality and productivity of the fruits [6]. Water is a limiting factor for the production of seedlings, it is fundamental to promote seed germination, the rooting of cuttings and the development of grafted seedlings [7]. The lack of water causes a reduction in the absorption of nutrients that can lead the plant to water stress, while the excess provides favorable environment for the development of

pathogens, contributing to nutrient leaching, besides generating waste which causes social and environmental problems [8].

The application of water in plants is most often done without technical criterion, resulting in the non-expression of maximum genetic potential and decrease of the crop production. Thus, the knowledge of the production system of seedlings, in a general way, allows the elaboration of protocols that optimize the production of seedlings with better quality, decreasing the waste of water resources, energy and nutrients [9,10].

In this context, it is necessary to evaluate the efficiency of water use for the production of papaya seedlings. This study aimed to evaluate the growth of papaya seedlings of the variety 'Rubi INCAPER 511' under different irrigation depths.

## 2. MATERIAL AND METHODS

The study took place at the Federal Institute of Espírito Santo Campus Itapina, in Colatina, a city located in the north of Espírito Santo state, Brazil, located at 19° 32' 22" South latitude and 40° 37' 50" West longitude, from September 9th to November 7th in 2016. The average, maximum and minimum temperatures in the experimental period are shown in Fig.1 and the average relative humidity in Fig. 2.




Fig. 1. Average, maximum and minimum temperatures during the experiment period in Colatina, Espírito Santo




**Fig. 2. Average relative humidity during the experiment period in Colatina, Espírito Santo**

The experiment took place in an agricultural greenhouse, with linear dimensions of 25 m x 5 m with height of 3 m, covered by transparent plastic film and black polypropylene screen with 50% shading. In the interior of the greenhouse were made six individualized environments (BOX), with 2.20 m long for 1.10 m wide, isolated by transparent plastic canvas on the sides. Each BOX contained six nebulizers of the GREEN MIST model of the NaanDanJain® company spaced by 0.8 m each other and located 1 m from the seedlings, with individualized irrigation maintained by centrifugal pumps of 0.5 CV and electronically controlled, operating by pulses of 10 hours per day, with a service pressure of 2 kgf cm<sup>-2</sup>.

The treatments consisted on the application of six irrigation depths: 4, 6, 8, 10, 12 and 14 mm d<sup>-1</sup>. The experimental design was completely randomized with 24 seedlings per treatment, totaling 144 in the experimental field.

For the preparation of the seedlings were used tubes with a volume of 280 cm<sup>3</sup>, arranged alternately in trays containing 54 cells to avoid shading which could lead to seedling. All tubes were pre-sanitized with 2% sodium hypochlorite solution. The tubes were filled with commercial substrate Tropstrato HT Vegetables plus Osmocote Plus 15-9-12 (3M), at the dosage of 3 g tube<sup>-1</sup>, with the following chemical composition: N = 15%, (7% ammoniacal and 8% nitrate), P<sub>2</sub>O<sub>5</sub> = 9%, K<sub>2</sub>O = 12%, Mg = 1.3%, S = 5.9%, Cu = 0.05%, Fe = 0.46%, Mn = 0.06% and Mo = 0.02%.

The papaya seeds used were of the 'Rubi INCAPER 511' variety, which has desirable agronomic characteristics and good market acceptance. It produces fruits with dark green skin, average weight of 1.47 kg, orange-reddish flesh and soluble solids of 10.2 °brix [11]. Two seeds were sown per tube and, at 10 days after sowing, thinning was carried out, with only one seedling per tube.

At 60 days after planting, the seedlings were evaluated for the following morphological characteristics: a) plant height (PH), measured with a graduated ruler, in cm; b) stem diameter (SD), measured 2 cm above the collection of the seedling, with pachymeter, in mm; c) length of the root system (LRS), measured from the base of the collection to the largest root length, in cm, with graduated ruler; d) number of leaves (NL), defined by the total leaf

count of the seedling; e) leaf area (LA), as measured by the LI-3100 equipment, in cm<sup>2</sup>; f) SPAD index, measured with SPAD 502 Plus equipment; g) dry mass of the aerial part (DMAP) and dry mass of the root system (DMRS), measured in g; h) Dickson Quality Index (DQI) according to Dickson *et al.* [12] given by:

$$DQI = \frac{DMAP + DMRS}{\frac{PH}{SD} + \frac{DMAP}{DMRS}}$$

105  
106  
107

In order to obtain dry mass of the aerial part (DMAP) and dry mass of the root system (DMRS) the plants were dried in a ventilated oven at 65 °C for 72 hours and weighed in a semi-analytical balance.

111  
112  
113  
114  
115  
116  
117

The results were submitted to analysis of variance and the effects of treatments were evaluated by the F test at 5% probability. When significant, regression models were fitted that best applied to the effects of the irrigation blade on the characteristics evaluated. The maximum points were obtained through the primary derivative of the regression equations.

### 118 3. RESULTS AND DISCUSSION

119

After analysis of variance, a significant difference ( $p < 0.05$ ) was observed for all the morphological characteristics evaluated in relation to the applied irrigation depths, attesting that they affected the growth of papaya seedlings of the variety 'Rubi INCAPER 511'.

123

Plant height (PH) increased linearly in relation to applied irrigation depths, with coefficient of determination ( $R^2$ ) of 0.8383 (Fig. 3A) showing that 83% of the height of the plant was explained by the irrigation depths. According to Klar and Jadoski [13], the water deficit causes a reduction in the growth of the seedlings. Higher plant height is more desirable because it offers more resistance to mechanical damage. Melo *et al.* [14], suggest that this characteristic is an important factor for the transportation of the seedlings, being those that have low stature are constantly damaged, affecting their development and, consequently, the final production.

131

132

The stem diameter (SD) showed quadratic adjustment, with maximum point in the irrigation depth of 13.92 mm d<sup>-1</sup> with a diameter of 6.42 mm and  $R^2$  of 0.9181 (Fig. 3B). Carneiro [15] cites that a larger diameter contributes to a smaller tipping of seedlings in the field. In this way, higher diameters are more desired for papaya seedlings. According to Melo *et al.* [14], the increase in diameter has a correlation with the amount of photoassimilates produced by the plant, being highlighted in the field, due to its higher vigor. Carneiro [15] and Dassie *et al.* [16] reported that plant height and stem diameter are easy to evaluate morphological characteristics used as the basis for the transplanting of seedlings to the field, the ones with the highest values of these characteristics being the most desirable.

141




**Fig. 3. Variation of plant height (A) and stem diameter (B), under different irrigation depths in papaya seedlings**

The length of the root system (LRS) presented an increasing linear adjustment in relation to the increase of the irrigation depths, with  $R^2$  of 0.9566 (Fig. 4). According to Scalón *et al.* [17], under water deficit conditions, the water concentrates only on the surface of the substrate, so some plant species have adaptation mechanisms to tolerate the available water quality. In this work, smaller values for the LRS were found in less water availability. This fact can be explained by the abscisic acid hormone, which is synthesized under conditions of water stress, stimulating the growth and lateral formation of the roots, optimizing the water absorption by the plants [18].




**Fig. 4. Variation of root system length under different irrigation depths in papaya seedlings**

157

158 The number of leaves (NL) presented quadratic effect, with determination coefficient of  
 159 0.7495 (Fig. 5A). The maximum number of leaves (around 7) was observed in the irrigation  
 160 depth of 13.08 mm d<sup>-1</sup> (Fig. 5A). So important for the development of plants, the number of  
 161 leaves indicates seedlings with more aptitude for the field conditions, because they present a  
 162 larger photosynthetically active area [14]. Klar and Jadoski [9] and Diniz *et al.* [19] suggest  
 163 that the stress state affects the metabolism of the plants, translocating the photoassimilates  
 164 to the osmotic adjustment, reducing the number of leaves.


165

166 The leaf area (LA) presented a simple linear effect in relation to the applied irrigation depths,  
 167 with R<sup>2</sup> of 0.9014 (Fig. 5B). Similar results were found by Lopes *et al.* [8] that studied the  
 168 effect of the irrigation depth on the development of *Eucalyptus grandis* W. (HILL ex.  
 169 MAIDEN) seedlings, showing that the leaves had reduced size under water deficit  
 170 conditions. According to Figueirôa *et al.* [20], the increase in water availability has a direct  
 171 influence on leaf area growth.

172

173 Mott *et al.* [21] reported that the environment directly interferes with leaf size and so, an  
 174 environment with high humidity implies larger leaves, whereas in water scarce environments  
 175 it infers into smaller leaves, reducing warming and leaf transpiration on warmer days. Larger  
 176 leaves produce more photosynthesis, but they present a larger surface, which implies higher  
 177 evaporation, causing exhaustion of the water present in the soil, besides having excessive  
 178 absorption of solar energy which causes damage, so the plants reduce their leaf area when  
 179 in stress due to lack of water [18].

180



181

182 **Fig. 5. Variation of number of leaves (A) and leaf area (B), under different**  
 183 **irrigation depths in papaya seedlings**


184

185 The chlorophyll content index SPAD presented linear decreasing behavior in relation to the  
 186 increase of the irrigation depths, with R<sup>2</sup> of 0.8541 (Fig. 6). According to Zuffo *et al.* [22], the  
 187 SPAD index determines the amount of green present in the leaves of the plant, being a very  
 188 efficient way of determining the leaf nitrogen content, since the chlorophyll (green  
 189 pigmentation) of the plants composition presents the nitrogen molecule. Its use allows the

190 generation of important information for decision making, as well as being a low-cost, non-  
191 destructive measure.

192

193 It was observed that as the irrigation depths increased, there was a significant reduction in  
194 the SPAD index. Lopes *et al.* [8] say that the excessive application of water in the production  
195 of seedlings implies leaching of the nutrients. Possibly, this fact explains the decrease in the  
196 green content of the plants influenced by the leaching of the nitrogen present in the  
197 substrate. The nutrient leaching caused by the constant use of irrigation is very common,  
198 prolonging the development time of the seedlings as a result of the lower use of fertilization  
199 [23].



200

201 **Fig. 6. Variation of the SPAD index, under different irrigation depths in papaya**  
202 **seedlings**

203

204 The dry mass of the aerial part (DMAP) presented linear regression in relation to applied  
205 irrigation depths, with  $R^2$  of 0.9079 (Fig. 7A). Plants under smaller irrigation depths have  
206 reduced DMAP. Similar results were obtained by Santiago *et al.* [24], that studied the growth  
207 of young plants of *Mimosa caesalpinifolia* BENTH cultivated under water stress, showing a  
208 decrease of the dry mass of the aerial part in plants submitted to water restriction.

209

210 The dry mass of the root system (DMRS) showed a quadratic effect with a maximum point  
211 on the irrigation depth of 12.65 mm d<sup>-1</sup> with weight of 0.72 g and  $R^2$  of 0.9354 (Figure 7B).  
212 Under the condition of lack or excess water in the root system, plants have limitations,  
213 increasing the amount of water in the roots contributes to an environment with low oxygen  
214 concentrations, reducing the photosynthetic rate of the plant, impairing the transport of  
215 photoassimilates to the root system and, consequently, the reduction in the dry mass of the  
216 root system [25,26].

217




**Fig. 7. Variation of the dry mass of the aerial part (A) and the dry mass of the root system (B) of papaya seedlings, under different irrigation depths**

The Dickson quality index (DQI) is a balanced formula that relates morphological characteristics, with plant height, stem diameter and dry biomass [27]. The DQI showed a quadratic effect with a higher index of 0.43 in the irrigation depth 12.66 mm d<sup>-1</sup> and a coefficient of determination of 0.9524 (Fig. 8). Posse *et al.* [28], evaluating seedlings of yellow passion fruit under different irrigation depths, detected DQI between 0.15 and 0.27, showing that values below 0.20 are not desirable. According to Binotto *et al.* [29], the DQI can be used to identify which seedlings have greater adaptability and development, conferring greater chances of survival in the field, and as higher this index as greater the robustness and vigor of the seedlings.




**Fig. 8. Variation of the Dickson quality index under different irrigation depths in papaya seedlings**

Posse *et al.* [28], reported that DQI should not be analyzed in isolation to determine the quality of a seedling. Characteristics such as plant height, number of leaves and stem diameter, besides being non-destructive methods, should be taken into account to ensure greater survival of the seedlings in the field.

#### **4. CONCLUSION**

Irrigation depths interfered with the growth of papaya seedlings, and the depths between 12.5 and 14 mm d<sup>-1</sup> presented higher Dickson quality index, stem diameter, plant height and number of leaves, which were used to identify better quality seedlings. For reasons of energy conservation and water resources the 12.5 mm d<sup>-1</sup> depth is the most recommended for the production of seedlings of the variety 'Rubi INCAPER 511'.

#### **REFERENCES**

1. Souza JMA, Ataíde EM, Silva MS. Post-harvest quality and correlation between physical and chemical characteristics of papaya fruits commercialized in Serra Talhada - PE. *Magistra*. 2014;26(4):554-560.
2. FAO – Food and Agriculture Organization of the United Nations. The Agricultural Production; 2016. Available: <http://www.faostat.org>.
3. IBGE – Instituto Brasileiro de Geografia e Estatística. Municipal Agricultural Production; 2016. Available: <http://www.sidra.ibge.gov.br>.
4. Danner MA, Citadin I, Junior AAF, Assmann AP, Mazaro SM, Sasso SAZ. Formation of jabuticabeira (*Plinia* sp.) Seedlings on different substrates and container sizes. *Revista Brasileira Fruticultura*. 2007;29(1):179-182.
5. Weckner FC, Campos MCC, Nascimento EP, Mantovanelli BC, Nascimento MF. Evaluation of the papaya seedlings under the effect of the application of different compositions of biofertilizers. *Revista da Universidade Vale do Rio Verde*. 2016; 14(1):700-706.
6. Marin SLD, Gomes JA, Salgado JS, Martins DS, Fullin EA. Recommendations for the cultivation of papaya from the groups Solo and Formosa in the State of Espírito Santo. 4.ed. Vitória: EMCAPA. 1995; 57.
7. Fronza D, Hamann JJ. Nurseries and seedlings propagation. Santa Maria: UFSM, Colégio Politécnico: Rede e-Tec Brasil. 2015; 142.
8. Lopes JLW, Guerrini IA, Saad JCC. Effects of irrigation slides on the production of *Eucalyptus grandis* W. (HILL EX. MAIDEN) seedlings on coconut fiber substrate. *Irriga*. 2005;10(2):123-134.
9. Gruber YBG. Optimization of the irrigation depth in the production of clonal eucalyptus seedlings (*Eucalyptus urophylla* x *Eucalyptus grandis* and *Eucalyptus urophylla* var. *Plathyphylla*). 2006; 144 p. Master Dissertation. Universidade de São Paulo; 2006.

- 285  
286 10. Delgado LGM, Silva RBG. Silva, MR. Morphological quality of ingá seedlings under  
287 different water management. *Irriga*. 2017;22(3):420-429.  
288
- 289 11. Cattaneo LF, Costa, AFS, Serrano LAL, Costa NA, Fanton CJ, Bravim AJB. 'Rubi  
290 INCAPER 511': First Papaya variety from the "Formosa" Group for Espírito Santo. Vitória,  
291 ES: INCAPER. 2010; 6.  
292
- 293 12. Dickson A, Leaf AL, Hosner JF. Quality appraisal of white spruce and white pine  
294 seedling stock in nurseries. *Forest Chronicles*. 1960;36:10-13.  
295
- 296 13. Klar AE, Jadoski, SO. Effects of irrigation and soil cover by black polyethylene on the  
297 morphological characteristics of the pepper. *Irriga*. 2002;7(3):154-167.  
298
- 299 14. Melo AS, Costa CX, Brito MEB, Viégas PRA, Silva Júnior CD. Production of papaya  
300 seedlings in different substrates and phosphorus doses. *Revista Brasileira de Ciências*  
301 *Agrárias*. 2007;2(4):257-261.  
302
- 303 15. Carneiro JGA. Production and quality control of seedlings. Curitiba: UFPR/FUPEF. 1995;  
304 451.  
305
- 306 16. Dassi LA, Aleman CC, Moreira ACM, Mignacca FA, Zanfolin PRL, Carvalho PR.  
307 Irrigated production of young finger pepper seedlings (*Capsicum baccatum*). *Colloquium*  
308 *Agrariae*. 2017;13(3):128-131.  
309
- 310 17. Scalón SPQ, Mussury RM, Euzébio VLM, Kodama FM, Kissmann C. Water stress in the  
311 metabolism and initial growth of Mutambo seedlings (*Guazuma ulmifolia* Lam.). *Ciência*  
312 *Florestal*. 2011;21(4):655-662.  
313
- 314 18. Taiz L, Zeiger E, Moller IM, Murphy A. Physiology and plant development. 6. ed. Porto  
315 Alegre: Artmed. 2017;858.  
316
- 317 19. Diniz GL, Sales GN, Sousa VFO, Andrade FHA, Silva SS, Nobre RG. Production of  
318 papaya seedlings under water salinity irrigation and phosphate fertilization. *Revista de*  
319 *Ciências Agrárias*. 2018;41(1):218-228.  
320
- 321 20. Figueirôa JM, Barbosa DCA, Simabukuro EA. Growth of young plants of *Myracrodruon*  
322 *urundeuva* Allemão (Anacardiaceae) under different water regimes. *Acta Botanica Brasilica*.  
323 2004;18(3):573-580.  
324
- 325 21. Mott KA, Gibson AC, O'leary JW. The adaptive significance of amphistomatic leaves.  
326 *Plant, Cell Environment*. 1982;5(6):455-460. □  
327
- 328 22. Zuffo AM, Andrade FR, Schossler TR. Milhomem M, Piauilino AC. Efficiency in the  
329 indirect determination of leaf nitrogen from the SPAD index. *Enciclopédia Biosfera*.  
330 2012;8(15):802-820.  
331
- 332 23. Fagundes MCP, Cruz MCM, Carvalho RP, Oliveira J, Soares BC. Hydroabsorbent  
333 polymer in the reduction of leached nutrients during the production of Yellow Passion fruit  
334 seedlings. *Revista Caatinga*. 2015;28(1):121-129.  
335
- 336 24. Santiago AMP, Nogueira RJMC, Lopes EC. Growth of young plants of *Mimosa*  
337 *caesalpiniiifolia* Benth grown under water stress. *Revista Ecosistema*. 2002; 26(1): 23-30.

338

339 25. Martinazzo EG, Perboni AT, Oliveira PV, Bianchi VJ, Bacarin MA. Photosynthetic activity  
340 in plum plants submitted to water deficit and flooding. *Ciência Rural*. 2013;43(1):35-41.

341

342 26. Queiroz TB, Rocha SMG, Fonseca FSA, Alvarenga ICA, Martins ER. Effects of water  
343 deficit on eucalyptus seedlings. *Irriga*. 2017;22(4):659-674.

344

345 27. Silva AK, Costa E, Santos ELL, Benett KSS, Benett CGS. Production of seedlings of  
346 'Formosa' papaya under thermoflective mesh effect and substrates. *Revista Brasileira de*  
347 *Ciências Agrárias*. 2013;8(1):42-48.

348

349 28. Posse RP, Valani F, Gonçalves AMS, Oliveira EC, Louzada JM, Quartezani WZ, Leite  
350 MCT. Growth and Quality of Yellow Passion Fruit Seedlings Produced under Different  
351 Irrigation Depths. *Journal of Experimental Agriculture International*. 2018;22(4):1-11.

352

353 29. Binotto AF, Lúcio AD, Lopes SJ. Correlations between growth variables and the Dickson  
354 quality index in forest seedlings. *Revista Cerne*. 2010;16(4):457-464.

355