Original Research Article

1 2 3

Urban extension in Calabar: A remotely sensed assessment

4 5 6

Abstract

7 Urban places in the developing world, like the advanced world, are experiencing 8 unprecedented extension or growth, although mostly in an uncontrolled manner. Calabar is no exception to such urban extension, with notable sprawl especially with increased 9 10 densification within the city and expansion along the fringes. This study aimed at assessing 11 landuse/ land-cover (LULC) changes in Calabar between the year 2000 and 2018. 70m ETM+ Landsat imager of both years were acquired from the Landsat Look platform. The 12 13 imageries were subjected to an unsupervised classification using the Iso Cluster and 14 Maximum Likelihood Classification tool. Measurements were done on the raster outputs to 15 allow for a comparison of the LULC statistics which assisted in identifying the pattern 16 extension in Calabar over the period. Results showed that in the year 2000, out of the total 17 164.3 square kilometers (sqkm) covered by the city, the extent of urban built-up was 28.7 18 sqkm and 62.2 sqkm for green areas. However, in the year 2018, the urban built-up area 19 increased to 44.8 sqm and green areas witnessed a reduction to 52.7 sqkm. Wet lands also 20 reduced in the area by 5.2 sqkm. The spate of urban extension and encroachment into green 21 fields and wet lands in Calabar is obvious and somewhat inevitable as by this study. It is thus recommended that necessary measures be taken by relevant government 22 23 agencies to monitor and manage this gradual extension, such that the development is 24 organized and sustainable.

25 Keywords: Urban extension, landuse, land-cover, geographic information systems, remote

add reference

26 sensing, Calabar.

Introduction

27

28

Globally, land cover today is altered principally by direct anthropogenic factors like

agriculture and livestock raising, forest harvesting and urban and suburban construction and 29

development. Due to rapid human activities, the earth surface is being progressively altered in such manner that man's existence on earth and his use of land to being greatly a flected

31

The fast pace of urbanization has been shown to be a 32

UNDER PEER REVIEW

63 64

33 serious global problem and is more evident in most of the developing countries. There is also 34 every indication that the trend will continue, adding approximately two billion people to the 35 urban population of the presently less-developed nations in the next 30 years (United Nations 36 Environment Programme, 2002) 37 The world is becoming increasingly urbanized, with 45 percent of the population already living in the urban areas in the year 2000. The projection was that half of the world 38 39 will live in urban areas by 2007 (Arnfield, 2003). It was also estimated that by the year 2025, 40 60 percent of the world's population will live in cities (United Nations Population Fund, 41 1999). Land transformation has been asserted to be one of the most important fields of human 42 induced environmental transformation (Fasal, 2000). 43 Urbanization is one of the several anthropogenic activities that impact on land use/land cover. 44 Urban population has been growing more rapidly than rural worldwide, particularly in 45 developing countries (Lambin, Geist, and Lepers, 2003). It is measured by the rate at which 46 the spatial extent of an urban settlement extends. In most countries, urban growth is 47 recognized as a crucial phenomenon of economic growth and social change, as it offers 48 increased opportunities for employment, specialization, production, goods and services 49 (Odjugo, Enaruvbe, and Isibor, 2015), which in turn initiates a large number of people 50 migrating from rural to urban areas (Abebe, 2013). 51 Several empirical studies have shown that unplanned changes of land use due to urbanization have become a major problem (Zhao, Dickson and Tian, 2004; Nanda, 2005). Most land use changes occur without a clear and logical planning without attention to their environmental 52 53 impacts. Major flooding, air pollution in large cities as well as deforestation, urban growth, 54 soil erosion, desertification, are all consequences of a mismanaged planning, without 55 considering environmental impacts of development planes. 56 57 This study focuses on urban extension in Calabar. Calabar is the capital city of Cross River State, in the southern region of Nigeria. The city lies between Longitudes 8°18'00"E to define 8°24'00"E and Latitudes 4°54'00"N to 5°04'00"N 58 8°24′00″E, and Latitudes 4°54′00″N to 5°04′00″N, sandwiched to between Odukpani(LGA) to 59 the north, the Calabar River to the west, Great Kwa River to the east, and the creeks of the 60 61 Cross River as it empties into the Atlantic Ocean in the South (Figure 1). The Metropolis covers an approximate land area of 137.039 square kilometers (sqkm) and had a population 62

of 328,878 in 1991, and 375,196 in 2006 according to the National Population Commission

(NPC) and a projected population of 529362 in 2015 (Njoku, Okon, Itu and Ahwen, 2017).

65 Calabar has witnessed observable urban extension over time. The built-up area continues to extend outward and have consumed prior agricultural and wet lands at a break-neck pace. Todd reference 66 67 Hectares of green areas are now covered by concrete and asphalt, as new roads are created and existing ones are extended. Over 5000 hectares of greenery have been taken over by built up 68 69 activities & Ekorinim, Esuk Utan, Edim-Otop, Anantigha, and Ikot Effanga areas of the metropolis (Atu, Offiong, Eni, Eja, and Esien, 2012). 70 accomodate 71 As the population of Calabar increases, so also is the desire to these population 72 thresholds through urbanization. The extension of the urban area in Calabar has resulted not 73 only in depletion of natural resources, but deterioration of the environment. The unregulated 74 and haphazard growth of urban development has adversely affected Calabar's ecosystem 75 which has putered to indirectly publication climate, all things and eventually had see feen 76 This study is constant at assessing spatial and temporal landuse and 77 land-cover changes in Calabar between the year 2000 and 2018, using remotely sensed data and GIS techniques. GIS-based multi-temporal land use data and analyses provided a 78 79 historical vehicle for determining and evaluating long-term changes in landuse due to urbanization. The collection of remotely sensed data facilitates the synoptic analyses of 80 81 changes on the earth surface at local, regional and global scales over time (Wilkie and Finn, 82 1996).




Figure 1: Map of Calabar Metropolis

Materials and Methods

The methodology of this study incorporated a reconnaissance survey, data acquisition, data processing and data analysis. The reconnaissance survey aimed at getting the researcher acquainted with the existing physical characteristics of the study area. A knowledge that was very useful for the selection of training sites before classification. The types of data used include Landsat imageries which were obtained from *Landsat Look* platform. Their attributes are presented in **Table 1.** Relevant literature materials were obtained from text books, journals and other existing literatures that are related to the research problem.

Table 1: Attributes of acquired satellite imager

	Data Type	Date of Acquisitions	Resolution	Source	
1.	Landsat image	2000	70m ETM+	Landsat Look	
2.	Landsat image	2018	70m ETM+	Landsat Look	

Imag

Image processing and classification

There was no need for georeferencing the images since they were already otho-rectified. The images were clipped with the boundary data of Calabar, using the clip tool in the ArcMap platform. However, the images were geometrically corrected to Universal Transverse Mercator (UTM) Zone 32 North coordinate system on the same platform. To detect changes in the land use/cover at different years, post classification comparison of the change detection

104 techniques was used.

The boundary data of Calabar metropolis was used to clip the Landsat imageries. The clipping helped to remove the extents outside the boundary of the satellite imageries (depicted in false color in Figure 2 and 3). After clipping, ERDAS imagine software was used for the pixel-based classification. The were subjected to an unsupervised classification using the Iso Cluster (IC) and Maximum Likelihood Classification (MLC) spatial analyst tool. The IC combines the functionalities of the IC and the MLC, while the MLC performs classification on a set of raster bands using a signature file from the IC tool as the input for MLC. The ArcMap software was afterwards used for the final color and the matter of the input for MLC.

the ERDAS outputs.

The classified raster output was converted to vector (polygons) to allow for measurements to be done. The area coverage of each of the LULC classes was measured sqkm, for each of the years under consideration using the calculate geometry tool in the same ArcMap platform. A comparison of the land cover statistics assists in identifying the change in sqkm/percentage, trend, and rate of change in Calabar over the period. In this study, in line with Anderson, Hardy, Roach and Witmer (1976) land-use/land-cover classification scheme in (Wakirwa, 2015), the various land-use/land-cover (LULC) types are modified and generalized into 4 classes within the study area as presented on Table 2.

lR .	REVIEW					
N	Jefine Jean Table 2: 1	LULC classification scheme adopted				
(6/N)	Land Use Category	Description				
l	Built-up	Land used for residential and transportation/communication purposes (i.e., settlements and roads, high residential area, industry and administrative block).				
3	Wet lands	Land covered characteristically saturated; a marsh.				
4	Water body	Areas covered by body of water e.g. dam, lake, rivers and swamps				
5	Green area	Areas that are spatially cultivated e.g., farmland, irrigation areas etc. as well as urban greenery, grasses, shrubs and grass-like plants.				




Figure 3: Landsat imagery of Calabar in 2018

159

Results and Discussions

129 Land use and land cover scenario in Calabar from 2000 to 2018 130 To assess the LULC situation of Calabar in the past and present, the LULC analysis was 131 executed. Tables, maps and charts were used to illustrate the LULC status quo at Calabar in 2000 and 2018 respectively. As symbolized by the legend, which brown is used to represent 132 the built-up area, the light green for green area, blue represent water body and the dark 133 green the forest areas. 134 As observed from the analyses outputs presented in Table 3, and Figures 4-5, there were 135 evident changes in the LULC of Calabar during the 18-year period considered. From the year 136 2000 to 2018, there was little change in the extent of the water bodies with only a slight 137 decrease 0.9 percent. The Calabar and Great Kwa rivers are tidal rivers mostly influenced 138 139 by their closeness to the Atlantic Ocean. The rivers are minimally influenced by urban 140 activities, as the wet land bordering them still shield the rivers from direct urban extension 141 impacts. 142 143 Within the same period under review, built-up areas increased by 16.1 sqm, with a coverage of 28.7 sqkm in the year 2000 and 44.8 in 2018. As pictured Figure 5, the city visibly 144 extended to the northern fringes, with new developments in the 8th mile, Ikot Nkebre, 145 Adiabo, and other areas at the northern fringes. There was also no significant extension in the 146 southern area which houses the Calabar South(LGA) The direct impact of the urban built-up 147 148 extension is on the green areas, which saw a reduction of 9.5 sqkm from an initial 62.2 sqkm 149 in the year 2000. The wetlands on both sides of the city also reduced by 5.2 sqkm. The extension in the urban buil up follows from the swelling population and increased socio-150 151 economic activities within the study area. Notably, the urban extension evident in these fringe 152 areas of Calabar are haphazard and uncoordinated, thus requiring the attention of relevant 153 agencies to ensure sustainable urban development in the area. 154 155 156 157 158

Table 3: LULC characteristics of Calabar in the year 2000 and 2018

2000			2018			
Land use class	Area coverage (Sqkm)	Percentage	Area coverage (Sqkm)	Percentage	Area coverage (+/-)	Percentage (+/-)
Built up	28.7	17.5	44.8	27.4	16.1	9.9
Green areas	62.2	37.8	52.7	32	-9.5	-5.8
Wet lands	55.9	34	50.7	30.8	-5.2	-3.2
Water bodies	17.5	10.7	16.1	9.8	-1.4	-0.9
Total	164.3	100	164.3	100	3705020	




Figure 4: Output of LULC analysis in the year 2000




Figure 5: Output of LULC analysis in the year 2018

162

163

164

165 166

177

179

180 181

182

183

184

185

186

187

188 189

190

Conclusion and Recommendation

- Typical to urban centers, urban extension in Calabar has been triggered by increased human 169
- 170 activities, resulting in LULC changes. The impacts of these changes are worrisome in the
- 171 face of uncoordinated planning policies and implementation. The application of remote
- 172 sensing and GIS provided quantification, estimation and understanding of LULC changes in
- 173 Calabar. There was apparent extension in the urban built-up area of Calabar with the greenery
- 174 mostly suffering the impacts of the extension.
- of this from the findings from the study, the proper recommend the study. 175
- 176 Landuse planning should be instituted and implemented in Calabar to ensure that the usage of land especially in the fringe is sustainable. This effort should involve sectoral integration of the relevant state ministries, boards and bodies of the Calabar 178 Municipal Council.
 - ii. Intensive sensitization should also be embarked upon and pursued holistically, with a view to ensuring that the inhabitants of the study area understand the negative effects and consequences that are associated with uncoordinated landuse development.
 - iii. Deliberate efforts such as the declaration, reservation and preservation of the ecologically fragile greenery and wetland areas of Calabar should be instituted and logically pursued by government.
 - iv. Finally, smart development of diverse-mixed landuse should be encouraged to reduce the rapid rate and large amount of urban land being converted for construction of homes, offices and commercial buildings.

191 References

- 192 Abebe, A.G. (2013). Quantifying Urban Growth Pattern in Developing Countries using
- 193 Remote Sensing and Spatial Metrics: A case Study in Kampala Uganda. Thesis submitted
- 194 to the faculty of Geo-information science and earth observation, University of Twente.
- 195 Pp 1-4
- 196 Arnfield AJ (2003). Two decades of urban climate research: A review of turbulence,
- 197 exchanges of energy and water, and the urban heat island. Int. J. Climatol., 23: 1-26.
- 198 Atu, J.E., Offiong, R.A., Eni, D.I., Eja, E.I. and Esien, O.E. (2012). The Effects of Urban
- 199 Sprawl on Peripheral Agricultural Lands in Calabar, Nigeria. International Review of
- 200 Social Sciences and Humanities. 2(2):68-76.
- 201 Fasal S (2000). Urban expansion and loss of agricultural land- A GIS based study of
- 202 Saharanpur City, India. Environ. Urbanisation, 12(2): 133-149.

UNDER PEER REVIEW

- Accessed 03 July 2018

- 203 Lambin, E.F., Geist, H., and Lepers, E. (2003) Dynamics of landuse and land Cover Changes
- in Tropical Regions. Annual Review of Environment Resources, 28: 205 -41 (online)
- availablehttp://www.globalrestorationnetwork.org/uploads/files/LiteratureAttachments/93_dy
- 206 namics-of-landuse- and-land-cover-change-in-tropical-regions.pdf
- 207 Nanda T (2005). Urban Sprawl and occupational change in Raipur City. (Unpublished M.Sc.
- 208 Thesis, Department of Geography, University of New Delhi, Delhi.
- Njoku, C. G., Okon, I., Itu, Y-C. & Ahwen, V. O (2017). Population Growth and Pipe-Borne
- 210 Water Supply in Calabar Metropolis: Finding the Balance. *International Journal of Research*
- 211 in Geography. 3:3, PP/51-59.
- Odjugo, P.A.O., Enaruvbe, G.O. and Isibor, H.O. (2015). Geospatial Approach to Spatio-
- 213 temporal Pattern of Urban Growth in Benin City, Nigeria. African Journal of
- Environmental Science and Technology. 9(3): 166-175
- 215 United Nations Environment Programme (2002). Global Environment Outlook 3. Available
- online at: http://www.unep.org/GEO/geo3/index.htm.
- United Nations Population Fund (1999). The State of World Population, pp. 76. New York:
- 218 United Nations Publications.
- Wakirwa, A. (2015). Analysis of Urban Sprawl Using Geospatial Techniques in Gombe
- 220 Metropolis, Gombe State, Nigeria. A Dissertation Submitted to the School of Postgraduate
- 221 Studies, Ahmadu Bello University, Zaria.
- Wilkie, D.S. and Finn, J.T. (1996) Remote Sensing Imagery for Natural Resources
- 223 Monitoring. Columbia University Press, New York. p. 295.
- 224 Zhao LM, Dickson RE, Tian YH (2004). Evidence for a significant urbanization effect on
- urban climate in china. P Natl. Acad. Sci. USA, 101: 9540- 9544.