

Urban extension in Calabar: A remotely sensed assessment

Abstract

Urban places in the developing world, like the advanced world, are experiencing unprecedented extension or growth, although mostly in an uncontrolled manner. Calabar is no exception to such urban extension, with notable sprawl, especially with increased densification within the city and expansion along the fringes. This study aimed at assessing land use/ land-cover (LULC) changes in Calabar between the year 2000 and 2018. 70m ETM+ Landsat imagery of both years was acquired from the Landsat Look platform. The imageries were subjected to an unsupervised classification using the Iso Cluster and Maximum Likelihood Classification tool. Measurements were done on the raster outputs to allow for a comparison of the LULC statistics which assisted in identifying the pattern extension in Calabar over the period. Results showed that in the year 2000, out of the total 164.3 square kilometers (sqkm) covered by the city, the extent of urban built-up was 28.7 sqkm and 62.2 sqkm for green areas. However, in the year 2018, the urban built-up area increased to 44.8 sqm and green areas witnessed a reduction to 52.7 sqkm. Wetlands also reduced in the area by 5.2 sqkm. The spate of urban extension and encroachment into green fields and wetlands in Calabar is obvious, and somewhat inevitable, as shown by this study. It is thus recommended that necessary measures be taken by relevant government agencies to monitor and manage this gradual extension, such that the development is organised and sustainable.

Keywords: Urban extension, land use, land-cover, geographic information systems, remote sensing, Calabar.

Introduction

Globally, land cover today is altered principally by direct anthropogenic factors like agriculture and livestock raising, forest harvesting and urban and suburban construction and development (Owoeye & Ibitoye, 2016). Due to rapid human activities, the earth surface is being progressively altered in such a manner that man's existence on earth and his use of land is being greatly affected. The fast pace of urbanization has been shown to be a serious global

33 problem and is more evident in most of the developing countries. There is also every
34 indication that the trend will continue, adding approximately two billion people to the urban
35 population of the presently less-developed nations in the next 30 years (United Nations
36 Environment Programme, 2002)

37 The world is becoming increasingly urbanised, with 45 percent of the population already
38 living in the urban areas in the year 2000. The projection then was that half of the world will
39 live in urban areas by 2007 (Arnfield, 2003). It was also estimated that by the year 2025, 60
40 percent of the world's population will live in cities (United Nations Population Fund, 1999).
41 The land transformation has been asserted to be one of the most important fields of human-
42 induced environmental transformation (Fasal, 2000).

43 Urbanization is one of the several anthropogenic activities that impact on land use/land cover.
44 Urban population has been growing more rapidly than rural population worldwide,
45 particularly in developing countries (Lambin et al., 2003). It is measured by the rate at which
46 the spatial extent of an urban settlement extends. In most countries, urban growth is
47 recognized as a crucial phenomenon of economic growth and social change, as it offers
48 increased opportunities for employment, specialization, production, goods and services
49 (Odjugo, Enaruvbe and Isibor, 2015), which in turn initiates a large number of people
50 migrating from rural to urban areas (Abebe, 2013).

51 Several empirical studies (Zhao, Dickson and Tian, 2004; Nanda, 2005) have shown that
52 unplanned changes of land use due to urbanisation become a major problem. Most land use
53 changes occur without a clear and logical planning activity and without attention to their
54 environmental impacts. In metropolitan cities, major flooding, air pollution as well as
55 deforestation, urban growth, soil erosion, desertification, are consequences of mismanaged
56 planning without considering the environmental impacts of development plans.

57 Calabar is the capital city of Cross River State, in the southern region of Nigeria. The city lies
58 between Longitudes 8°18'00"E to 8°24'00"E, and latitudes 4°54'00"N to 5°04'00"N,
59 sandwiched between Odukpani Local Government Area (LGA) to the north, the Calabar River
60 to the west, Great Kwa River to the east, and the creeks of the Cross River as it empties into
61 the Atlantic Ocean in the South (Figure 1). The Metropolis covers an approximate land area
62 of 137.039 square kilometers (sqkm), and had a population of 328,878 in 1991, and 375,196
63 in 2006 according to the National Population Commission (NPC), and a projected population
64 of 529,362 in 2015 (Njoku, Okon, Itu and Ahwen, 2017). Calabar has witnessed observable

65 urban extension over time. The built-up area continues to extend outward and has consumed
66 prior agricultural and wetlands at a break-neck pace. Hectares of green areas are now covered
67 by concrete and asphalt, as new roads are created and existing ones are extended. Over 5000
68 hectares of greenery has been taken over by built-up activities in the Ekorinim, Esuk Utan,
69 Edim-Otop, Anantigha, and Ikot Effanga areas of the metropolis (Atu et al., 2012).

70 As the population of Calabar increases, so also is the desire to accommodate these population
71 thresholds through urbanisation. The extension of the urban area in Calabar has resulted not
72 only in depletion of natural resources but is deterioration of the environment. The unregulated
73 and haphazard growth of urban development has adversely affected Calabar's ecosystem
74 (Offiong & Eteng, 2014). This study is aimed at assessing spatial and temporal land use and
75 land-cover changes in Calabar between the year 2000 and 2018, using remotely sensed data
76 and GIS techniques. GIS-based multi-temporal land use data and analyses provide a historical
77 vehicle for determining and evaluating long-term changes in land use due to urbanisation.
78 The collection of remotely sensed data facilitates the synoptic analyses of changes on the
79 earth surface at local, regional and global scales over time (Wilkie and Finn, 1996).




Figure 1: Map of the Calabar Metropolis

Materials and Methods

The methodology of this study incorporated a reconnaissance survey, data acquisition, data processing and data analysis. The reconnaissance survey aimed at getting the researcher acquainted with the existing physical characteristics of the study area. A knowledge that was very useful for the selection of training sites before classification. The types of data used include Landsat imageries which were obtained from *Landsat Look* platform. Their attributes are presented in **Table 1**. Relevant literature materials were obtained from textbooks, journals and other existing literature that are related to the research problem.

Table 1: Attributes of acquired satellite imagery

	Data Type	Date of Acquisitions	Resolution	Source
1.	Landsat image	2000	70m ETM+	Landsat Look
2.	Landsat image	2018	70m ETM+	Landsat Look

Image processing and classification

There was no need for georeferencing the images since they were already ortho-rectified. The images were clipped with the boundary data of Calabar using the clip tool in the ArcMap platform. However, the images were geometrically corrected to the Universal Transverse Mercator (UTM) Zone 32 North coordinate system on the same platform. To detect changes in the land use/cover at different years, post classification comparison of the change detection techniques was used.

The boundary data of Calabar metropolis was used to clip the Landsat imagery. The clipping helped to remove the extents outside the boundary of the satellite imagery (depicted in false colour in Figure 2 and 3). After clipping, Earth Resource Development Assessment System (ERDAS) imagine software was used for the pixel-based classification. The images were subjected to an unsupervised classification using the Iso Cluster (IC) and Maximum Likelihood Classification (MLC) spatial analyst tool. The IC combines the functionalities of the IC and the MLC, while the MLC performs classification on a set of raster bands using a signature file from the IC tool as the input for MLC. The ArcMap software was afterwards used for the final processing of the ERDAS outputs.

The classified raster output was converted to vectors (polygons) to allow for measurements to be done. The area coverage of each of the LULC classes was measured in sqkm, for each of the years under consideration, using the calculate geometry tool in the same ArcMap platform. A comparison of the land cover statistics assists in identifying the change in sqkm/percentage, trend, and rate of change in Calabar over the period. In this study, in line with Anderson, Hardy, Roach and Witmer (1976) the land-use/land-cover classification scheme in (Wakirwa, 2015), the various land-use/land-cover (LULC) types are modified and generalised into 4 classes within the study area as presented in Table 2.

Table 2: LULC classification scheme adopted

Serial No.	Land Use Category	Description
1	Built-up	Land used for residential and transportation/communication purposes (i.e., settlements and roads, high residential area, industry and administrative block).
3	Wet lands	Land covered characteristically saturated; a marsh.
4	Water body	Areas covered by body of water e.g., dam, lake, rivers and swamps
5	Green area	Areas that are spatially cultivated e.g. farmland, irrigation areas etc. as well as urban greenery, grasses, shrubs and grass-like plants.




Figure 2: Landsat imagery of Calabar in 2000

Figure 3: Landsat imagery of Calabar in 2018

Results and Discussions

Land use and land cover scenario in Calabar from 2000 to 2018

To assess the LULC situation of Calabar in the past and present, the LULC analysis was executed. Tables, maps, and charts were used to illustrate the LULC status quo in Calabar during 2000 and 2018 respectively. As symbolised by the legend, the colour brown is used to represent the built-up area, the light green for green area, the blue colour represents water bodies and the dark green, the forest areas.

As observed from the analysis outputs presented in Table 3, and Figures 4-5, there were evident changes in the LULC of Calabar during the 18-year period. From the year 2000 to 2018, there was little change in the extent of the water bodies, with only a slight decrease of 0.9 percent. The Calabar and Great Kwa rivers are tidal rivers, mostly influenced by their closeness to the Atlantic Ocean. The rivers are minimally influenced by urban activities, as the wet land bordering them still shield the rivers from direct urban extension impacts.

Within the same period under review, built-up areas increased by 16.1 sqm, with a coverage of 28.7 sqkm in the year 2000 and 44.8 in 2018. As pictured in Figure 5, the city visibly extended mostly to the northern fringes with new developments in the 8th mile, Ikot Nkebre, Adiabo, and other areas in the northern fringes. There was also no significant extension in the southern area, which houses the Calabar South Local Government Area (LGA). The direct impact of the urban built-up extension is on the green areas, which saw a reduction of 9.5 sqkm from an initial 62.2 sqkm in the year 2000. The wetlands on both sides of the city were also reduced by 5.2 sqkm. The extension in the urban build-up follows from the swelling population and increased socio-economic activities within the study area. Notably, the urban extension evident in these fringe areas of Calabar are haphazard and uncoordinated, thus requiring the attention of relevant agencies to ensure sustainable urban development in the area.

156

157 Table 3: LULC characteristics of Calabar in the year 2000 and 2018

	2000		2018			
Land use class	Area coverage (Sqkm)	Percentage	Area coverage (Sqkm)	Percentage	Area coverage (+/-)	Percentage (+/-)
Built up	28.7	17.5	44.8	27.4	16.1	9.9
Green areas	62.2	37.8	52.7	32	-9.5	-5.8
Wet lands	55.9	34	50.7	30.8	-5.2	-3.2
Water bodies	17.5	10.7	16.1	9.8	-1.4	-0.9
Total	164.3	100	164.3	100		

158




Figure 4: Output of LULC analysis in the year 2000




Figure 5: Output of LULC analysis in the year 2018

159

160

161

162

Conclusion and Recommendation

Typical to urban centers, urban extension in Calabar has been triggered by increased human activities, resulting in LULC changes. The impacts of these changes are worrisome in the face of uncoordinated planning policies and implementation. The application of remote sensing and GIS provided quantification, estimation and understanding of LULC changes in Calabar. There was an apparent extension in the urban built-up area of Calabar with the greenery mostly suffering the impacts of the extension.

From the findings of this study, the following is recommended:

- i. Land use planning should be instituted and implemented in Calabar to ensure that the usage of land, especially in the fringe, is sustainable. This effort should involve the sectoral integration of the relevant state ministries, boards, and bodies of the Calabar Municipal Council.
- ii. Intensive sensitisation should also be embarked upon and pursued holistically, with a view to ensuring that the inhabitants of the study area understand the negative effects and consequences that are associated with uncoordinated landuse development.
- iii. Deliberate efforts such as the declaration, reservation, and preservation of the ecologically fragile greenery and wetland areas of Calabar should be instituted and logically pursued by the government.
- iv. Finally, smart development of diverse-mixed land use should be encouraged, to reduce the rapid rate and large amount of urban land being converted for construction of homes, offices and commercial buildings.

References

- Abebe, A.G. (2013). Quantifying Urban Growth Pattern in Developing Countries using Remote Sensing and Spatial Metrics: A case Study in Kampala Uganda. Thesis submitted to the faculty of Geo-information science and earth observation, University of Twente. Pp 1-4
- Arnfield AJ (2003). Two decades of urban climate research: A review of turbulence, exchanges of energy and water, and the urban heat island. Int. J. Climatol., 23: 1-26.
- Atu, J.E., Offiong, R.A., Eni, D.I., Eja, E.I. and Esien, O.E. (2012). The Effects of Urban Sprawl on Peripheral Agricultural Lands in Calabar, Nigeria. International Review of Social Sciences and Humanities. 2(2):68-76.

198 Fasal S (2000). Urban expansion and loss of agricultural land- A GIS based study of
199 Saharanpur City, India. *Environ. Urbanisation*, 12(2): 133-149.

200 Lambin, E.F., Geist, H., and Lepers, E. (2003) Dynamics of landuse and land Cover Changes
201 in Tropical Regions. *Annual Review of Environment Resources*, 28: 205 -41 (online)
202 available http://www.globalrestorationnetwork.org/uploads/files/LiteratureAttachments/93_dynamics-of-landuse-and-land-cover-change-in-tropical-regions.pdf
203

204 Nanda T (2005). Urban Sprawl and occupational change in Raipur City. (Unpublished M.Sc.
205 Thesis, Department of Geography, University of New Delhi, Delhi.

206 Njoku, C. G., Okon, I., Itu, P-C. & Ahwen, V. O (2017). Population Growth and Pipe-Borne
207 Water Supply in Calabar Metropolis: Finding the Balance. *International Journal of Research
208 in Geography*. 3:3, PP 51-59.

209 Odjugo, P.A.O., Enaruvbe, G.O. and Isibor, H.O. (2015). Geospatial Approach to Spatio-
210 temporal Pattern of Urban Growth in Benin City, Nigeria. *African Journal of
211 Environmental Science and Technology*. 9(3): 166-175

212 Owoeye, J. O., & Ibitoye, O. A. (2016). Analysis of Akure urban land use change detection
213 from remote imagery perspective. *Urban Studies Research*, 2016. pp 9.
214 <http://dx.doi.org/10.1155/2016/4673019>
215

216 Offiong, R. A., & Eteng, O. E. (2014). Effect of urbanization on greenareas in Calabar
217 metropolis. *Int. J. Eng. Sci*, 3, 71-75.
218
219

220 United Nations Environment Programme (2002). Global Environment Outlook 3. Available
221 online at: <http://www.unep.org/GEO/geo3/index.htm>.

222 United Nations Population Fund (1999). The State of World Population, pp. 76. New York:
223 United Nations Publications.

224 Wakirwa, A. (2015). Analysis of Urban Sprawl Using Geospatial Techniques in Gombe
225 Metropolis, Gombe State, Nigeria. A Dissertation Submitted to the School of Postgraduate
226 Studies, Ahmadu Bello University, Zaria.

227 Wilkie, D.S. and Finn, J.T. (1996) Remote Sensing Imagery for Natural Resources
228 Monitoring. Columbia University Press, New York. p. 295.

229 Zhao LM, Dickson RE, Tian YH (2004). Evidence for a significant urbanization effect on
230 urban climate in china. *P Natl. Acad. Sci. USA*, 101: 9540- 9544.