

Original Research Article

Survey of Ethnomedicinal importance of plants used in the managements of diseases in Irepodun Ifelodun Local Government Area, Ado-Ekiti, Southwest, Nigeria.

*Olanipekun Mary K and Olowokere Enitan

Department of Plant Science and Biotechnology.

Ekiti State University Ado-Ekiti.

Corresponding Author

* Olanipekun Mary K, Department of Plant Science and Biotechnology, Ekiti State University
Ado-Ekiti.

e-mail:mary.olanipekun@eksu.edu.ng

Abstract The use of plants in treating diseases traditionally was found to be part of culture of the inhabitants in the study area. It was shown that the greater number of respondents (61, 50.8%) preferred the use of herbs when compared to the number who used synthetic (32, 26.6%) medicines. A total number of 50 botanicals belonging to 30 families were documented. Members of the family Fabaceae had the highest number (9) of plant species, followed by Euphorbiaceae with five (5) plant species, Cucurbitaceae (3 plants) and Rutaceae (3 plants) respectively. Leguminosae, Apocynaceae, Araceae, Anacardiaceae, Poaceae and Bignoniaceae had two species each, while other families had only one species each. *Azadiracta indica* and *Alstonia boonei* were high on the frequency of citation. These plants were mostly used for the treatment of malaria and body pains. The botanical names, common names, family names, forms of plant, plant part used and diseases treated were identified. Tree was found to be the most used plants followed by shrubs, herbs, underground stem, grass, climber, creeper and weed. Leaves were the plants parts frequently mentioned. It was revealed that 34 plants out of all the 50 plants were common. The methods of preparation and mode of administration revealed that plants were prepared traditionally with the combination of inert materials such as camphor, local gin or alcohol. Deforestation that lead to depletion of the plants and lack of government recognition were among the constraints to utilization of the traditional knowledge. Efforts on the development of cheaper medical services in the state should be more focused.

Key words: Respondents, herbal medicine, diseases, plants, citation, constrains

1.0 INTRODUCTION

Nigeria is endowed with diversity of plants and animals that are naturally used as foods, medicine, clothing and shelter. Traditional medicine is undoubtedly a reliable alternative approach to health delivery because it cheaper, easily accessible and effective as reported by kayode *et al.*, (2009). Although the use of plants traditionally as medicine has been discouraged in the time past by many medical practitioners because of the lack of clarity of chemicals

composition, dosages and toxicity level of plants used traditionally (Elujoba 2005., Lowel *et al.*, 2001; Adjanohoun *et al.*, 1991).

Medicinal plants include different types of plants such as *Alstonei boonei*, *Azadirachta indica*, *Mangifera indica*, *Anacadium occidentale*, *Zingiber officinale*, *Allium sativum*, *Allium cepa*, *Venona amygdalina*, *Momordica charanta*, *Chromoleana odorata*, etc were used in treating various health challenges because of the presence of different potent bioactive ingredients they contained (Olanipekun, *et al.*, 2016; Arowosegbe *et al.*, 2015; Olanipekun, and Kayode, 2014; kayode *et al.*, 2009). These medicinal plants are considered as rich resources of ingredients which can be used in **drug synthesis and development** for the treatment of various diseases (WHO, 2003). Moreover, some plants such as ginger, green tea, walnuts and some others plants as reported by (Tapsell *et al.*, 2006) are considered as important source of nutrition and as a result they are recommended for their therapeutic values (Adodo, 2004). Plants have not only **therapeutic values**, **they are important as food shelter, income**. Plants have the ability to synthesize a wide variety of chemical compounds that are useful in performing important biological functions Khan, et al., (2006). **The use of herbs to treat disease is a common phenomenon among non-industrialized or rural societies and is often more affordable than purchasing modern pharmaceuticals** (Akobundu and Agyakwa, 2003; Gbolade and Soremekun, 2000). The World Health Organization (WHO) in 2006 estimated that 80% of the population of some Asian and African countries presently used herbal medicine for some aspect of primary health care. **Though, studies** in the United States and Europe shown that the use of herbal drugs are less common in clinical settings, but has becoming increasingly more common in recent years as scientific evidences about the effectiveness of herbal medicine becoming more widely available (Akobundu and Agyakwa, 2003).

It was noted by Falconer *et al.*, (2002) the high rate of self treatment with **herbal drugs and pharmaceuticals are highly recognized recently**. The use of herbs involves the use of different parts of the plants, ranging from roots and root bark, bark and stems, latex and sap, leaves, buds and flowers, and seeds. For example, Falconer *et al.*, 2002, Diallo *et al.*, 1999 reported the use of leaves and stem bark of *Alstonia boonei* can be mashed and pound (in the treatment of rheumatism and swellings, powdered root and leaf sap of *Hoslundia opposita* as an antiseptic and jaundice or the stewed leaf and buds of mango to treat fevers or oil extracted from seeds. There is also a range of different modes of application and administration, some are ingested as decoction and infusions and tisanes, some applied direct as poultices or rubs and lotions, as enemas and eye drops, gargle, or nasal drops and as snuff (Falconer *et al.*, 2002, Diallo *et al.*, 1999; Coatney, 1963). Recently, there is a sort of paradigm shift from synthetic pharmaceutical to traditional medicine. This however, has lead to the reduction of plants population which serves as the primary source of plants. Many plants that are used and effective to treat diseases are no more available due to human influence and advancement in technology which have shifted the various subsistence farming to commercial farming, thereby leading to the genetic erosion of the plants population. Also, it was asserted by Momma (1994) and Olanipekun 2015 that there is an increase in the reduction in the availability of medicinal plants

species, because harvesting of plants is predatory and annihilative, this quite often leads to the destruction of the plants mother source. Similarly, medicinal plant resources are relevant, although scattered, eroded, wantonly destroyed and needed to be consolidated. Most indigenous plants that are found in the wild and semi-wild habitats are presently suffering from unprecedented genetic erosion due to large scale deforestation, fire and other several uses at which plants are put into by the people in the study area. Incidentally, the custodians of indigenous knowledge are now old and being reduced by death. It has been observed however, that some of these plants are not well documented and they are at the danger of extinction. Despite the medicinal value they possessed, a lot of them are being destroyed or removed and the land is used as commercial farming where mostly exotic products are planted.

In-lieu of the above, the objective of the study is to identified, document the medicinal plants used in treating diseases with the aim of identifying their abundance status.

MATERIALS AND METHODS

2.1 Study area

The study area was Igbemo in Irepodun/Ifelodun Local Government Area, Ekiti State, Nigeria. Irepodun/Ifelodun is one of the sixteen Local Government areas in Ekiti-State where Igbemo is one of the rural town in the local Government area. The major occupation of the inhabitants is farming, local trading and civil servants.

2.2 Experimental materials

The plant materials for the study were collected from the survey through the use of Semi-Structured Questionnaires through interview at the local markets in the study area. Recognizance surveys and visits were made in the Communities selected in order to intimate the residence of the intension to visit their communities for the study and to enlist their support, co-operation and active participation during the exercise and during the collection of samples of botanicals used in the communities.

Individual and group of three and four respondents were interviewed to ascertain the group consensus of the knowledge of the traditional use of plants. The plants were identified, collected and the information on the indigenous preparation and use of the plants were documented (Gbile, 2002). The voucher specimens of the identified plants were prepared and deposited at the herbarium unit of Plant Science and Biotechnology Department of Ekiti State University, Ado Ekiti. The scientific name, family name, parts used, abundance status, methods of preparation and mode of administration were documented. Similarly, the frequency of citation and informants consensus agreement which revealed the reliability and credibility on the utilization reports were also obtained. The data were spread on Excel sheet and encoded using Statistical Package for Socio Sciences (SPSS). Descriptive statistical tools (percentages, frequencies and mean) were used to summarize the data.

Frequency of citation (%) (F_c) was calculated by using the following methods provided by (Trotter and Logan, 1991) and (Heinrich *et al.*, 2009), formula:

$$(F_c) = \frac{\text{Number of informants who cited the species}}{\text{Total number of informants interviewed}} \times 100$$

Total number of informants interviewed

$$F_{ic} = \frac{N_{uR} - N_{TAXA}}{N_{uR} - 1}$$

F_{ic} = Factor of informant consensus N_{uR} = Number of used reports in a particular ailment

N_{TAXA} = Number of taxa used to treat that particular ailment

Results and discussion

Results: The use of plants in treating diseases traditionally was found to be part of culture of the inhabitants in the study area. It was shown that the greater number of respondents (61, 50.8%) preferred the use of herbs when compared to the number who used orthodox (32, 26.6%) medicines. Plants were observed to be readily available, less toxic, not resistance to diseases etc. However, a quiet number of 27(22.6%) respondents were of the opinion that the use of the combination of herbs and orthodox medicine provided a better, synergistic and fast healing than when only synthetic or herbal formulation is used (Table 1). Also, a total number of 50 botanicals belonging to 30 families were documented. Members of the family Fabaceae had the highest number (9) of plant species, followed by Euphorbiaceae which has five (5) plant species, Cucurbitaceae (3 plants) and Rutaceae (3 plants) respectively. However, Leguminosae, Apocynaceae, Araceae, Anacardiaceae, Poaceae and Bignoniaceae had two species each, while other families had only one species. *Azadiracta indica* and *Alstonia boonei* were frequently mentioned on the frequency of citation. These plants were mostly used for the treatment of malaria and pains by the respondents in the study area. The botanical names, common names, family names, forms of plant, plant part used and disease treated were presented in (Table 3). The plants mentioned were represented by various plants forms. Tree was found to be the most used plants followed by shrubs, herbs, underground stem, grass, climber, creeper and weed. The various plants parts mentioned include fruit, leaves, root, seed, stem bark and whole plants. Table 4 shows the availability of the identified plants in the study area. It was revealed that 34 plants out of all the 50 plants were common. These plants could be fetched within 20 to 30 minutes within the study area. The plants were cultivated sexually during the wet season of the year, thereby making the plants available. Similarly, it was shown in table 5, the methods of preparation and mode of administration of the identified plants used in treating diseases in the study area. The plants were mostly prepared traditionally with the combination of inert materials such as camphor, local gin or alcohol. Then, the concoction is taken as herbs, soup or use to bath. About 97% of the respondents complained of forest destruction, 87.2% complained of lack of

government recognition while 97.4% complained of concealment of knowledge by those who have it (Table 6). They only reveal the knowledge to their trusted children.

Table 1: List of treatment used in treating diseases by the respondents in the study area.

Type of treatment	No. of respondents	Reasons for the treatment
Use of herbs	61 (50.8%)	Readily available, cheaper, resistance to diseases
Orthodox	32 (26.6%)	Effective, easy access, no need of special preparation
Both methods	27(22.6%)	Synergy, effectiveness, non-resistance to diseases

Table 3: List of identified botanicals used in treating diseases in the study area.

S/N	Botanical Name/Forms of plants	Family name	Common Name	Part used	Disease treated	Frequency of citation %
1	<i>Abrus precatorius</i> (L)	Leguminosae	Egboirejeje	Leaf	rashes	0.8
2	<i>Acanthospermum hispidum</i> (DC)	Asteraceae	Dagunro	Leaf , bark, root	Pain	0.8
3	<i>Aframomum melegueta</i> (K. Schum), Leaf	Zingiberaceae	EgboAtare	Root	Rashes	2.3
4	<i>Allium sativum</i> (L), Leaf	Amaryllidaceae	Alubosaelewe	Leaf	Malaria	1.7
5	<i>Aleovera</i> (L)Burm.F. Leaf	Xanthorrhoeaceae	Aleovera	Leaf	Arthritis	4.4
6	<i>Alstonia boonei</i> (De wild), Tree	Apocynaceae	IgiAhun	Fruit	Yellow fever/ malaria	6.7
7	<i>Anchomanes difformis</i> (BI), Tree	Araceae	Ewe Abiorusoko	Leaf	Measles	1.7
8	<i>Anacardium occidentale</i> (L), Tree	Anacardiaceae	Cashew leaf	Leaf and bark	Malaria	4.2
9	<i>Azadirachta indica</i> (A.Juss), Tree	Meliaceae	Dogoyaro	Leaf	Malaria	8.3
10	<i>Bambusa vulgaris</i> (schrads. Ex	Poaceae	Ewe oparun	Leaf	Chicken pox	1.7

	Wendel), Tree					
11	<i>Boerhavia</i> <i>coccinea</i> (Mill)	Nyctaginace	EgboEtupa elila	Leaf and root	Liver problem	0.8
12	<i>Brachystegia</i> <i>eurycoma</i> (Harms), Tree	Fabaceae (Caesalpinoidea e)	EpoEku	Root	Malaria	1.7
13	<i>Buchholzia</i> <i>coriacea</i> (Engl.)	Capparaceae	Wonderful cola	Leaf	Typhoid	3.3
14	<i>Caesalpinia</i> <i>bondac</i> (L)Roxb	Fabaceae	Ewe ayo	Root and leaf	Chicken pox	1.7
15	<i>Cajanus cajan</i> (L)	Fabaceae	Feregede	Leaf	Measles	2.3
16	<i>Calliandra</i> <i>haematocephala</i> (Hassk)	Fabaceae	Tude	Leaf, root	Measle	0.8
17	<i>Citrullus lanatus</i> (thumb), leaf	Cucurbitaceae	Egunsibara	Leaf, seed, fruit	Gonorrhea	3.3
18	<i>Citrus</i> <i>aurantifolia</i> (Christ m).swingle	Rutaceae	osanwewe	Leaf	Tuberculosis	3.7
19	<i>Citrus aurantium</i> L.	Rutaceae	Osanganyi nganyin	Leaf	Tuberculosis	0.8
20	<i>Citrus paradise</i> (Macfad), Tree	Rutaceae	Grape	Fruit	Tuberculosis	2.5
21	<i>Croton</i> <i>Zambesicus</i> (Muell,Arg.)	Euphorbiaceae	Ewe Ajeobale	Leaf	Tetanus	1.7
22	<i>Elaeis guenensis</i> (Jacq)	Arecaceae	Ogbaranee yin	Seed, fruit	Headaches, Gonnorrhea	5.3
23	<i>Euphorbia</i> <i>Lateriflora</i> (schum&thonn); Herbs	Euphorbiaceae	Enuopire	Leaf	Viral disease	3.4
24	<i>Euphorbia hirta</i> (L.) Herbs	Euphorbiaceae	Ewe emile	Seed	Tetanus	0.8
25	Harungana <i>madagascariensis</i> (Lam ex poir), Tree	Cuttifereae	Arunje	Bark and leaf	Malaria	4.2
26	<i>Irvingia</i>	Irvingiaceae	Egbooro	Root	Rashes	0.8

	<i>gabonensis</i> (Aubry1-lecomte) Tree					
27	<i>Jatropha curcas</i> (L), Herb	Euphorbiaceae	Ewe lapalapa	Leaf	Malaria	1.2
28	<i>Kigelia Africana</i> (Lam),Benth. Tree	Bignoniaceae	Ewe pandoro	Leaf	Malaria	1.2
29	<i>Lagenaria</i> <i>breviflora</i> (benth. Roberty) Vine	Cucurbitaceae	Tangiri	Leaf	Measles	0.8
30	<i>Lawsonia inermis</i> (L), Shrubs	Lythraceae	Ewe laali	Leaf	Fever or high temperature	1.2
31	<i>Mangifera indica</i> (L), Trees	Anacardiaceae	Mango leaf	Leaf	Malaria	5.8
32	<i>Mormodica</i> <i>charantia</i> (L.) Herbs	Curcubitaceae	Ejinrin	Fruit	Gonorrhea/Pile	0.8
33	<i>Milicia excelsa</i> (welw), Tree	Moraceae	Ewe iroko	Leaf	Measles	0.8
34	<i>Morinda lucida</i> (Benth.) Trees	Moringaceae	Egbooruwo	Seed and root	Malaria	4.0
35	<i>Newbouldia leavis</i> (Seem), Tree	Rubiaceae	Ewe Akoko	Leaf, root, Bark	Arthritis, diarrhea and dysentery	0.8
36	<i>Parquetina</i> <i>nigrensens</i> (Afzel), Vine	Apocynaceae	Egbo ewe ogbo	Fruit	Rashes	2.5
37	<i>Phyllanthus</i> <i>muellerianus</i> (kuntze) Shrubs	Phyllanthaceae	Ewe egigun-eja	Leaf	Typhoid	0.8
38	<i>Piper guineense</i> (Schum.and Thonn.) vine	Piperaceae	EsoIyere	Seed	Tetanus	1.7
39	<i>Piptadeniastrum</i> <i>africanum</i> (Hook.f) Trees	Fabaceae	Agboyin	Leaf	Cough, headache and genitor- urinary infectious	0.8
40	<i>Plumbago</i> <i>zeylanica</i> (L) Herbs	Plumbaginaceae	Ewe inabiri	Root	Influenza flu	0.8

41	<i>Rauvolfia vomitoria</i> (Afzel). Shrubs	Apocynaceae	Ewe Asofeyeje	Leaf, root, bark,	Hypertension	0.8
42	<i>Senna siamea</i> (Lam) Trees	Fabaceae	Cashia leaf	Leaf	Malaria	3.3
43	<i>Spigelia anthelmia</i> (L), Shrubs	Loganiaceae	Paran-funfun	Root	Hepatitis	0.8
44	<i>Termnalia superba</i> Engl&Diels), Tree	Combretaceae	Epo igiafara	Root	Malaria	0.8
45	<i>Tetrapleura tetraptera</i> Tree (Schumach)	Fabaceae	Esoaidan	Leaf, root, bark	Tetanus	1.2
46	<i>Trema orientalis</i> (L), Tree	Cannabaceae	Egbo igiafefe	Fruit	Cough	0.8
47	<i>Uraria pizta</i> (Jacq), Shrubs	Fabaceae	Ewe apada	Leaf	Rashes	0.8
48	<i>Xylopia aethiopica</i> (Dunal) A.Rich. Trees	Annonaceae	Egboeruru	Root, fruit	Chicken pox	4.2
49	<i>Zea mays</i> (L.) Grass	Poaceae	Ewe agbado	Leaf, seed, husk	Chicken Pox	0.8
50	<i>Zingiber officinale</i> (Rosc), Herbs.	Zingiberaceae	Ginger	Seed	Arthritis	1.2

Table 4: The Occurrences of the Identified Plants in the Study Area.

S/N	Botanical name	Availability of the plant	Materials for propagation	Season
1	<i>Abrus precatorius</i>	Common	Seed	Wet
2	<i>Acanthospermum hispidum</i>	Common	Seed/cutting	Wet
3	<i>Aframomum melegueta</i>	Scarce	Seed	Wet
4	<i>Allium sativum</i>	Common	Seed	Dry
5	<i>Aloe vera</i>	Common	Stem cutting	Wet
6	<i>Alstonia boonei</i>	Common	Seed	Wet/dry
7	<i>Anchomanes difformis</i>	Common	Tuber	Wet
8	<i>Anacardium occidentale</i>	Common	Seed	Dry
9	<i>Azadirachta indiica</i>	Common	Seed	Wet
10	<i>Bambusa vulgaris</i>	Common	Sucker	Seed
11	<i>Boerhavia coccinea</i>	Scarce	Seed/root cutting	Dry
12	<i>Brachystegia eurycoma</i>	Scarce	Seed	Dry

13	<i>Buchholza coniacca</i>	Scarce	Bulb	Dry
14	<i>Caesalpinia bonnac</i>	Scarce	Seed	Wet
15	<i>Cajanus cajan</i>	Common	Seed	Dry
16	<i>Calliandra haematocephala</i>	Rare	Seed	Wet
17	<i>Citrullus lanatus</i>	Common	Seed	Wet
18	<i>Citrus aurantifolia</i>	Common	Seed	Dry
19	<i>citrus aurantium</i>	Common	Seed	Wet
20	<i>Citrus paradise</i>	Common	Seed/root cutting	Dry/wet
21	<i>Croton zambesicus</i>	Common	Seed	Wet
22	<i>Elaeis guenensis</i>	Common	Seed	Wet
23	<i>Euphorbia laterflora</i>	Common	Cutting	Wet
24	<i>Euphorbia hirta</i>	Common	Cutting	Wet
25	<i>Harungana madagascariensis</i>	Common	Seed	Dry/wet
26	<i>Irvingia gabonensis</i>	Common	Seed	Wet
27	<i>Jatropha curcas</i>	Common	Seed	Wet
28	<i>Kigelia Africana</i>	Common	Stem cutting	Wet
29	<i>Lawsonia inermis</i>	Scarce	Seed	Wet
30	<i>Magnifera indica</i>	Common	Seed	Wet
31	<i>Momondica charantia</i>	Common	Seed	Wet
32	<i>Milicia excelsia</i>	Common	Seed	Wet
33	<i>Morinda lucida</i>	Common	Root cutting	Wet
34	<i>Newbouldia leavis</i>	Common	Seed	Wet
35	<i>Laganaria breviflorus</i>	Common	Seed	Wet
36	<i>Parquetina nigrescens</i>	Scarce	Seed/root cutting	Wet
37	<i>Phyllanthus muellerianus</i>	Scarce	Seed	Wet
38	<i>Piper guineense</i>	Scarce	Seed	Dry
39	<i>Piptadeniastrum africanum</i>	Common	Seed	Dry
40	<i>Plumbago zeyianica</i>	Scarce	Seed/root cutting	Wet
41	<i>Rauvolfia vomitoria</i>	Common	Seed	Wet
42	<i>Senna siamea</i>	Common	Seed	Dry/wet
43	<i>Spigelia anthelmia</i>	Rare	Seed	Dry/wet
44	<i>Termnalia superbia</i>	Common	Vegetative/seed	Wet
45	<i>Tetrapleura tetraptera</i>	Common	Seed	Dry/wet
46	<i>Trema orientalis</i>	Common	Seed	Wet
47	<i>Uraria pizta</i>	Common	Seed/cutting	Wet
48	<i>Xylopia aethiopica</i>	Common	Seed	Wet
49	<i>Zea mays</i>	Common	Seed	Wet
50	<i>Zingiber officinale</i>	Common	Rhizomes	Wet

Table 5: Methods of preparation and mode of administration of the identified plants used in treating diseases in the study area.

Name of diseases	Medium of contact	Method of preparation and administration
Arthritis	Injury or surgical procedure.	1.Aloe plants, grind with Naphtalene (kafura) and mixed with black soap for bathing at morning and night. 2.Grind camphor, Naphtalene (kafura) and <i>Buchholza coriacea</i> (Wonderful kola) mixed together and pour it into bottle of gin. 1 teaspoon 3 times daily.
Chicken pox	Body contact Coughing and sneezing.	<i>Caesalpinia bonduc</i> leaf, <i>Bambusa vulgare</i> leaf, <i>Zea may</i> leaf <i>Xylopia aethiopica</i> leaf/cooked together. To be taken 3(three) times daily for 3 days.
Gonorrhea	Sexual contact	1. <i>Citrillus lanatus</i> , palmwine, <i>Momordica charantia</i> squeeze together and drink immediately. 2. <i>Citrillus lanatus</i> , peel it inside and soak for 2 days before drinking.
Hepatitis	Sexual contact Sharing of needle or sharp object	Egbo iyaoya. Wash and grind with <i>Piper guineense</i> seed, Full snail, <i>Boerhovia coccinea</i> , sun dry and grind then cook with bush meat and eat.
Influenza flu	Airborne e.g coughing and sneezing	<i>Acanthospermum hispidum</i> leaf, <i>Plumbago zeylanica</i> leaf, boil and drink 1 cup in the morning afternoon and night. Chewing of cashew stick.
Rashes	Body contact nasal secretion	Powder Efu ado (yellow colour) Etutu-sun dry Korofoeyinadie Grind it very well and start rubbing soap. Root and vine of <i>Parquetina nigrescens</i> with the root of <i>Abrus precatorius</i> root of <i>Chemopodium ambrosioides</i> (aruntata), root of <i>Irvingia gabonensis</i> , root of <i>Aframomum melegueta</i> , root of <i>Piper guineense</i> grind and mixed with adin apaku with black soap . For bathing morning and night.

Malaria	Insect	1. Root of khaya spp, Bark of <i>Terminalia superba</i> , Bark of <i>Brashystegia eurycoma</i> , Root of <i>Morinda lucida</i> . Wash and cut into pieces and cooked together.
Measles	Body contact	1. Rub <i>Anchomanes difformis</i> leaf, Fresh <i>Cajanus cajan</i> leaf, Efun ado (African chalk), Grind together with <i>Calliandra portoricensis</i> then boiled together and add Native Hot (ogogoro). For bathing Drink 2 spoonful Morning/night
Tetanus	Insect or bacteria	Seed of <i>Tetrapleura tetrapetra</i> , root of <i>Piper guineese</i> , root of <i>Aframomum melegueta</i> , <i>Croton zambesicus</i> , <i>Euphorbia hirta</i> , <i>Xylopia aethiopia</i> grind together and used it to wash.
Septic wounds	Insect or bacteria bites	<i>Parquetina nigrescens</i> (201) leaves, <i>Xylopia aethiopia</i> (9) leaves; Adin-Ifun Adie(Foul intestine) Male-akuko Female-Abo Adie.Grind all together and used peripe(foul feathers) to add to the wound.
Typhoid	Contaminated water E.G when typhi-bacteria get into water for drinking.	(1) 2. <i>Buchlozia coriacea</i> , 9 (nine) kolanut, cut into pieces inside a bottle add sevenup and soak for seven hours before drinking (2) Unripe <i>Carica papaya</i> , leaf of <i>Phyllanthus muellerianus</i> , bark of <i>Alstonia boonei</i> , leaf of <i>Anacardium occidentale</i> , leaf of <i>Azadirachta indica</i> , bark of <i>Mangifera indica</i> , cooked together and drink 3 (three) times daily for 3 days.
Tuberculosis	Airborne e.g coughing and sneezing	Leave of <i>Citrus aurantifolia</i> , Leave of <i>Citrus paradisiaca</i> , cooked together and add small potash and after cooking, leave it for 10 minutes before drinking, drink three times daily
Viral disease	Sexual contact	Peel <i>Citrullus lanatus</i> inside palm wine and soak for 12 hours. Drink 1 (one) cup per meal daily
Yellow fever	Insect bit	Bark of <i>Alstonia boonei</i> , bark of <i>Mangifera indica</i> , bark of <i>Harungana madagascenriensis</i> boiled together Drinking and bathing for 3 days.

Table 6: Constraints to Utilization of Ethno-botanical Knowledge

Methods	*Frequency	Percentage
Deforestation	110	96.9
Lack of Government recognition	98	87.2
Western education	76	77.5
Concealment of knowledge	113	97.4
Bush burning	91	84.1

Discussion

Plants have been a major source of medicine for human kind. The demand for traditional herbs is increasing very rapidly, mainly because of their effectiveness, cheaper price and disease resistance against orthodox drugs. The study area is endowed with an enormous diversity of plants and animals that are both domesticated and wild, living in different habitats and ecosystems. This heritage sustains the provision of food, medicine, clothing, shelter, spiritual, recreational, and other needs of her population. The information given on the local names of plants and the recipes used in the treatment of various diseases made the plants to be easily accessed for identification and collection. This was in consonance with Singh (2008) who reported that plants are generally known by their local names in every part of the world. Also, it was reported by (Jouad, *et al.*, 2001; Kayode *et al.*, 2009) that local names rendered a useful service as a means of plants identification especially in the rural areas, they may certainly be considered as a tool for search of new useful plants. **Though local names are not recommended as a substitutes to scientific names as they lack uniformity and consistency. Arowosegbe *et al.*, 2016; Olanipekun *et. al.*, 2016; Kayode *et al.*, 2009 had earlier reported that plants are identified and justified using their scientific names.** The use of plant parts such as leaves, roots and stem barks have be found to be efficient in the management of various diseases in the study area. The effectiveness of the use of the leaves and other plants parts in treating diseases such as Rashes, malaria, arthritis, measles tuberculosis and so on as shown in Table 1 revealed the presence of plants secondary metabolites. **Leaves are the site of photosynthesis where there is manufacture of food and accumulation of phytochemicals that are responsible for the various healing. This corroborated the assertion of researchers such as Olanipekun *et al.*, 2016; Ebong, *et al.*, 2015; Kayode *et al.*, 2015; Satrija *et al*; 1995; Sokomba *et al*; 1983; Dalziel, 1939; Iwu 1989; Aggarwal, (1995); Gbile (1991).** They reported the use of *Abrus precatorius*, *Aframomum melegueata*, *Parquetina nigrensecns* as plants for rashes, *Allium sativum*, *Alstonia boonei*, *Anacardium occidentale* are effective against malaria, *Aloevera*, *Citrus aurantifolia* for the treatment of tuberculosis among others etc.

The plants mentioned were represented by all plants forms. Trees were found to be the most used plants followed by shrubs, herbs, underground stem, grass, climber, creeper and weed. The respondents view on the increase in the demand of medicinal plants as herbal remedies for prevention and cure of diseases is because of the cost and effectiveness of plants as compared to the cost of orthodox drugs. Mostly, plants identified are propagated sexually, therefore planting of the plants is highly advocated for, in order to sustain the availability of the plants. This is in accordance to the reports of Kayode *et al.*, 2015; Olanipekun, 2014 who said there is tendency for the total eradication of medicinal plants if the rare ones are not afforested and conserved, as plants are facing a lots of pressures on their existence due to the increase in populations, advancements in science and technology, changing in farming system to commercial production among others. The methods of preparation varies between decoctions and infusions, where the preparation are in combination of other plants and some inert materials. The use of the combination of two or more plants traditionally are effective against various common and febrile illnesses among traditional healers. This exert synergy, thereby ensuring adequate treatment. This is in accordance with the reports of Erinoso and Aworinde, (2012); Oladunmoye and Kehinde, (2011); Etuk *et al.*, (2010); Tahraoui *et al.*, (2007); Ogbole *et al.*, (2010); Ajaiyeoba *et al.*, (2006) Osowole *et al.*, (2005), Lyoussi, (2007). Also, there were no scientific precautions such as the sterilization of the tools to prevent outbreak of diseases and the consideration of toxicity and concentration of the plants extracts used, hence the methods are purely traditional (Kayode *et al.*, 2015). Although, it was reported that there were rare occurrences of outbreak of diseases, because there are measures taken as an alternative for precautions such as dosages with the use of calibrated cups, teaspoons, proper washing of the plants materials before processing, with the full assurance that the measures are enough without any side effects. This is in accordance to the reports of Mustapha 2014., Ajaiyeoba, *et al.*, 2006; Osowole *et al.*, 2005 who reported that the methods of preparation of plants traditionally does not need a special measures and precautions during preparation as against the preparation of orthodox drugs. There are challenges where the respondents are not willing to give relevant information due to fear of losing their major source of their income, some demanded money prior to interview as they claimed to have “intellectual properties” stocked with knowledge of medicinal plants, while some castigated government for neglecting them and sending researchers to come and exploit their indigenous knowledge. The preferences of some aged men sharing the knowledge on a television programme rather than disseminating the information to researchers is a believe that this will rather and better advertising their names and services.

The constraint such as over exploitation of plants and conversion of some wild habitat to commercial farming are at the detriment to the availability of plants, hence serves as threat to the existence to the traditional knowledge (Usman, *et al.*, 2015). Also these resources are threatened by forest removal and bush burning, rapid socio-economic, technological and environmental changes as many youths are into western education and they preferred western medicine which they considered less stressful. Hence plants and the indigenous knowledge are in danger of extinction as older people die and younger generations fail to learn the traditional way of life.

Conclusion

This work provides additional information to the relevance of plants and their usefulness in the treatment of diseases in the study area. The traditional knowledge of plants have their benefits that cannot be eradicated among the users in the rural areas because plants are cheaper, available, disease resistance and effective. However, the dosage of the concoction, the toxicity and the concentration should be subjected to further studies.

REFERENCES

- Adodo, A. (2004) Nature Power, A Christian Approach to Herbal Medicine. *3rd Edition*. *Benedictine Publication Nigeria*. pp. 103–111.
- Adjanohoun, E., Ahiyi, M.R.A., Ake-Assi, L., Dramane, K., Eewude, J.A., Fadoju, S.O, Gbile Z.O, Goudote, E., Johnson, C.L.A., Keita, A., Morakinyo, O., Ojewole, J.A.O., Olatunji, A.O., Sofowora, E.A, (1991). Traditional Medicine and Pharmacopoeia: Contribution to ethnobotanical and floristic studies in western Nigeria. *Organization of African Unity/Scientific Technical and Research Commission*, 420pp,
- Aggarwal, A.(1995) “Indigenous and Scientific knowledge: Some critical comments.” *Indigenous Knowledge and Development Monitor* 2(3): 3-6.
- Ajaiyeoba, E.O., Ogbole O.O., Ogundipe, O.O. (2006). Ethnobotanical survey of plants used in the traditional management of viral infections in Ogun State of Nigeria. *European Journal of Scientific Research* 13(1): 64-73
- Akobundu, I.O. and C.W. Agyakwa, (2003). A Handbook of West African Weeds. *International Institute of Tropical Agriculture*, Oyo State, Ibadan, Nigeria.
- Anita, M. (2004). ‘Medicinal and aromatic plants: Monitoring the effectiveness of nbiological conservation. www.confound.org/global/global.html.
- Arowosegbe, S, Olanipekun, M.K, Kayode, J. (2015). Ethnobotanical survey of medicinal plants used for the treatment of Diabetes mellitus in Ekiti-State Senatorial districts, Nigeria. *European. Journal of Botany and Plant Science Phytology*. 2(4):1-8.
- Coatney, G.R. (1963). Pitfalls in a discovery: the chronicle of chloroquine. *American Journal*

- of Tropical Medicine* 12:121-128.
- Dalziel, J. M. (1937) Useful plants of West Tropical Africa. Crown Agents. London Pp. 612, 309, 384-386.
- Diallo D B, Hveem MA, Mahmoud G, Betge BS, Paulsen and A Maiga (1999). An ethnobotanical survey of herbal drugs of gourmadistrict, Mali. *Phamaceutical Biology* 37: 80-91
- Ebong, O. Nwauche, C, Ogbuehi I, Ifeyinwa N. Chijioke, N, Chinwe T. Umoh E, Aniekan Afia G, Pius Z (2015). Is this Evidence of Success in Malaria Prevention and Control Measures? *Greener Journal of Medical Sciences* 5 (1): 001-010.
- Elujoba, AA, Odaleye OM and Ogunyemi CM (2005) Traditional medicinal development for medical and dental primary health care delivery system in Africa. *Africa Journal of Traditional complementary and Alternative Medicine*. Vol. 3 Pp46.
- Erinoso, S.M. and Aworinde, D.O. (2012) Ethnobotanical Survey of Some Medicinal Plants Used in Traditional Health Care in Abeokuta Areas of Ogun State, Nigeria. *African Journal of Pharmacy and Pharmacology*, 6, 1352-1362.
- Etuk, E.U., Bello, S.O., Isezuo, S.A., Mohammed, B.J. (2010). Ethnobotanical survey of medicinal plants used for the treatment of diabetes mellitus in North-waestrern region of Nigeria. *Asian Journal of Experimental Biological Science* 1(1): 55-59
- Falconer, D.A., Moore, R.L., Gary, G.A. (2002) Correlation of the Coronal Mass Ejection Productivity of Solar Active Regions with Measures of Their Global Nonpotentiality from Vector Magnetograms: Baseline Results. *The Astrophysical Journal*, Volume 569, Issue 2, pp. 1016-1025.
- Gbile, Z. O. (1991) Identification and Pharmaceutical Importance of medicinal plants. Ibadan: Forestry Research Institute of Nigeria, Unpublished paper.
- Gbile, Z.O. (2002) Vernacular Names of Nigeria Plants (Yoruba). Forestry Research Institute

- of Nigeria (FRIN), Ibadan, 124. Nigeria. *Tropical Journal of Pharmaceutical Research*, 9, 110-118.
- Gbolade and Soremekun, A.A, (2000). Medicinal and pesticidal potential of Nigerian plants. Proceedings of International Symposium on Natural Products in Developing Countries: Trends and Needs.
- Heinrich M, Moerman, ES and Leonti M (2009) Ethnopharmacological field studies; A critical assessment of their conceptual basis and methods. *J ethnopharmacological*, 124;1-17.
- Iwu, M. (1993) Food for medicine in : M. Iwu (Ed) Dietary plants and Masticetories as sources of biologically active substances, University of Ife Press, Pp.303-310.
- Jouad, H., Haloui, M., Rhiouani, H., EL Hilaly, J., Eddouks, M. (2001). Ethnobotanical survey of medicinal plants used for the treatment of diabetes, cardiac and renal diseases in North-central region of Morocco (Fez-Boulemane). *Journal of Ethnopharmacology* 77(2-3): 175-182
- Kayode, J, Michael, A.O, Modupe, J.A., Ayodele, A.O (2015). Stem Barks and Roots Extravitism in Ekiti State Nigeria: Need for Conservation as a Sustainable Innovation in Healthcare Management in Rural Areas. *American Journal of Bioscience*. 3(2):28-33.
- Kayode, J., Olanipekun, M.K, Tedela, P.O (2009). Medicobotanicals studies in relation to veterinary in Ekiti-State, Nigeria Conservation of botanicals used for the treatment of poultry diseases. *Ethnobotany of Leaf*. 13:273-280.
- Khan, N.A, Rashid, A.Z.M., (2006). A study on the indigenous medicinal plants and healing practices in Chittagong hill tracts (Bangladesh). *African Journal of Traditional Medicine*. 2: 37 – 47.
- Mustafa, A.A., Fawibe, O.O., Ajiboye, A.A., Agboola, D.A. (2014). Ethnobotanical survey of medicinal plants used in the treatment of diabetes in Irepodun Local Government area of Osun State, Nigeria. *Greener Journal of Biological Sciences*. 4(2): 59-68

- Ogbole O, Abolade A, Ajayeoba E (2010). Ethnobotanical survey of plants used in treatment of inflammatory diseases in Ogun State of Nigeria. *European Journal of Scientific Research* (3)79-124
- Ogbole, O.O. and Ajaiyeoba, E.O. (2010). The Practice and Ethnobotanical Survey. *African Journal of Traditional Medicine, Complementary and Alternative Medicines*,
- Oladunmoye, M.K. and Kehinde, F.Y. (2011) Ethnobotanical Survey of Medicinal Plants Used in Treating Viral Infections among Yoruba Tribe of South Western Nigeria. *African Journal of Microbiology Research*, 5, 2991-3004.
- Olanipekun, M.K., Kayode, J. (2014). Ethno-veterinary studies on botanicals used for treating carnivorous in Ekiti State, Nigeria. *Ind. Journal of Science and Technology* 2(3): 98-104
- Osowole, O., Ajaiyeoba, E., Bolaji, O, Akinboye, D., Fawole, O., Gbotosho, G., Ogbole, O, Ashidi, J., Abiodun, O., Falade, C., Sama, W., Oladepo, O., Itiola, O., Oduola, A., (2005). A survey of treatment practices for febrile illnesses among Traditional Healers in the Nigerian (Middle Belt Zone). *African Journal of Traditional Medicine* 2, 337 – 344.
- Satrija F, Nanseen, P. Murtini, S. and HESI (1995) Anthelmintic activity of papaya latex against *Heligmosmoides polygyrus* infections in mice. *Journal of Ethnopharmacology* 48:161-164.
- Singh, M.P., Malla, S.B., Manadkar, A (1980). Medicinal plants of Nepal. *Ecology and Botany* 33 (2): 185-193
- Singh, H. (2008) Importance of Local Names of Some Useful Plants in Ethnobotanical Study. *Indian Journal of Traditional Knowledge*, 7, 365-370.
- Tahraoui, A., El-Hilaly, J., Istraili, Z.H., Lyoussi, B. (2007). Ethnopharmacological survey of plants used in the traditional treatment of hypertension and diabetes in South-eastern Morocco (Errachidia province). *Journal of Ethnopharmacology* 110: 105-117.

- Tapsell, L.C., I. Hemphill, L. Cobiack, C.S. Patch and Sullivan D.R (2006). Health benefits of herbs and spices: The past, the present, the future. *Medical Journal of Australia*, 185: S4-S24.
- Trotter R and Logan M (1991). Informant consensus; a new approach for identifying potentially effective medicinal plants and diet: biobehavioural approaches. Edited by Etkin N.L. Bedford hills, New York: Redgrave Publishers;1986: 91- 112.
- Usman, I. S., A. U. Mani and I. D. Mohammed (2015). Indigenous Foot and Mouth Disease Control Methods among Nomadic Cattle Fulanis in Adamawa State, Nigeria. *Alexandria Journal of Veterinary Science*. 45:71-78.
- World Health Organisation. (1985). Diabetes mellitus report of WHO study group on diabetes mellitus. Geneva: *WHO Technical Report*. pp 727.
- WHO (2006). Traditional medicine – growing needs and potential. WHO Policy Perspectives Med. 2:1–6.