

2 **EFFECTS OF *CAPSICUM FRUTESCENS* SUPPLEMENTED DIET**  
3 **(C.F.S.D) ON FASTING BLOOD GLUCOSE LEVEL AND BIOCHEMICAL**  
4 **PARAMETERS IN ALLOXAN INDUCED DIABETIC WISTAR RATS.**

5 Ojeh, E. Anthony,<sup>1</sup> Adegbor, C. Ese,<sup>2</sup> Ewhre O. Lawrence<sup>3</sup>

6 <sup>1,2</sup>Department of Physiology, Faculty of Basic Medical Sciences,  
7 Delta State University, Abraka, Nigeria.

8 <sup>3</sup>Department of Pharmacology and Therapeutics, Faculty of Basic  
9 Medical Sciences, Delta State University, Abraka, Nigeria.

---

10  
11  
12  
13 **ABSTRACT**  
14

**Aim of the study:** Assessment of the effects of *Capsicum frutescens* supplemented diet (C.F.S.D) on biochemical parameters in alloxan induced diabetic Wistar rats.

**Experimental Design:** Forty male Wistar rats weighing between 130 to 150g were divided into four groups. Group 1 served as a normal control and received normal feed. Group 2 (Diabetic control) received normal feed. Group 3 (Diabetic test 1) received normal feed + 1% C.F. Group 4 (Diabetic test 2) received normal feed + 2% C.F.

**Place and Duration of study:** This study was carried out in the department of Physiology, Faculty of Basic Medical Sciences, Delta State University, Abraka and the feeding lasted for three weeks. At the end of the experiments, the animals were sacrificed, blood samples were collected and the serum was further subjected to biochemical analysis using biochemical analyzer (Reflotron Plus). Indexes investigated include; AST, ALT, ALP, GGT, Creatinine, Uric acid, total cholesterol, high density lipoprotein cholesterol (HDL-c) and fasting blood sugar level.

**Results:** Serum AST, ALT, ALP, GGT, Creatinine, Uric acid, total cholesterol and fasting blood sugar level were increased while serum high density lipoprotein cholesterol (HDL-c) was decreased in diabetic control (group 2), when compared with normal control (group 1). The incorporation of *Capsicum frutescens* in the diet at 1% and 2 % doses significantly ( $P < 0.05$ ) reduced the fasting blood glucose level as well as the serum level of AST, ALT, ALP, GGT, Creatinine, Uric acid, total cholesterol when compared with diabetic control. Serum HDL was also significantly increased when compared with diabetic control (Table 1). Decrease in body weight in diabetic control group and increased in body weight of 1% and 2% C.F.S.D groups were also observed (Table 2).

**Conclusion:** The observed improvement in the biochemical parameters of alloxan induced

oxidative stressed Wistar rats by 1% and 2% *Capsicum frutescens* supplemented diet suggests *Capsicum frutescens* to possess, cardio-protective and anti-diabetic properties. This could be attributed to its Phytochemical constituents.

**Recommendation:** The incorporation of *Capsicum frutescens* in the diet of patients susceptible to oxidative imbalance such as diabetes mellitus is worthy of recommendation.

15

16 **E-mail:** [tonniojie@yahoo.com](mailto:tonniojie@yahoo.com), [tonniojie@gmail.com](mailto:tonniojie@gmail.com)

17 **Tel:** +234-803700697

18 **Postal address:** P.O Box 91, Abraka, Delta State, Nigeria.

19

20 **Keywords:** Capsicum Frutescens, Fasting Blood Glucose, Liver  
21 enzymes, Capsaicin, Thermogenesis.

22

23

## 24 1. INTRODUCTION

25

26 Diabetes mellitus (DM) has been described as a multifactorial disease that is characterized  
27 by hyperglycemia and lipoprotein disorders [1], increased basal metabolic rate [2], defect in  
28 reactive oxygen species scavenging enzymes, as well as altered intermediary metabolism of  
29 major food substances [2]. Diabetes is a major degenerative disease in the world today [3],  
30 affecting at least 15 million people and having complications which include hypertension,  
31 atherosclerosis and microcirculatory disorders.

32 At least 80% of Africans rely on plant medicine for their healthcare [4]. Today, medicinal  
33 plants are increasingly being used in most parts of the world as: hypolipidemic [5];  
34 antihypertensive [6]; treatment for skin diseases [7] and hypoglycemic [8].

35 For the past 25 years, epidemiological studies have revealed a diminished risk of chronic  
36 diseases in populations consuming diets fortified with fruits and vegetables, [9]. It has been  
37 suggested that antioxidants found in large quantities in fruits and vegetables may be  
38 responsible for this protective effect, [10]. In the past three decades, it has been  
39 experimentally documented that several common spices can also exert health beneficial  
40 physiological effects, [11; 12]. These physiological effects of spices in most instances have  
41 been traced to the bioactive chemicals in them. Among these physiological effects of spices  
42 documented are hypolipidemic and antioxidant properties with beneficial health implications,  
43 [13].

44 One of such phytomedicine is *Capsicum frutescens*, a short lived evergreen shrub that  
45 usually grows from 1 to 1.5m in height and 1 to 3cm in basal stem diameter. It is commonly  
46 recognized by its fruit, the large red, orange, or yellow chili peppers that the plant produces.

\* Tel.: +xx xx 265xxxxx; fax: +xx aa 462xxxxx.

E-mail address: xyz@abc.com.

47 *Capsicum frutescens* fruits grow as long pods, and when ripe they develop their  
48 characteristic warm coloring. Its species likely originated in south or Central America. It  
49 spread quickly throughout the subtropical regions in the area and still grows wild today. The  
50 plant grows in tropical climates, because it needs a warm, humid climate to survive. It had  
51 been reportedly used in the treatment of various ailments such as Diabetes, Blood pressure  
52 [high/ low], Bronchitis, Burning feet, Arthritis, among others, [14].

53 A number of studies have shown multiple pharmacological effects of *Capsicum* on a variety  
54 of physiological systems such as cardiovascular system, gastro-intestinal tract, metabolic  
55 rate, and pain relief, [15].

56 Previous research had shown the Chemo-Protective effect of spices among which are;  
57 *Turmeric, Capsicum frutescens, Cloves* and *Cardamom* on Correcting Iron Overload-  
58 Induced Liver Injury, Oxidative Stress and Serum Lipid Profile in Rat Model. The  
59 incorporation of chili (*Capsicum frutescens*) in the diet at 2 % significantly restored the  
60 enzyme activities of the liver AST, ALT, and ALP to normal level. The mean values of lipid  
61 profile, the MDA and serum total bilirubin were also reduced, [16].

62 The active substance in *Capsicum frutescens* that gives the hot and spicy flavor was  
63 identified as capsaicin, [15]. Red chili (RC) (*Capsicum frutescens*) is widely used as a spice  
64 for flavoring foods, particularly in South- East Asian and Latin-American countries. Several  
65 studies indicate that capsaicin (red pepper) is an appetite suppressant that can slightly  
66 increase metabolism. Spicing up one's foods with capsaicin-containing spices and using red  
67 pepper as a condiment can aid in increasing the rate of fat burning or thermogenesis. In an  
68 article published in the British Journal of Nutrition, Yoshioka et al (2001)<sup>17</sup> concluded that the  
69 consumption of red pepper and caffeine can induce a considerable change in energy  
70 balance when individuals are given free access to foods. Pungent capsaicinoids (capsaicin,  
71 dihydrocapsaicin), antioxidant vitamins (ascorbic acid, vitamin E), carotenoids ( $\beta$ -carotene,  
72  $\beta$ - cryptoxanthine) and several organic acids and minerals are the major active ingredients of  
73 *Capsicum frutescens*, [18]. Capsaicin (8-methyl-*N*-vanillyl-6-nonamide) is an irritant for  
74 mammals, including humans, and produces a sensation of burning in any tissue with which it  
75 comes into contact. Capsaicin and several related compounds are called capsaicinoids and  
76 are produced as a secondary metabolite probably as deterrents against certain herbivores  
77 and fungi. The burning and painful sensations associated with capsaicin result from its  
78 chemical interaction with sensory neurons. Capsaicin, as a member of the vanilloid family,  
79 binds to a receptor called the vanilloid receptor subtype 1 (VR1), [19].

80 Diabetes mellitus which arise as a result of insulin insufficiency is associated with altered  
81 activity of various biochemical parameters such as alkaline phosphatase (ALP), alanine  
82 transaminase (ALT), aspartate transaminase (AST), serum electrolyte, lipid profile, among  
83 other biochemical parameters, [20; 21].

84 Because the liver plays a critical role in the maintenance of carbohydrate  
85 homeostasis, glucoregulation, and insulin degradation, it is not surprising that its functions  
86 may be affected as a result of diabetes mellitus.

87 However, scientific information on the effects of *Capsicum frutescens* supplemented  
88 diet on biochemical parameters of alloxan induced diabetic Wistar rats is lacking. It is against  
89 this background that this study was designed.

90

## 91 **2. MATERIAL AND METHODS**

92

### 93 **Chemicals and equipments:**

94 All chemical used in the research were procured as follows:

95 Red Chili (*Capsicum frutescens*), purchased from Abraka market in Ethiopia East local  
96 government area, Delta State. Alloxan monohydrate (Sigma, alpha Aesar, 25g. A15324,  
97 CAS:2244-11-3. Cotton wool, Hand gloves, Dissecting kit, Centrifuge, Pipettes, Growers  
98 mash ,Beakers, Electronic weighing balance, Syringes and needles, Marker pen, Oncall  
99 Redii Glucometer and Reflorton plus<sup>(R)</sup> reflectance photometer (Roch Diagnostic GmbH, D-  
100 68298).

### 101 **COLLECTION AND IDENTIFICATION OF *CAPSICUM FRUTESCENS***

102 The plant *Capsicum frutescens* fruit was purchased from Abraka market in Ethiopia East  
103 Local Government of Delta sate were most people usually get it from and was authenticated  
104 by Dr. (Mrs). N.E. Edema in the department of Botany, Faculty of Science, Delta State  
105 University, Abraka. It was then air-dried at room temperature ( $22\pm 1^{\circ}\text{C}$ ) for 14 days until a  
106 constant weight was attained and was then blended with the aid of a grinding machine and  
107 stored in an airtight container for use in the experiment.

108



109



110

111

112

**FRESH AND DRIED *CAPISCUM FRUTESCENS* FRUITS**

\* Tel.: +xx xx 265xxxxx; fax: +xx aa 462xxxxx.  
E-mail address: xyz@abc.com.

113

114 **PREPARATION OF PEPPER SUPPLEMENTED DIET**

115 1% and 2% *Capsicum frutescence* supplemented diet were prepared weighing 1g and 2g  
116 of powdered *Capsicum frutescence* and mixing them with 99g and 98g of animal feed  
117 (growers mash) respectively.

118 **COMPOSITION OF THE GROWERS MARSH**

119 Protein-19.0%

120 Fat -2.85%

121 Fibre – 6.00%

122 Calcium – 1.00%

123 Available phosphate – 0.45%

124 Energy – 2875 KGC

125 (Animal Care Services Konsult (NIG) LTD).

126

127 **HANDLING OF EXPERIMENTAL ANIMALS**

128 Forty (40) Male Wister rats weighing 130-150g were procured from the International institute  
129 of tropical agriculture, (IITA), Ibadan Nigeria. They were acclimatized for 14-days at in the  
130 animal house unit in the Department of Pharmacology, Faculty of Basic Medical Science,  
131 Delta State University Abraka before commencement of the experiment. The rats were kept  
132 in well ventilated wooden cages. They were exposed to 12 hours of natural daylight and  
133 darkness and fed standard rat feed and water *ad libitum*. Procedures followed in raising the  
134 experimental animals were in accordance with the ethical standards of the Institutional  
135 Animals Ethics Committee (IAEC). And permission for the use of animals and animal  
136 protocol was obtained from the Research Ethics Committee of Delta State University,  
137 Abraka.

138 **Induction of diabetes**

139 Thirty (30) animals were food deprived for 24hours (but with free access to water) and later  
140 rendered diabetic by a single intraperitoneal dose of alloxan monohydrate (150mg/kg)  
141 prepared in stock of 1500mg/50ml and a concentration of 30mg/ml. Three days after  
142 induction of diabetes, rats with fasting blood glucose concentration above 200mg/dl were  
143 confirmed diabetic and were randomly selected for the study. Diabetic state was maintained  
144 for three days for well establishment of diabetes.

145 **EXPERIMENTAL PROCEDURE**

146 Rats with evidence of diabetes mellitus were randomized into different groups alongside with  
147 non diabetic rats as follows;

148 **Group 1:** Non diabetic rats received normal diet (normal control)

149 **Group 2:** Diabetic rats received normal diet (diabetic control)  
 150 **Group 3:** Diabetic rats received 1% *Capsicum frutescens* supplemented diet (test 1 group)  
 151 **Group 4:** diabetic rats received 2% *Capsicum frutescens* supplemented diet (test 2 group).

152

153 Animal feed was formulated with 1% and 2% *Capsicum frutescens* and treatment was done  
 154 twice daily for twenty one days. Rats' initial body weight prior to commencement of treatment  
 155 was recorded. Inclusion criteria in this study were; non diabetic that were not induced with  
 156 diabetes (which served as positive control), and animals with evidence of diabetes.  
 157 Exclusion criteria include those animals that died during the maintenance of diabetes. Thus  
 158 higher numbers of animals were allocated to groups 1, 2 and 3.

159 **BLOOD COLLECTION AND BIOCHEMICAL ASSAY**

160 After twenty one days of treatment, all overnight fasted rats were anaesthetized using  
 161 chloroform and then sacrificed. Blood samples collected by cardiac puncture were delivered  
 162 into lithium heparin bottles. The tubes were then centrifuged at 4000rpm for ten minutes to  
 163 obtain clear serum which were later subjected to biochemical evaluation for ALT, AST, ALP,  
 164 GGT, URIC ACID, CREATININE, HDL, and TOTAL CHOLESTEROL using Reflotron plus  
 165 kit.

166 Fasting blood glucose level was determined with the aid of glucose analyzer machine  
 167 (Oncall- Redii glucometer) by collecting blood samples from tail veins of overnight fasted  
 168 animals. Values were expressed in mg/dl.

169 **STATISTICAL ANALYSIS**

170 The result of this study were expressed as mean  $\pm$  SEM, and were analyzed by one way  
 171 analyses of variance (ANOVA) using statistical package for social science (SPSS, 16).  
 172 Difference between the means were tested with post Hoc- Tukey's test for multiple  
 173 comparison and significance was considered when  $p < 0.05$ . Student's dependent t-test was  
 174 used to analyze the significant difference between body weight before treatment and after  
 175 treatment.

176

177 **3. RESULTS AND DISCUSSION**

178

179 **Table 1:**

180 **Effects of *Capsicum frutescens* supplemented diet on biochemical parameters of**  
 181 **alloxan induced diabetic Wistar.**

	<b>Group 1: Normal control</b>	<b>Group 2: Diabetic control</b>	<b>Group 3: Diabetic +1% C.F.S.D</b>	<b>Group 4: Diabetic + 2% C.F.S.D.</b>
<b>Creatinine (IU/L)</b>	0.42 $\pm$ 0.03	0.94 $\pm$ 0.17 <sup>a</sup>	0.4 $\pm$ 0.3 <sup>b</sup>	0.54 $\pm$ 0.07 <sup>b</sup>

\* Tel.: +xx xx 265xxxxx; fax: +xx aa 462xxxxx.  
 E-mail address: xyz@abc.com.

<b>Uric acid (IU/L)</b>		5.49 ± 0.2	7.87 ± 0.85 <sup>a</sup>	5.03 ± 0.2 <sup>b</sup>	6.3 ± 0.7
<b>GGT (IU/L)</b>		223.4 ± 7.5	275.0 ± 10.7 <sup>a</sup>	221.8 ± 6.4 <sup>b</sup>	224.8 ± 6.0 <sup>b</sup>
<b>AST (IU/L)</b>		278.4 ± 19.6	325.2 ± 26.1	247.2 ± 10.8 <sup>b</sup>	251.8 ± 12.3
<b>ALP (IU/L)</b>		251 ± 6.81*	316.4 ± 37.7*	327.6 ± 27.6*	243.8 ± 4.53*
<b>ALT (IU/L)</b>		61.7 ± 1.03*	128.2 ± 32.97*	98.98 ± 8.74*	87.86 ± 8.54*
<b>HDL (mg/dl)</b>		47.98 ± 1.8 <sup>ns</sup>	43.1 ± 2.8 <sup>ns</sup>	46.8 ± 1.6 <sup>ns</sup>	46.0 ± 1.4 <sup>ns</sup>
<b>T. Cholesterol (mg/dl)</b>		85.6 ± 5.6	79.2 ± 4.4	101.6 ± 3.3 <sup>b</sup>	61.5 ± 3.4 <sup>abc</sup>
<b>Initial Blood glucose level (mg/dl)</b>		88.8 ± 6.22	380.2 ± 16.6	363.8 ± 24.3 <sup>d</sup>	382.2 ± 14.7 <sup>d</sup>
<b>Final Blood glucose level (mg/dl)</b>		94.8 ± 6.18	370.0 ± 19.81 <sup>a</sup>	182.8 ± 16.82 <sup>abd</sup>	146.6 ± 14.8 <sup>bd</sup>
		(6.8%)	(-2.63%)	(-49.8%)	(-61.6%)

182 **Values are expressed as mean ± Standard error of mean (S.E.M), n=10 \*P<0.05:**  
183 **Significant as determined by one way analysis of variance. Significant difference (<sup>abc</sup>P**  
184 **< 0.05): (a) compared to group 1, (b): to group 2, (c): to group 3.<sup>d</sup>P<0.05: Significant**  
185 **when initial and final fasting blood glucose level were compared in groups 3 and 4.**  
186 **Values in parenthesis depict the percentage change in FBGL when initial and final**  
187 **values were compared.**

188

189 **AST- (Aspartate Transaminase)**

190 **ALT- (Alanine amino Transaminase)**

191 **ALP- (Alkaline Phosphatase)**

192 **GGT- (Gamma Glutamyl Transpeptidase)**

193 **They are all liver enzymes(biomarkers) of liver damage.**

194

195

196 **Table 2:**

197 **Effects of Capsicum frutescens (C.F.) supplemented diet on body weight of alloxan**  
198 **induced diabetic rats.**

	<b>Body weight before treatment Week 0 (g)</b>	<b>Body weight after treatment Week 3 (g)</b>
<b>Group 1 (Normal control)</b>	131 ± 9.8	195 ± 17.2



		(48.9%)
<b>Group 2 (Diabetic control)</b>	140 ± 9.6	120 ± 7.9 (-16.7%)
<b>Group 3 (Diabetic, 1% C.F.S.D)</b>	125 ± 6.7	134 ± 19.2 (7.2%)
<b>Group 4 (Diabetic, 2% C.F.S.D)</b>	140 ± 7.2	152 ± 16.9 (8.5%)

199 **Values are expressed as mean ± Standard error of mean (SEM), n = five animals per**  
200 **group. C.F: *Capsicum frutescens*.**

201

202 Table 1 above depicts the effects of *Capsicum frutescens* on biochemical  
203 parameters of alloxan induced diabetic Wistar rats.

204 From the result of serum creatinine, group 2 ( $0.94 \pm 0.17$ ) significantly increased  
205 serum creatinine level when compared with group 1 ( $0.42 \pm 0.03$ ). Group 3 ( $0.40 \pm 0.3$ ) and  
206 group 4 ( $0.54 \pm 0.07$ ) significantly reduced ( $P < 0.05$ ) serum creatinine level when compared  
207 with group 2.

208 From the result of serum uric acid, group 2 ( $7.87 \pm 0.85$ ) significantly increased  
209 serum uric acid level when compared with group 1 ( $5.49 \pm 0.2$ ). Group 3 ( $5.03 \pm 0.2$ )  
210 significantly reduced serum uric acid level when compared with group 2. Group 4 ( $6.3 \pm 0.7$ )  
211 reduced serum creatinine level when compared with group 1, but did not attain statistical  
212 significance ( $P > 0.05$ ).

213 From the result of serum gamma glutamyl transferase (GGT), group 2 ( $275.0 \pm 10.7$ )  
214 significantly ( $P < 0.05$ ) increased serum GGT level when compared with group 1 ( $223.4 \pm$   
215  $7.5$ ). Groups 3 ( $221.8 \pm 6.4$ ) and 4 ( $224.8 \pm 6.0$ ) significantly ( $P < 0.05$ ) reduced serum GGT  
216 level when compared with group 2.

217 From the result of serum aspartate transaminase (AST) group 2 ( $325.2 \pm 26.1$ )  
218 increased serum AST level but did not attain statistical significant ( $P > 0.05$ ) when compared  
219 with group 1 ( $278.4 \pm 19.6$ ). However, group 3 ( $247.2 \pm 10.8$ ) significantly ( $P = 0.030$ ) reduced  
220 serum AST level when compared with group 2. Group 4 ( $251.8 \pm 12.3$ ) reduced the serum  
221 level of AST when compared with group 2, but was not statistically significant ( $P > 0.05$ ).

222 From the result of serum alkaline phosphatase (ALP), there was an overall  
223 significant difference ( $P < 0.05$ ) as determined by one way ANOVA. However Turkey's post  
224 hoc test did not reveal any significant difference between groups. However, there was an  
225 increase in group 2 ( $316.4 \pm 37.7$ ) serum ALP level when compared to other groups. Group  
226 1 ( $327.6 \pm 27.6$ ) was increased among other groups while group 4 ( $243.8 \pm 4.53$ ) reduced its  
227 level and group 3 ( $327.6 \pm 27.6$ ) increased its level its level.

228 From the result of serum alanine transaminase (ALT), a significant difference was  
229 observed as determined by one way ANOVA. Turkey's post hoc test did not reveal any  
230 statistical significant. However, serum ALT mean value was highest in group 2 ( $128.2 \pm$   
231  $32.97$ ) followed by group 3 ( $98.98 \pm 8.74$ ), next to group 4 ( $87.86 \pm 8.54$ ) and least in group 1  
232 ( $61.7 \pm 1.03$ ).

233 From the result of serum high density lipoprotein cholesterol (HDL) there was no  
234 significant difference as determined by one way analyses of variance (ANOVA), ( $P > 0.05$ ).  
235 However, serum HDL level was highest in group 1 ( $47.98 \pm 1.8$ ) followed by group 3 ( $46.8 \pm$   
236  $1.6$ ) next to group 4 ( $46.0 \pm 1.4$ ) and least in group 2 ( $43.1 \pm 2.8$ ).

237 From the result of serum total cholesterol level, there was a significant difference as  
238 determined by one way ANOVA. Post hoc Turkey's test showed that the group 3 ( $101.6 \pm$   
239  $3.3$ ) significantly increased serum total cholesterol level when compared with group 2 ( $79.2 \pm$   
240  $4.4$ ). Group 4 ( $61.5 \pm 3.4$ ) significantly ( $P < 0.05$ ) reduced serum total cholesterol level when  
241 compared with group 2 and 3 ( $101.6 \pm 3.3$ ).

242 From the result of blood glucose level, one way ANOVA revealed an overall  
243 significant difference ( $P < 0.05$ ) among group means. Turkey's post hoc test showed that  
244 group 2 ( $370.0 \pm 19.81$ ) significantly increased FBGL when compared with group 1 ( $94.8 \pm$   
245  $6.18$ ). Groups 3 ( $182.8 \pm 16.82$ ) and 4 ( $146.6 \pm 14.8$ ) significantly decreased FBGL when  
246 compared with group 2 ( $370.0 \pm 19.81$ ). There was no significant difference ( $P > 0.05$ ) when  
247 initial and final FBGL of groups 1 and 2 were compared. However, group 3 and 4  
248 significantly reduced FBGL after treatment when compared with initial value.

249 From table 2 above, body weight of normal rats (group 1) was significantly ( $P < 0.05$ )  
250 increased after treatment period. Body weight of diabetic control rats (group 2) was  
251 significantly ( $P < 0.05$ ) decreased after treatment. Body weight of 1% C.F.S.D treated rats  
252 (group 3) was increased after treatment. Body weight of 2% C.F.S.D treated rats (group 4)  
253 was increased after treatment. Percentage change in body weight (between before  
254 treatment and after treatment) were expressed in percentage.

255

## 256 DISCUSSION

257 The present study was undertaken to investigate the effect of *Capsicum frutescens*  
258 supplemented diet on biochemical parameters in alloxan induced diabetic Wistar rats. The  
259 action of capsaicin is mediated by TPRV1 (vanilloid receptor), which belongs to an ion  
260 channel group. VR1 when activated permits cations to pass through the cell membrane and  
261 into the cell resulting in depolarization of the neuron stimulating it to signal the brain. By  
262 binding to the VR1 receptor, the capsaicin molecule produces the same sensation that

263 excessive heat or abrasive damage would cause, explaining why the spiciness of capsaicin  
264 is described as a burning sensation. The inflammation resulting from exposure to Capsaicin  
265 is believed to be the result of the body's reaction to nerve excitement rather than just  
266 chemical burn or any direct tissue damage when chili peppers are the source of exposure.

267 Alloxan is a well- known diabetogenic agent widely used to induce Type 11 diabetes in  
268 animals [22]. Alloxan is a urea derivative which causes selective necrosis of the pancreatic  
269 islet  $\beta$ -cells. Alloxan and its reduction product dialuric acid establish a redox cycle with the  
270 formation of superoxide radicals. These radicals undergo dismutation to hydrogen peroxide.  
271 The action of reactive oxygen species with a simultaneous massive increase in cytosolic  
272 calcium concentration causes rapid destruction of beta cells, [23]. Alloxan which has been  
273 reported to destroy the beta cells of the pancreas causing reduction in insulin secretion  
274 thereby increasing blood glucose level and decreasing in body weight gain [24]. From results  
275 of the present study, the diabetic rats induced with alloxan showed these changes by  
276 decreasing body weight from  $(140 \pm 9.6)$ , before treatment to  $(120 \pm 7.9)$ , after treatment  
277 [Table 2]. Body weight of 1% and 2% *Capsicum frutescens* supplemented diet treated  
278 groups were increased more than rats in group 2. This could be traced to the recovery  
279 effects of *Capsicum frutescens* against weight loss associated with diabetes mellitus caused  
280 by alloxan monohydrate.

281 Alloxan induced diabetic is characterized by Increase in blood glucose (hyperglycemia)  
282 above normal level (normoglycemia), [24]. Increased in fasting blood glucose level (FBGL) in  
283 group 2 could be attributed to the diabetogenic effect of alloxan. Significant reduction in  
284 FBGL in 1% (group 3) and 2% (group 4) C.F.S.D treated groups may be attributed to the  
285 presence of hypoglycemic agents in *Capsicum frutescens*. Studies had shown that  
286 *Capsicum frutescens* is used to treat diabetes mellitus by traditional healers in Jamaica, [25].  
287 Pharmacokinetic and the effect of Capsaicin in *Capsicum Frutescens* on decreasing Plasma  
288 Glucose Level in a crossover study of 12 healthy volunteers by performing the OGTT while  
289 receiving placebo or 5 grams of capsicum had been documented [26].

290 In this study polydipsia and excess voiding of urine observed in group 2 rats (diabetic  
291 control) was most predominant when compared with groups 1, 3 and 4. In diabetes, the  
292 obligatory renal water loss combined with the hyperosmolarity tends to deplete  
293 intracellular water, triggering the osmoreceptor of the thirst centre of the brain and  
294 polydipsia which leads to increase in water intake, [27]. Reduce diauresis and excessive  
295 taste observed in groups 3 and 4 could be attributed to the effects of *Capsicum*  
296 *frutesence* in the diet of such rats.

297 Impaired carbohydrate utilization in the diabetic also leads to accelerated lipolysis, which  
298 results in elevated plasma triglycerides levels (hyperlipidemia), [28]. The observed  
299 abnormalities of triglyceride and HDL metabolism are in accordance with reports on early  
300 manifestation of insulin resistance, the precursor to diabetes [29; 30]. From the result of the  
301 study, 2% C.F.S.D treated group elicited reduction in serum level of total cholesterol than 1%  
302 treated group. The physiological effects of most spices had been documented to exhibit  
303 hypolipidemic and antioxidant properties with beneficial health implication, [13].

304 Individuals with type 2 diabetes had also been reported to have a higher incidence of  
305 liver function test abnormalities than non diabetic individuals. Mild chronic elevations of  
306 transaminases often reflect underlying insulin resistance. Diabetes mellitus can arise as a  
307 result of insulin insufficiency, which is associated with altered activity of various liver  
308 enzymes, [20]. Grossi, *et al.*, (1998)<sup>21</sup> had also reported that values of serum ALP can be  
309 raised in diabetic patients. The liver releases alanine aminotransferase (ALT) and an  
310 elevation in plasma concentrations are an indicator of liver damage, [30]. The levels of  
311 aspartate aminotransferase (AST), alanine amino transaminase (ALT) and alkaline  
312 phosphatase (ALP) had been reported to be increased in alloxan-induced diabetic rats, [31].  
313 Increased in serum liver enzymes parameters in diabetic control group observed in the  
314 present investigation corroborates these findings. Reduction in liver enzyme levels in group  
315 3 (1%, C.F.S.D.) and 4 (2% C.F.S.D.) clearly indicates the therapeutic role of *Capsicum*  
316 *frutescens* against increased in serum liver enzyme parameters correlated with alloxan  
317 induced diabetes. In previous research, *Capsicum frutescens* had been documented to  
318 protect against iron overload liver injury by reducing plasma liver parameters level to normal,  
319 [16].

320 There was a significant increase in serum creatinine level of group 2. An increase in plasma  
321 creatinine levels may be a sign of impaired renal function which is associated with diabetes.  
322 The elevation in the plasma creatinine concentration indirectly suggests kidney damage  
323 specifically the renal filtration mechanism, [32]. Significant reduction observed in the serum  
324 creatinine levels of the diabetic rats treated with 1% and 2% C.F.S.D in this study suggests  
325 protective effect by *Capsicum frutescens* against kidney disorders associated with diabetes  
326 mellitus.

327 Another characteristic feature of severe diabetic is an elevated excretion of urea whose  
328 concentration may be five times higher than the normal value [35]. As corroborated by  
329 this study, serum uric acid level of group 2 (Diabetic control) was significantly increased  
330 when compared with group 1 (Normal control). The significant reduction in serum uric acid  
331 level observed in the group 3 conferred protections against elevated uric acid associated

332 with diabetes mellitus. The significant reduction could be attributed to the main active  
333 principles present in *Capsicum frutescens*.

334

#### 335 **4. CONCLUSION**

336

337 From the above study increased in serum liver enzymes (AST, ALT, ALP, GGT) levels,  
338 increased in serum uric acid, creatinine, total cholesterol, fasting blood glucose level and  
339 reduced high density lipoprotein (HDL) cholesterol associated with alloxan induced diabetes  
340 mellitus were reversed after treatment with 1% and 2% C.F.S.D. Such remarkable changes  
341 observed in this study could be traced to the active ingredients [capsaicin,  
342 dihydrocapsaicin, antioxidant vitamins (ascorbic acid, vitamin E), carotenoids ( $\beta$ -carotene,  
343  $\beta$ -cryptoxanthine) and several organic acids and minerals present in *Capsicum frutescens*.  
344 Its therefore recommended that *Capsicum frutescens* be added to diet especially of diabetic  
345 patients.

346

347

348

349

350

#### 351 **AUTHORS' CONTRIBUTIONS**

352 Author 1 designed the study and wrote the first draft of the manuscript. Author 2 managed  
353 the literature searches; author 3 performed the statistical analysis and managed the  
354 analyses of the study. All authors read and approved the final manuscript.

355

#### 356 **REFERENCES**

357

- 358 1. Scoppola, A., Montecchi, F.R., Mezinger, G. and Lala, A. (2001). Urinary  
359 mevalonate excretion rate in type 2 diabetes: role of metabolic control. *Atherosclerosis* **156**:  
360 357-361.
- 361 2. Avesani, C.M., Cuppari, L., Silva, A.C., Sigulem, D. M., Cendoroglo, M., Sesso, R. and  
362 Draibe, S.A. (2001). Resting energy expenditure in predialysis diabetic patients. *Nephrol.*  
363 *Dial. Transplant.* **16**: 556-560.
- 364 3. Ogonnia SO, Odimegwu JI, Enwuru VN (2008). Evaluation of Hypoglycemic and  
365 Hypolipidemic Effects of Aqueous Ethanol Extract of *Treculia africana* Decne and  
366 *Bryophyllum Pinnatum* Lam and Their Mixture on Streptozotocin(STZ)-induced Diabetic  
367 Rats. *Afr. J. Biotechnol.*, **7**(15): 2535-2539.
- 368 4. Sofowora, A. (1993). *Medicinal Plants and Traditional Medicine in Africa*, 2nd Edition,  
369 Spectrum Books, Ibadan, Nigeria, pp. 26 -100.

- 370 5. Ugochukwu, N.H., Babady, N.E., Cobourne, M. and Gasset, S.R. (2003). The effect of  
371 *Gangronema latifolium* extracts on serum lipid profile and oxidative stress in hepatocytes of  
372 diabetic rats. *Journal of Biosciences* **28** (1): 1-5.
- 373 6. Ojewole, J.A.O. and Adewole, S.O. (2007). Hypoglycemic and hypotensive effects of  
374 *Globimetula cupulata* leaf extract in rats. *Cardiovascular. J. S. Africa.* **18**(1):9-15.
- 375 7. Ajose, F.O.A. (2007). Some Nigeria plants of dermatologic importance. *Int. J.*  
376 *Dermatology* **46** (1): 48-55.
- 377 8. Eddouks M, Jouad H, Maghrani M. Lemhadri A and Burcelin R (2003). Inhibition of  
378 endogenous glucose production accounts for hypoglycemic effect of *Spergularia purpurea* in  
379 streptozotocin mice. *Phytomedicine: International Journal of Phytotherapy andn*  
380 *Phytopharmacology* **10** (6-7): 594-599.
- 381 9. Pryor W, Stahl W and Rock C. (2000). Beta carotene: Bio chemistry to clinical Trials. *Nutr*  
382 *Rev*; **58**: 39-53.
- 383 10. Halliwell B. (1994). Antioxidants sense or speculation. *Nutr Today.* **29**: 15-19.
- 384 11. Srinivasan MR, and Chandrasekhara N. (1992). Comparative influence of vanillin and  
385 capsaicin on liver and blood lipids in the rat. *Ind. J. Med. Res.*, **96**: 133-135.
- 386 12. Srinivasan K. (2005). Role of spices beyond food flavouring: Nutraceuticals with multiple  
387 health effects. *Food Reviews Int.*, **21**: 167–188.
- 388 13. Manjunatha H, Srinivasan K (2008). Hypolipidemic and antioxidant potency of heat  
389 processed turmeric and red pepper in experimental Rats. *Afr. J. Food Sci.* **2**: 1-6.
- 390 14. Dewitt, Dave. Stock, Melissa T. and Hunter, Kellye (1998). *The Healing Powers of Hot*  
391 *Peppers.* (17-22) Three Rivers Press, NY.
- 392
- 393 15. Chaiyata P. (2003). Effect of chili pepper (*Capsicum frutescens*) ingestion on glucose  
394 response, metabolic rate, lipid profile, lipid peroxidation, thrombogenic and fibrinolytic  
395 activities in hyperlipidemic thai women. Doctoral dissertation. Bangkok: Research Unit  
396 Nutrition Faculty of Medicine Ramathibodi Hospital Mahidol University.
- 397
- 398 16. Eman A.Sadeek and Fatma H. Abd El-Razek (2010). The Chemo-Protective Effect of  
399 Turmeric, Chili, Cloves and Cardamom on Correcting Iron Overload-Induced Liver Injury,  
400 Oxidative Stress and Serum Lipid Profile in Rat Models. *Journal of American Science.* **6**  
401 (10): 42- 47.
- 402 17. Yoshioka M, Doucet E, Dropeau V, Dionne I, Tremblay A. Combined Effect of Red  
403 Pepper and Caffeine Consumption on 24 Hour Energy Balance in Subjects Given Free  
404 Access to Foods. *British Journal of Nutrition.* **85**: 203-211.

- 405 18. Antonious, G. F.; Meyer, J.; and Snyder, J. C. (2006). Toxicity and repellency of hot  
406 pepper extracts to spider mite, *Tetranychus urticae* Koch. *J. Environ. Sci. Health.* **41**, 1383-  
407 1391.
- 408 19. Story GM, Crus-Orengo L (2007). "Feel the burn". *American Scientist* .**95** (4): 326–333.
- 409 20. Siddiqui SA, Cheema AM, and Waheed M (2005). Study of serum insulin, liver profile  
410 and protein levels of insulin resistant type-2 diabetics in Pakistan population. *Pak J*  
411 *BiochemMol Biol* ; **38**(3-4); 92-7.
- 412 21. Grossi SG, Genco RJ, and Machtei E. (1988). Periodontal disease and diabetes mellitus:  
413 A two way relationship. *J Ann Periodontol*; 3: 51-61.
- 414 22. Viana G.S, Medeiros A.C, Lacerda A.M, Leal L.K, Vale T.G, and Matos F.J. (2004).  
415 Hypoglycemic and anti-lipemic effects of the aqueous extract from *Cissus sicyoides*. *BMC*  
416 *Pharmacol*, **8**: 4-9.
- 417 23. Szkudelski T. (2001). The mechanism of alloxan and streptozotocin action B cells of the  
418 rat pancreas. *Physiology, Res.* **50**:536-546.
- 419 24. Al-Kalifa, A., Mathew T.C., Al-Zaid N.S., Mathew E. and Dashti H.M. (2009).  
420 Therapeutic role of low-carbohydrate ketogenic diet in diabetes. *Nutrition*, **25**: 1177-1185.
- 421 25. Tolan I, Ragoobirsingh D, and Morrison EY (2004). Isolation and purification of the  
422 hypoglycaemic principle present in *Capsicum frutescens*. *Phytotherapy Research.*; **18**(1):95-  
423 6.
- 424 26. Kamon Chaiyasit, Weerapan Khovidhunkit, and Supeecha Wittayalerpanya. (2009).  
425 Pharmacokinetic and The Effect of Capsaicin in *Capsicum frutescens* on Decreasing Plasma  
426 Glucose Level. *J Med Assoc Thai.* **92** (1): 108-113
- 427 27. UKPDS (UK Prospective Diabetes Study Group), (1998). Intensive blood glucose with  
428 sulphonylureas or insulin compared with conventional treatment and risk of complications in  
429 patients with Type 2 diabetes. *Lancet.* **353**: 837-853.
- 430 28. Granner DK, Mayes PA, and Rodwell VW (1996). Harper's Biochemistry, ed 24,  
431 Connecticut, USA, Appleton and Lange, pp. 586-587.
- 432 29. Frederickson DS, and Lee RS. (1965). A system for phenotyping hyperlipidemia.  
433 *Circulation*, **31**: 321-327.

- 434 30. Lyons TJ (1992). Lipoprotein glycation and its metabolic complications. *Diabetes*.  
435 **41**(Suppl 2): 67-73.
- 436 31. Claudia ENM, Julius EO, Dagobert T, and Etienne D (2006). Antidiabetic and  
437 Hypolipidemic effects of *Laportea ovalifolia* (URTICACEAE) in alloxan induced diabetic rats.  
438 *Afr. J. Tradit., Complement. Altern. Med.* **3**(1): 36-43.
- 439 32. Akah P.A, Alemji, J.A, Salawu O. A., Okoye T.C. and Offiah N.V, (2009). Effects of  
440 *Vernonia amygdalina* on Biochemical and Hematological Parameters in Diabetic Rats. *Asian*  
441 *Journal of Medical Sciences* **1**(3): 108-113.
- 442 33. Wasan KM, NajafiS, Wong J, Kwong M (2001). Assessing plasma lipid levels, body  
443 weight, and hepatic and renal toxicity following chronic oral administration of a water soluble  
444 phytostanol compound FMVP4, to gerbils. *J. Pharm. Sci.* **4**(3): 228-234.
- 445 34. Lehninger, A.L., (1998). Principles of Biochemistry. CBS Publishers and Distributors  
446 Pvt. Ltd., India, pp: 531-535.
- 447