

# Assessment of Nutrition Knowledge Among University Students in Ankara

**Aims:** Nutrition knowledge is one of the factors that could influence an university student's nutritional behaviours. This research was aimed to determined nutrition knowledge of university students in Ankara

**Study design:** The study was conducted in Ankara, Turkey between May and July, 2014 in university students.

**Place and Duration of Study:** This was a cross sectional study.

**Methodology:** Out of 341 students, 66.3% % were female (n=226), 33.7% % were male (n=115). Our questionnaire included a demographic section, and 25 true-false nutrition knowledge questions. For the reliability of the questionnaire, the internal consistency coefficient was calculated, and the Kuder Richardson (KR-20) value was found to be 0.82. For the statistical analyses of the data, table showing mean, standard deviation ( $\bar{X} \pm SD$ ) and percentage (%) values were prepared. When identifying the nutrition knowledge of students, the "independent t test" was used for the as taking gender and age.

**Results:** The mean nutrition knowledge score was  $15.8 \pm 4.9$ . The mean score for gender was  $16.6 \pm 4.3$  in females, and  $14.2 \pm 5.5$  in males, and found statistically significant ( $p=.000$ ).

**Conclusion:** In this study, was determined that nutrition information scores of the participants were moderate. This situation is the result of increasing the level of nutrition knowledge in young people. Nutrition knowledge may be effective in increasing the quality of life and decreasing the prevalence of some diseases.

*Keywords: nutrition knowledge, gender, age, university*

## 1. INTRODUCTION

Being able to keep human force at its highest level in terms of physical and mental functions is closely related to human nutrition. One can be unaware of nutrition values of various foods; what foods are suitable for health or what he knows about them might be wrong [1]. Lack of nutritional knowledge or wrong knowledge over nutrition could lead to serious health problems based on nutrition (obesity, diabetes, cardiovascular diseases etc.) in the future [2-4].

Providing knowledge of nutrition is realized through true nutrition education. Educational programs for nutrition have a direct impact on the knowledge of nutrition and nutritional behaviors [5] Knowledge of nutrition is a great factor having an effect on the nutritional behaviors of families and communities [1]. Basic aim in nutritional education is to give the information with regard to its relation to nutrition and which healthy food to eat [6].

With an increase in health service activities, improving dietary and there will be a decrease in health complications and untimely deaths [7].

University life is a period when some significant changes occur in the life of individuals [8-9]. Differentiating together with university, life style could have some changes in the nutritional behaviors of the students. Changing nutritional behaviors do not only deal with the mental and physical status of the university student, That's why, Increasing the nutrition information of university students has an impact on maintaining a healthier life [10-13]. In the current study, was aimed to determined nutrition knowledge of university students according to gender and age.

31 **2. MATERIAL AND METHODS**

32  
33 The sampling of the research was made up of 341 volunteer students attending to various universities  
34 in Ankara. Ankara is the capital city of Turkey. The number of participants has been determined by the  
35 duration of the study. Participants were included in the study by voluntarily obtaining the informed  
36 consent form. The study is a descriptive research. The research data were collected through a  
37 questionnaire and face to face interviews. The questionnaire form was composed of two sections, the  
38 first of which was designed to obtain information about the demographic characteristics of the  
39 students and the second part contained statements related to nutrition knowledge. Statements were  
40 prepared for an examination of the relationship between nutrition and health. In order to evaluate their  
41 knowledge on nutrition, the students who participated in the study were given 25 statements which  
42 they can reply as "true" or "false". At the stage of developing items, some sources were used [1,  
43 14,15]. All authors hereby declare that all experiments have been examined and approved by the  
44 appropriate ethics committee and have therefore been performed in accordance with the ethical  
45 standards laid down in the 1964 Declaration of Helsinki. This research was prepared in accordance  
46 with the Helsinki Declaration principles Ethics Committee Approval is obtained from Assessment  
47 Commission of Non-Interventional Research of Ankara University; (273 numbered decision in  
48 12.12.2013 year).

49

50 **Statistical Analysis**

51 After administering the questionnaire to the individuals and assessing it, reliability test was applied.  
52 For the reliability of the questionnaire, "Kuder Richardson", the internal consistency coefficient, was  
53 calculated, and the KR-20 value was found to be 0.82. As the results of reliability were not low, all the  
54 item were not included. Accordingly, it was agreed that the "Nutrition knowledge" scale was a reliable  
55 instrument.

56 While the nutrition knowledge was being evaluated, 1 point was given to each correct answer,  
57 whereas wrong answers were not given any points. The students' knowledge status was evaluated  
58 with 25 points by giving a score in the right answer for each item with the data collection tool. The  
59 data of the study were evaluated using SPSS statistical package program. Nutritional knowledge of  
60 the students were examined in terms of age and gender variables. For the statistical analyses of the  
61 data, table showing mean, standard deviation ( $\bar{x} \pm SD$ ) and percentage (%) values were prepared.  
62 When identifying the nutrition knowledge of students, the "independent t test" was used for the as  
63 taking age and gender. A criterion alpha level of < 0.05 was used to determine statistical significance.

64

65 **3. RESULTS AND DISCUSSION**

66

67 **Descriptive data**

68 Of the participants of the research, 66.3 % (n:226) were girls and 33.7% % (n:115) were boys. The  
69 mean age of the students was 21.06±1.56 (girl: 21.0±0.6 years, boy: 21.2±1.5 years)

70 Nutritional knowledge has a direct effect on the health of individuals. It is likely to prevent a great  
71 many health problems with an awareness of healthy nutritional principles and their application [1]. The  
72 percentages of true answers of the items asked to students with regard to nutrition were given in  
73 Table 1.

74 **Table 1. Nutritional knowledge of students**

Statements	%
Consuming fish 2-3 times a week decreases the risk of cardiovascular diseases (T).	80.9
Lack of vitamin C leads to tooth gum bleeding (T).	65.7
When not taking enough vitamin D, children could have the disease of rachitism (T).	80.6
Overconsumption of salt could lead to hypertension (T).	86.8
Overconsumption of fried food increases the risk of cancer (T).	90.9
An increase in the energy from fat in diet increases the risk of cardiovascular diseases (T).	57.8
Overconsumption of fiber food increases the risk of intestinal cancer (F).	35.5
The risk of having anemia is more in those not consuming red meat (T).	55.1

Consuming iodine salt is not effective in preventing the development of goitre (F).	52.8
The risk of having diabetes in obese people is higher than the thin ones (T).	72.7
Consuming an egg a day by healthy people does not affect cholesterol level (T).	47.2
Consuming less fiber food causes constipation (T).	51.3
Inadequate flour intake causes tooth decay (T).	77.1
Inadequate flour intake leads to mental retardation (T).	41.6
Consuming milk and dairy products less leads to osteoporosis only in women (F).	32.0
Inadequate vitamin D taking leads to softening in bones and teeth (T).	72.7
Consuming tea in meals leads to diminishing iron absorption of food and causes anemia (T).	80.6
Green leaved vegetables consumed at breakfast help to prevent anemia (T).	64.2
Consuming less food than needed by diabetics leads to decrease in blood sugar (T).	56.9
Overconsumption of food with saturated fat and having high cholesterol content leads to cardiovascular diseases (T).	87.4
Consuming leguminous food increases bad cholesterol (LDL) (F).	31.7
A, C, E vitamins help to prevent cancer by protecting cells (T).	59.2
Eat fast and chewing less could lead to (T).	78.9
Fast food menus could lead to hypertension because of their sodium content besides salt (T).	60.1
Skipping main meal leads to obesity (T).	59.8

Note: (T) = true, (F) = false.

75  
76

77  
78

According to Table 1, the first three expressions answered correctly in high rate: (Overconsumption of fried food increases the risk of cancer 90.9%, Overconsumption of food with saturated fat and having high cholesterol content leads to cardiovascular diseases 87.4%, Overconsumption of salt could lead to hypertension 86.8%). High rates of wrong answers given phrases (Consuming leguminous food increases bad cholesterol (LDL) 31.7%, Consuming milk and dairy products less leads to osteoporosis only in women 32.0%, Overconsumption of fiber food increases the risk of intestinal cancer 35.5%).

84

85

Basic element in protecting oneself from coronary diseases is the change in life and nutrition style. The item of "Consuming fish 2-3 times a week decreases the risk of cardiovascular diseases" replied by 80.9% of the participants as true. Important fat acids in fish have a positive impact on cardiovascular health [16]. As for the item of "An increase in the energy from fat in diet increases the risk of cardiovascular diseases", the rate of answering it was 57.8%. Besides decreasing the energy coming from fat in diet, providing a variety of diet fat types should not be neglected [7]. An increase in taking mono and multi fat acids and a decrease in saturated fat acids could decrease the risk of cardiovascular disease risk to a great extent [16]. Due to the fact that egg is a sample of protein, there is no problem with consuming an egg a day by the ones having no health problems [17].

93

94

The rate of the ones knowing that overconsumption of the food having a high content of saturated fat and cholesterol could lead to cardiovascular diseases was 87.4%. However, the important point here is to decrease total daily fat consumption. The rate of answering the item "Consuming leguminous food increases bad cholesterol (LDL)" as true saying that it is false was 31.7%. It would be possible to prevent cholesterol accumulation in veins with a decrease of LDL-cholesterol in diet and an increase in HDL-cholesterol [18]. Due to the fiber content, legume is helpful in showing this effect.

99

100

101

102

103

104

105

106

107

108

109

110

111

Since vitamin C sources are not well benefitted in winter months in rural areas, low and medium level vitamin C deficiency symptoms could be seen [1]. It was found that the participants answered the item "Lack of vitamin C leads to tooth gum bleeding" as true at the rate of 65.7%. And 55.1% of the participants knew that those not consuming red meat have a higher possibility of anemia. Iron absorption in animal based food is higher. In the prevention of a great variety of anemia, consumption of animal based foods are effective. There is a need to inform people over increasing the consumption of such kind of foods. It is also of a great importance to increase this rate in the university years which are considered to the last period of the development. The tannins in tea ~~decreases~~ decrease the bioavailability of iron [1]. The item of "Consuming tea in meals leads to diminishing iron absorption of food and causes anemia" was answered correctly at a high rate (80.6%). The rate of answering the item of "Green leaved vegetables consumed at breakfast help to prevent anemia" correctly was 64.2%. In particular, due to the fact that green leaved vegetables

112 consumed at breakfast are rich in vitamin C, they are significant nutrients increasing iron absorption  
113 [17].

114 Goitre is an important health problem [ef in Turkey](#). The item of “Consuming iodine salt is not effective  
115 in preventing the development of goitre” was answered correctly by half of the participants (52.8% [%](#)).  
116 Increasing iodine salt consumption will be effective in preventing such kind of diseases. Only 41.6%  
117 % of the students knew the item of inadequate iodine intake could lead to mental retardation. As a  
118 result of iodine deficiency, the hormones passing into blood through thyroid gland cannot be produced  
119 much enough and there might occur some problems in the development of organs and in their  
120 functions. Mental functions retard [19]. The rate of saying true for the item “Inadequate florin intake  
121 causes tooth decay” asked to students were 77.1% [%](#). Mentioning more about florin in toothpaste  
122 commercials show that there is an awareness in this issue. Vitamins have important functions in  
123 preventing various cancer types. A, C and E vitamins could prevent the development of cancer cells  
124 by preventing the formation of free radicals [20]. The rate of answering the item “A, C, E vitamins help  
125 to prevent cancer by protecting cells” true depending on this information was 59.2% [%](#).

126 In societies having a higher average salt consumption, blood pressure increases with age [21].  
127 Cutting down on salt consumption leads to a decrease in blood pressure and this case means a  
128 decrease in a significant risk factor for cardiovascular diseases. The fact that the rate of those saying  
129 that overconsumption of salt leads to hypertension was high (86.8% [%](#)) is good news. In the fast food  
130 style of nutrition, vitamin C, A, calcium and fiber intake is inadequate and fat and sodium consumption  
131 is higher [22]. As for the item of “Fast food menus could lead to hypertension because of their sodium  
132 content besides salt”, the rate of true answer was 60.1% [%](#). Relatively low rate in this item could  
133 result from unawareness of the fact that the rate of salt in fast food products is not known clearly.  
134 Some healthy foods (qualitative protein, low fat and whole-wheat products, salads enriched with  
135 lemon and vinegar) have been included in the menus of fast food restaurants recently. Therefore, it is  
136 necessary that the young should be informed to prefer healthy menus in fast food restaurants.

137 Faulty processes of the food products could increase the risk of developing cancer. Since the method  
138 of frying changes the structure of oil, it increases the formation of carcinogenic materials. The item  
139 “Frequent consumption of fried foods increases the risk of cancer” asked in this respect was replied  
140 true at a high rate (90.9% [%](#)). Fiber arranges intestinal activity and prevents constipation. Two items  
141 were asked with regard to diet fiber. The rate of the students being aware of the fact that consuming  
142 fiber food more reduces the risk of large bowel was 35.5% [%](#), while that of the ones knowing that  
143 consuming fiber food less could lead to constipation was 51.3% [%](#).

144 Obesity could lead to a great many health problems due to the negative effect on body systems and  
145 psychosocial cases [1]. Skipping meals, having frequent snacks and eating fast are among most  
146 import faulty behaviors causing obesity. The rate of those knowing that skipping main meals could  
147 lead to obesity was 59.8% [%](#) and the ones knowing that eating fast and chewing less could also lead  
148 to obesity was 78.9% [%](#).

149 Obesity one of the preventable risk factors in diabetes development. Increased weight gaining and the  
150 duration of obesity could also increase the risk of developing diabetes [22]. In the research, two items  
151 were included with regard to measuring diabetes awareness. The rate of participants saying that “The  
152 risk of having diabetes in obese people is higher than the thin ones” was 72.7% [%](#) while that of the  
153 ones saying “Consuming less food than needed by diabetics leads to decrease in blood sugar” was  
154 56.9% [%](#).

155 Among vitamin D and calcium functions are the maintenance of bone and tooth health. In the case of  
156 deficiency, bone mineralization is broken and rachitism could be seen at children while osteoporosis  
157 could be encountered at the elderly [23]. Of the students, 80.8% [%](#) knew that the disease of rachitism  
158 could be encountered at children in the case of not taking enough vitamin D and 72.7% [%](#) of them  
159 knew that when vitamin D is taken inadequately, there might occur some softening in bones and  
160 teeth. In the maintenance of bone health, it is of great importance to consume milk and dairy products  
161 as they are a good source of calcium [17]. The rate of those answering the item of “Consuming milk  
162 and dairy products less leads to osteoporosis only in women” wrongly saying “Yes” was 68.8% [%](#). It is  
163 thought that the answer given this item correctly at the rate of 1/3 could result from the fact that  
164 osteoporosis is known to be a woman disease.

165 **Nutrition knowledge score**

166 The mean nutritional scores, standard deviation and t test results of the students in terms of gender  
167 and age group variable were given in Table 2.

**Table 2. Mean nutritional scores of students in terms of variables**

Age(year)	n	$\bar{X}$	SD	df	t	P
18-21	218	16.14	4.77	339	1.73	.084
22-24	123	15.18	5.07			
<b>Gender</b>						
Female	226	16.61	4.34	339	4.41	.000
Male	115	14.20	5.50			
<b>Total</b>	<b>341</b>	<b>15.80</b>	<b>4.90</b>			

168

169

170 Nutrition information; is one of the factors affecting the nutritional status and habits of individuals,  
171 families and societies [1]. It is known that education that will be given to the individual is a dynamic  
172 process and must be continuous in order to comply with the differences as conditions are constantly  
173 changing [24]. In our study, the highest points that can be taken 25. In general sampling, mean  
174 nutritional knowledge score was 15.80±4.90. Depending on the age group, it was 16.14±4.77 at 18-21  
175 age range and 15.18±5.07 at 22-24 age range. In terms of gender, mean nutritional knowledge scores  
176 of girls (16.61±4.34) was higher than those of boys (14.20±5.50). Mean nutritional knowledge scores  
177 taken depending on gender are of significance statistically ( $P= .000$ ). In a study, İlhan et al. [9] and  
178 Şanlıer et al. [25] found that nutritional knowledge scores of girls were higher than those of boys. İlhan  
179 et al. [9] found in a study with regard to the healthy life style behaviours of university students that  
180 mean nutritional knowledge score of girls was 15.55±3.32 and it was 14.81±3.14 at boys. Şanlıer et  
181 al. [25] found that mean nutritional knowledge score of boys attending to a university was 5.65±5.55  
182 and it was 6.05±2.38 at girls, and that the score difference between genders was statistically  
183 significant ( $P= .05$ ). They found that although the female students had more knowledge, the nutrition  
184 information of the young people was inadequate and inadequate information could not turn into habits  
185 and behavior [25]. In an other study conducted, 63.1% % of the students were found to have  
186 moderate nutritional knowledge, 9.0% % poor, 27.0% % good and 0.9% % very good [26].

187

188

#### 4. CONCLUSION

189

190 It was determined that nutrition information scores of the participants were moderate. Nutritional  
191 knowledge has a direct impact on the nutritional status of individuals and also on their habits.  
192 Therefore, it is necessary that the importance of nutritional education should be taken into  
193 consideration in the maintenance and development of health. As nutrition is an indispensable part of  
194 human being, it is also required that students should be made to review their nutritional knowledge. It  
195 is thought that nutrition education programs for young is needed not only to get correct nutrition  
196 knowledge but also to promote affirmative dietary behavior and the volition to practice nutritionally  
197 balanced meals and to induce changes in nutritional behavior. It is known that Universities provide to  
198 various opportunities to increase consciousness awareness of nutritional among university students.  
199 Including nutrition courses in the programs of higher education, placing it in the state politics and  
200 providing its maintenance is believed to have a considerable contribution to the awareness of the  
201 issue.

202

203

#### STRENGTHS AND LIMITATION

204

205 It is advantageous to work with young groups. Findings cannot generally be generalized to the study  
206 population or community, because this research could not be done at various universities. The limited  
207 time for research has led to the limitation of the number of samples. Failure to obtain a daily nutrient  
208 intake lead to a restriction of study.

209

210 **CONSENT**

211

212 All authors declare that 'written informed consent' was obtained from the participants for publication of  
213 this study.

214

215

216 **REFERENCES**

217

218 1. Baysal A. Beslenme. Yenilenmiş 17th ed. Ankara, Turkey: Hatipoğlu Yayınları; 2017. Turkish.

219 2. Omid MG, Torabi R, Hojjati M, Ghanian M, Kitterlin M. Factors influencing Iranian consumers'  
220 attitudes toward fast-food consumption. *British Food Journal*. 2018;120(2):409-23.

221 3. Bazzano LA. The high cost of not consuming fruits and vegetables. *Journal of the American*  
222 *Dietetic Association*. 2006;106(9):1364-68.

223 4. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels:  
224 surveillance progress pitfalls and prospects. *Lancet*. 2012;380:247-57.

225 5. Jonides L, Bushbacher MS, Barlow SE. Management of child and adolescent obesity:  
226 psychological, emotional and behavioral assessment. *Pediatrics*. 2002;110(1):215-21.

227 6. Kyrkou C, Tsakoumaki F, Fotiou M, Dimitropoulou A, Symeonidou M, Menexes G, et al. Changing  
228 trends in nutritional behavior among university students in Greece, between 2006 and 2016.  
229 *Nutrients*. 2018;10(1), 64.

230 7. Benjelloun S. Nutrition transition in Morocco. *Public Health Nutr*. 2002;5(1A):135-40.

231 8. Aceijas C, Waldhäusl S, Lambert N, Cassar S, Bello-Corassa R. Determinants of health-related  
232 lifestyles among university students. *Perspectives in public health*. 2017;137(4):227-36.

233 9. İlhan N, Batmaz M, Akhan, LU. Üniversite öğrencilerinin sağlıklı yaşam biçimi davranışları. *Maltepe*  
234 *Üniversitesi Hemşirelik Bilim ve Sanatı Dergisi*. 2010;3(3):34-44. Turkish.

235 10. Keiner F, Hasbargen U. Emergencies associated with pregnancy and delivery: peripartum  
236 hemorrhage. *Dtsch Arztebl Int*. 2008;105(37):629-38.

237 11. Elsayes KM, Trout AT, Friedkin AM, Liu PS, Bude RO, Platt JF, et al. Imaging of the placenta: a  
238 multimodality pictorial review. *Radiographics*. 2009;29:1371-91.

239 12. Rao AP, Bojahr H, Beski S, MacCallum PK, Renfrew I. Role of interventional radiology in the  
240 management of morbidly adherent placenta. *J Obstet Gynecol*. 2010;30:687-89.

241 13. Wong HS, Cheung YK, Williams E. Antenatal ultrasound assessment of placental/myometrial  
242 involvement in morbidly adherent placenta. *Aust N Z J Obstet Gynaecol*. 2012;52(1):67-72.

243 14. Brown A. Understanding food principles and preparation. Wadsworth: USA. 2000.

244 15. Şahin H. Obezite'de beslenme tedavisi ve doğru bilinen yanlışlar. P Arslan, A Dağ, Türkmen EG  
245 (eds.), *Her Yönüyle Obezite: Önleme ve Tedavi Yöntemleri*. 1th ed. Ankara: 2012. Turkish.

246 16. Baysal A. Beslenme, lipoproteinler ve koroner kalp hastalığı. *Beslenme Diyet Dergisi*.  
247 2006;34(1):5-9. Turkish.

248 17. Turkey Dietary Guidelines. Ministry of Turkey Health Publication No: 1046, Kayhan Ajans,  
249 Ankara: Turkey. 2016. p. 293.

- 250 18. Joseph N, Chettuvatti K, Yadav H, Bharadwaj H, Kotian SM. Assessment of risk of metabolic  
251 syndrome and cardio vascular diseases among medical students in India. Journal of Cardiovascular  
252 Disease Research. 2017; 8(3).
- 253 19. Pekcan G. İyot yetersizliği hastalıkları, çocuklarda önemi ve önlenmesi. Klinik Çocuk Forumu  
254 Dergisi. Kasım-Aralık. 2004;12-24. Turkish.
- 255 20. Aksoy M. Kanserde beslenme. Diyet El Kitabı, 9th ed. Ankara, Turkey: Hatipoğlu Yayınevi; 2016.  
256 Turkish.
- 257 21. Sağlık Bakanlığı Temel Sağlık Hizmetleri Genel Müdürlüğü Beslenme ve Fiziksel Aktiviteler Daire  
258 Başkanlığı, Türkiye aşırı tuz tüketiminin azaltılması programı. Yayın No:835, Deniz matbacılık,  
259 Ankara. 2011. s. 52. Turkish.
- 260 22. Whitney E, Rolfes SR. Understanding nutrition. 15th ed. USA: Cengage Learning; 2018.
- 261 23. Jones KD, Hachmeister CU, Khasira M, Cox L, Schoenmakers I, Munyi C, et al. Vitamin D  
262 deficiency causes rickets in an urban informal settlement in Kenya and is associated with malnutrition.  
263 Maternal & child nutrition. 2018;14(1).
- 264 24. Merdol TK. Beslenme eğitimi ve danışmanlığı. Basım 2. Yayın No: 726. Reklam Kurdu  
265 Ajansı.Sağlık Bakanlığı. Ankara. 2012. Turkish.
- 266 25. Şanlıer N, Konaklıoğlu E, Güçer E. Gençlerin beslenme bilgi, alışkanlık ve davranışları ile beden  
267 kütle indeksleri arasındaki ilişki. Gazi Eğitim Fakültesi Dergisi. 2009;29(2):333-52. Turkish.
- 268 26. Tütüncü İ, Karaismailoğlu E. Üniversite öğrencilerinin beslenme bilgi düzeylerinin belirlenmesi.  
269 Uluslararası Hakemli Akademik Spor Sağlık ve Tıp Bilimleri Dergisi, 2013;3(6):29-42. Turkish.