

Original Research Article

BARBING EQUIPMENTS: TOOLS FOR TRANSMISSION OF TINEA CAPITIS

Abstracts

This efficacy of sterilization and cleaning methods used in the barbing salons and the possibility of those equipments acting as vehicles of transmission of fungal pathogens from one customer to another was assessed in this study. Thirty barbing saloons were randomly selected from the five wards in Calabar South. One hundred and fifty samples were aseptically collected using sterile swab sticks to brush out debris from combs, hair brushes and sterilized clippers into sterile paper envelopes. Samples were subjected to microscopy, culture and physiological tests. The recovery rate of fungal organisms from barbing equipments was 52(34.7%). Dermatophytes recovery rates was 40(26.7%) and non dermatophytes recovery rates was 12(8.0%). There was a statistically significant relationship between dermatophytes and barbing equipments recovered from ($\chi^2 = 25.0$, $p = 0.01$). *Trichophyton interdigitale* 12(30.0%). was the most common dermatophyte recovered from barbing equipments, followed by *Trichophyton rubrum* 9(22.5%). *Malassezia* species 5(41.6%) was the most common non dermatophytes recovered, followed by *Aspergillus flavus* 3(25.0%). Hair brushes were the most contaminated equipments (60.0%) by dermatophytes while the least contaminated were the hair clippers 17.5%. Dermatophytes and non dermatophytes were recovered from barbing equipments which depicts that they could serve as vehicles for transmission of fungal infections. Although the clippers were sterilized by all Barbers, dermatophytes and few non dermatophytes were still recovered from them. This points to incomplete sterilization either by disinfectant inefficacy or methods inefficiency.

Key words: Barbing equipments; Tine capitis, sterilization; transmission

Introduction

Barbers are important professionals in the community. Barber's shops are owned and managed by individuals in the community. Barbers use instruments such as knives, blades, clippers, brushes and combs which may serve as vehicles for transmission of fungal infections [1-2].

From 2002 till date, The establishment of barbing salons in our locality have been on the increase. Majority are owned and controlled by people with little or no training and knowledge on infection controlled practices. Researchers have shown the possibility of

37 customers returning home with one infection or the other despite the aesthetics in the
38 barbing industry [3].

39 Tinea capitis affects the scalp and hair. The infection is caused by *Trichophyton*,
40 *Microsporum* and *Epidermophyton* species of dermatophytes [4]. The dermatophytes
41 invade hair follicles and keratinized layer of hairy skin leading to scaling, kerion, hair loss,
42 folliculitis, favus and black dot [4]. The infection could be anthropophilic, zoophilic or
43 geophilic [4-5]. Prepubertal school age children between 6 to 10 years are usually more
44 affected, it is rare in adults. The infection occurs in both sexes with high infection rates
45 among males [4, 6].

46 *Tinea capitis* is of public health importance but not a notifiable disease therefore little
47 is known on its prevalence in many endemic areas [7-8]. Tinea capitis can be transmitted
48 via infected persons, animal vectors and fallen infected hairs. Spread of tinea capitis by
49 fomites (contaminated barbershop instruments, hairbrushes, combs and shared hats) is
50 common [9-10]. Factors affecting disease transmission are personal hygiene, overcrowding
51 and low socioeconomic status. Shed hairs may harbour infectious fungal agents up to a
52 year. Eradication is difficult due to common asymptomatic carriers [11]. Barbering
53 operations are still under little or no scrutiny despite the possible risk of spreading infectious
54 diseases [10].

55 Barbers use equipments which are always contaminated after some customer
56 usage. Their sterilization methods include disinfecting and flaming. Some of the materials
57 used are never sterilized, while some are sterilized but the sterilization efficacy is
58 questionable since customers still get infected after saloon visits. This study was designed
59 to assess the possibility of transmission of tinea infections from otherwise sterilized barbing
60 equipments in Calabar, Nigeria.

61

62 **Materials and methods**

63 The study area for this research was Calabar South Local Government Area of
64 Cross River State. It has an area of 264km² and the population of 191,630 with the
65 headquarters at Anantigha. Thirty barbing saloons were randomly selected from the five
66 wards in Calabar South. Ethical approval was not sought since samples were obtained from
67 inanimate objects. One hundred and fifty samples were aseptically collected using sterile
68 swab sticks to brush out debris from combs, hair brushes and sterilized clippers into sterile
69 paper envelopes. Samples were transported to the Department of Medical
70 Microbiology/Parasitology, University of Calabar Teaching Hospital, Calabar for analysis.

71 Samples were subjected to microscopy using 10% KOH. The samples were cultured
72 on dermatophyte test medium (DTM) with actidione (LabM Limited, Lancashire, UK) and
73 Sabouraud dextrose agar with chloramphenicol [12]. Plates were incubated at room tem-
74 perature and at 37°C, and examined for growth up to 2 weeks.

75 Hair fragments from the barbing equipments were placed on the microscope slide. A
76 drop of 10%KOH was added and covered with cover slip. The slides were examined
77 microscopically for arthroconidia in an ectothrix (outside the hair shaft) or endothrix (within
78 the hair shaft) patterns [13].

79 Pure culture of the isolates were obtained by sub culturing individual isolates onto
80 fresh Sabouraud dextrose agar (SDA) at room temperature (26°C-29°C) for up to 10-14
81 days. In cases of no or slow sporulation, slide culture was carried out [12]. Isolates were
82 identified by colonial morphology, microscopy and physiological tests. The physiological
83 tests used in the identification of the isolates include: in vitro hair perforation test and
84 urease test [1].

85 Isolates were identified based on colonial morphology: (color, pigmentation, rate of
86 growth and texture), microscopy and physiological tests. The Physiological tests carried out

87 include: urease test, hair penetration test, growth on rice grains, tween assimilation and lipid
88 test [1, 14-17]. The isolates were compared to published descriptions of these organisms.

89 Pure cultures of isolates were inoculated on Christensen's urea slants. A change in
90 colour from yellow to pink indicated a positive urease production.

91 Hair perforation test was performed in a Petri dish containing 20ml of deionized water
92 with 2-3 drops of 10% yeast extract. Sterilized hair fragments from a prepubescent child
93 was placed in the dish and inoculated with the isolate to be tested. The petri dishes were
94 incubated at room temperature for up to 4 weeks. Positive and negative controls were also
95 set up with *Trichophyton interdigitale* and *Trichophyton rubrum* respectively. The hair
96 fragments are examined weekly under the microscope as wet preparations in a drop of
97 water. The observation of a cone shaped perforation on the hair shaft means positive.
98 When viewed from the top they appear as round holes [13].

99 In vitro hair perforation test and urease test was used to differentiate *T. soudanense*
100 and *T. Tonsurans*. *Trichophyton soudanense* is usually urease negative and hair perforation
101 test negative while *T. Tonsurans* is urease positive and hair perforation test variable.

102 Growth on rice grains was used to distinguish *M. gypseum* from *M. audouinii*. Few
103 grains of polish rice were were placed in a small flask and covered with distilled water
104 before autoclaving. The isolates were inoculated onto the surface of the grains and
105 incubated for 7-14 days. *Microsporum gypseum* produces good growth, with no
106 pigmentation but with variable conidia formation. *Microsporum audouinii* does not grow on
107 rice grain. The identification scheme for the dermatophytes is shown in Table 1.

108 Catalase test, lipid test and tween assimilation test were used in the identification of
109 *Malassezia* species [16-18]. *Aspergillus* species were identified based on their colonial and
110 microscopic morphology.

111

112

Table 1 Identification scheme for dermatophytes

Dermatophytes isolates	Invitro hair perforation test	Hair invasion	Growth on rice grains	urease
<i>Microsporum gypseum</i>	+	Large spores ectothrix	Good growth no pigmentation. (+) or (-) conidia	+
<i>Microsporum audouinii</i>	-	Ectothrix	Poor with no conidia	(-) or (+)
<i>Trichophyton interdigitale</i>	- (positive) Meaning please????	Negative	-	+
<i>Trichophyton rubrum</i>	-	Negative	N/A	+
<i>Trichophyton tonsurans</i>	Variable	Endothrix	N/A	+
<i>Trichophyton soudanense</i>	Negative	Endothrix	N/A	Usually negative

113

114

115 Data analysis

116 The data obtained in this study were analyzed with Epi-Info CDC, 2012 data analysis
 117 package. Descriptive statistics were carried out. Frequencies were calculated for
 118 categorical variables. Interactions between specific categorical variables were tested for
 119 significance using the χ^2 test. A p-value of 0.05 was considered statistically significant.

120

121 Results

122 Out of the 150 samples collected from hair barbing equipments 52 (34.7%) were infected
 123 with fungal organisms. Figure 1 shows the distribution of fungal isolates by barbing
 124 equipments. Hair brushes were the most contaminated barbing items 27 (54.0%) followed
 125 by combs 15 (30.0%). Only 10 (20.0%) isolates were recovered from hair clippers.

126 Dermatophytes recovery rates was 40(26.7%). The most common dermatophyte was
 127 *Trichophyton interdigitale* 12 (30.0%), followed by and *Trichophyton rubrum* 9(22.5%) and
 128 *T. tonsurans* 6 (15.0%). *Trichophyton soudanense* 3 (7.5%) was the least recovered isolate.

Hair brushes were the most contaminated equipments 24 (60.0%) by dermatophytes and the least contaminated were the hair clippers 7 (17.5%). There was a statistically significant relationship between the barbing equipments and dermatophytes recovery rates ($\chi^2 = 25.0$, $p = 0.01$) (Table 2).

Non dermatophytes recovery rates was 12(8.0%). The most common non-dermatophyte recovered from barbing equipments was *Malassezia* species 5(41.6%) followed by *Aspergillus flavus* 3(25.0%). Combs were the the most contaminated barbing items 6(50.0%) by non-dermatophytes. There was no statistically significant relationship between non-dermatophytes recovery rate and barbing equipments ($\chi^2 = 11.0$, $p = 0.16$) (Table 3).




Fig 1 Distribution of fungal isolates in barbing equipments

Table 2 Distribution of dermatophytes on barbing equipments

Dermatophytes isolates	Barbing equipments (n=150)
	No (%) of isolates

	Clippers	Brushes	Combs	Total
<i>Microsporum audouinii</i>	0(0.0)	4(16.6)	1(11.1)	5(12.5)
<i>Microsporum gypseum</i>	0(0.0)	4(16.6)	1(11.1)	5(12.5)
<i>Trichophyton interdigitale</i>	2(28.5)	7(29.1)	3(33.3)	12(30.0)
<i>Trichophyton rubrum</i>	2(28.5)	5(20.8)	2(22.2)	9(22.5)
<i>Trichophyton soudanense</i>	0(0.0)	1(4.2)	2(22.2)	3(7.5)
<i>Trichophyton tonsurans</i>	3(42.8)	3(12.5)	0(0.0)	6(15.0)
TOTAL	7(17.5)	24(60.0)	9(22.5)	40(26.7)

Table 3 Distribution of non-dermatophyte isolates on barbing equipments

Non-dermatophyte isolates	Clippers	Brush	Combs	Total
<i>Aspergillus flavus</i>	0(0.0)	2(66.7)	1(16.7)	3(25.0)
<i>Aspergillus fumigatus</i>	2(66.7)	0(0.0)	0(0.0)	2(16.7)
<i>Aspergillus niger</i>	0(0.0)	0(0.0)	2(33.3)	2(16.7)
<i>Malassezia species</i>	1(33.3)	1(33.3)	3(50.0)	5(41.6)

TOTAL	3(25.0)	3(25.0)	6(50.0)	12(8.0)
-------	---------	---------	---------	---------

149

150 **Discussion and conclusion**

151 This present study assessed the efficacy of the sterilization and cleaning methods
 152 used in the barbing salons and the possibility of those equipments acting as vehicles of
 153 transmission of fungal pathogens from one customer to the other.

154 The barbers in the salons recruited for the study were of wide range of barbing
 155 experience with mean years of practice 6.4 ± 3.5 yrs. They could reasonably be reflective of
 156 overall attitude of the barbers in the locality. Barbers in this part of the world are barely
 157 educated and often primary or secondary school leavers. Becoming a barber usually
 158 requires a few months of apprenticeship; the actual length of time varies, and is often
 159 dependent on the agreement between the apprentice and his master; no form of
 160 certification or license is involved. We found out that majority of the barbers were aware that
 161 their clippers could transmit ringworm infection, but were ignorant about other items doing
 162 same.

163 In this study, all the barbers decontaminated their clippers with the application of
 164 methylated spirit followed by flaming for 2-5 minutes after every customer usage. Other
 165 barbing equipments were rarely decontaminated while some barbers washed the combs
 166 and brushes with detergents and 10% hypochlorites once or twice a month at room
 167 temperature.

168 The recovery rate of fungal pathogens may depict the infectivity rate in the locality or
 169 the inefficiency of the sterilization processes in the barbing salons. In this study
 170 *Trichophyton interdigitale* 12(30.0%) was the most common dermatophyte recovered from
 171 barbing equipments, followed by *Trichophyton rubrum* 9(22.5%). This is in agreement with

the study of Oyeka and Eze [19], Ayanbimpe *et al* [9] and Mbata and Nwajagu [14] who reported *Trichophyton interdigitale* and *Trichophyton rubrum* as the most predominant species causing tinea capitis in the South Eastern and North Central zones of Nigeria. However, our findings differ from the study of Oke *et al* [20] who reported *Microsporum audouinii* (28.0%) as the commonest dermatophyte species among their subjects in South western Nigeria. These differences may be due to variation in distribution of dermatophytes species in different geographical areas in the country.

In this study hair brushes were the most contaminated equipments 54.0% by fungal organisms while the least contaminated were the hair clippers 20.0%. This shows that the sterilization method of the clippers by the application of methylated spirit followed by flaming is more effective than washing with detergents and 10% hypochlorite which is applicable to the brushes and combs.

The 17.5% recovery rates of dermatophytes from clippers depict incomplete sterilization leading to mere reduction in microbial load. The location with the highest recovery rate was Malabor, University of Calabar male hostel. This may be due to high patronage caused by high population of students which leaves the barbers with little or no time for sterilization of equipments.

The presence of non-dermatophyte moulds on the scalp is becoming increasingly common. Adefemi *et al* [21] and Chepchirchir *et al* [22] attributed this to the presence of spores in the environment which enables them to be carried transiently on healthy skin. The non-dermatophyte moulds identified in the study were *Aspergillus flavus* 25.0%, *Aspergillus fumigatus* 16.7% and *Aspergillus niger* 16.7%. Oke *et al* [20] in their study among children in South Western Nigeria also isolated lower rates of *Aspergillus fumigatus* (8.4%), and *Aspergillus niger* (6.4%). This suggests the possibility of transmission of these moulds from improperly sterilized barbing equipment from one customer to the other.

The limitation of the study was the inability to obtain multiple samples from the barbing equipments after each customer usage for laboratory studies. This could have given an insight unto whether the non-Dermatophytic moulds were contaminants or pathogens. This limitation was because of the uncertainty of when each customer will return for another barbing.

Conclusion

Dermatophytes and non dermatophytes were recovered from barbing equipments which depicts that they could serve as vehicles for transmission of fungal infections. Although the clippers were sterilized by all Barbers, dermatophytes and few non dermatophytes were still recovered from them. This point to incomplete sterilization. Customers that cannot afford hair clippers should insist on longer sterilization exposure time while combs and brushes should be personalized.

Financial Disclosure: The authors declared no financial support.

References

1. Chanda SK & Khan KH. Sharing of Razor Blade in Salons and Risk of Spreading HIV in Bangedeth. *Clin. Dermatol.* 2004; 28:197-201.
2. Wazir MS, Mehmood S, Ahed A. & Jadoon HR. Awareness among barbers about health hazards associated with their profession. *J Ajub Med. Coll. Abbottabad.* 2008; 20: 35-38.
3. Kondo M, Nakano N & Shiraki Y. A Chinese-Japanese Boy with Black Dot Ringworm due to *Trichophyton violaceum*. *J Dermatol.* 2006; 33(3): 165-168
4. Moto JN, Maingi JM and Nyamache AK. Prevalence of *Tinea capitis* in school going children from Mathare, informal settlement in Nairobi, Kenya. *BMC Research Notes* 2015; 8:274 DOI: 10.1186/s13104-015-1240-7
5. Oguzkaya AM, Koc AN, Baykan Z & Buldu H. Prevalence of *Tinea capitis* in primary school children. *Int J Med Investig.* 2013; 1(2):104-110.
6. Abdel-Rahman SM, Herron J, Fallon-Friedlander S, Hauffe S, Horowitz A & Rivière GJ. Pharmacokinetics of terbinafine in young children treated for *Tinea capitis*. *Pediatr Infect Dis J* 2005; 24(10):886-891.
7. Anosike JC, Keke IR & Uwaezuoke JC. Prevalence and distribution of ringworm infection in primary schools in parts of Eastern Nigeria. *J Appl Sci Environ Manag.* 2005, 9:21-25.

8. Sidat MM, Correia D & Buene TP. Tinea capitis among children at one suburban primary school in the city of Maputo, Mozambique. *Rev Soc Brasl Med Trop J.* 2007; 40:473-475.
9. Elewski BE. "Tinea capitis: A current perspective". *Journal of the American Academy of Dermatology* 2000; 42: 1-20.
10. Ayanbimpe GM, Taghir H, Diya A & Wapwera S. "Tinea capitis among primary school children in some parts of central Nigeria," *Mycoses* 2008; 51(4): 336–340
11. Pai VV, Hanumanthayya K, Tophakhane RS, Nandihal NN & Kikkeri NS. Clinical study of Tinea capitis in Northern Karnataka: A three-year experience at a single institute. *Indian Dermatol Online J.* 2013; 4(1): 22–26. doi: [10.4103/2229-5178.105461](https://doi.org/10.4103/2229-5178.105461)
12. Sahin I, Oksuz S, Kaya D, Sencan I & Çetinkaya R. "Dermatophytes in the rural area of Duzce, Turkey". *Mycoses* 2004; 47(11-2): 470-474
13. Tosti A, Piraccini BM, Lorenzi S, Iorizzo M. Treatment of non dermatophyte mold and *Candida onychomycosis*. *Dermatol Clin* 2003; 21:491-497.
14. Reiss E, Shadomy HJ, Lyod GM. Dermatophytosis and dermatomycoses (Superficial cutaneous mycoses): In: *Fundamental medical mycology* 2012; 1st ed. John Wiley & Sons, Inc. Pp. 525-563.
15. Mbata CC & Nwajagu C. Dermatophytes and Other Fungi Associated with Hair-Scalp of Nursery and Primary School Children in Awka, Nigeria. *The International Journal of Microbiology.* 2007; 3:2.
16. Baraikio D, Ogba OM & Abia-Bassey LN. Second Hand Covered Shoes: Transmission route of fungal foot infections. *Mary Slessor Journal of Medicine* 2014; 13(1): 26-33.
17. Nengimoyo B, Ogba OM & Abia-Bassey LN. Dandruff Aetiology and the Effects of edible lipids on the growth of Isolates. *European Journal of Medical and Pharmaceutical Research.* 2016; 3(9), 71-76.
18. Ali H, Shahram J, Mansour B, Ali SA, Parviz M, Siamak MR. Identification of Different *Malassezia* Species Isolated from Skin of Healthy Dog Owners in Tabriz, Iran (2010-2011). *Advances in Environmental Biology*, 2011; 5(11): 3688.
19. Midgley, G. The lipophilic yeasts: state of the art and prospects. *Medical Mycology*, 2000; 38: 9-16.
20. Onyeka CA & Eze II. "Fungal skin infections among prison inmates in Abakaliki, Nigeria," *Mycoses* 2008; 51(1): 50–54
21. Oke OO, Onayemi O, Olasode OA, Omisore AG & Oninla OA. The Prevalence and Pattern of Superficial Fungal Infections among School Children in Ile-Ife, South-Western Nigeria. *Dermatology Research and Practice*; 2014: 7 <http://dx.doi.org/10.1155/2014/842917>
22. Adefemi SA, Odeigah LO & Alabi KM. Prevalence of dermatophytosis among primary school children in Oke-oyi community of Kwara state. *Nigerian Journal of Clinical Practice* 2011; 14(1): 23–28.
23. Chepchirchir A, Bii C & Ndinya-Achola JO. Dermatophyte infections in primary school children in Kibera slums of Nairobi. *East African Medical Journal* 2009; 86(2): 59–68.